
Overview

Useful For

Evaluation of pediatric bone marrow and peripheral blood specimens by fluorescence in situ hybridization probe analysis for classic rearrangements and chromosomal copy number changes associated with T-cell acute lymphoblastic leukemia in patients being considered for enrollment in Children's Oncology Group clinical trials and research protocols

Testing Algorithm

This test is only performed on specimens from pediatric patients being considered for enrollment in a Children's Oncology Group (COG) protocol.

The fluorescence in situ hybridization (FISH) panel includes testing for the following abnormalities using the FISH probes listed:

+9/9p-, CDKN2A/D9Z1

t(9;22) or ABL1 amplification, ABL1/BCR

11q23 rearrangement, MLL (KMT2A) break-apart

-17/17p-, TP53/D17Z1

t(5;14), TLX3/BCL11B fusion

7q34 rearrangement, TRB [break-apart](#)

14q11.2 rearrangement, TRAD break-apart

t(10;11), MLLT10/PICALM fusion

1p33 rearrangement, TAL1/STIL

When an MLL (KMT2A) rearrangement is identified, reflex testing will be performed to identify the translocation partner. Probes include identification of:

t(11;19)(q23;p13.3) MLL/MLLT1

t(6;11)(q27;q23) MLLT4(AFDN)/MLL

t(4;11)(q21;q23) AFF1/MLL

t(9;11)(p22;q23) MLLT3/MLL

t(10;11)(p12;q23) MLLT10/MLL

t(11;19)(q23;p13.1) MLL/ELL

When a TRAD rearrangement is identified, reflex testing will be performed to identify the translocation partner. Probes include identification of:

t(11;14)(p15;q11.2) LMO1/TRAD

t(8;14)(q24.1;q11.2) MYC/TRAD

t(10;14)(q24;q11.2) TLX1(HOX11)/TRAD

t(11;14)(p13;q11.2) LMO2/TRAD

When a TRB rearrangement is identified, reflex testing will be performed to identify the translocation partner. Probes include identification of:

t(7;10)(q34;q24) TRB/TLX1

t(7;11)(q34;p15) TRB/LMO1

t(7;11)(q34;p13) TRB/LMO2

t(6;7)(q23;q34) MYB/TRB

In the absence of *BCR/ABL1* fusion or apparent episomal *ABL1* amplification, when an extra *ABL1* signal is identified, reflex testing will be performed using the *ABL1* break-apart probe set to evaluate for the presence or absence of an *ABL1* rearrangement.

Reflex Tests

Test Id	Reporting Name	Available Separately	Always Performed
COGTB	Probe, Each Additional (COGTF)	No, (Bill Only)	No

Method Name

Fluorescence In Situ Hybridization (FISH)

NY State Available

Yes

Specimen**Specimen Type**

Varies

Ordering Guidance

This test is only performed on specimens from pediatric patients being considered for enrollment in a Children's Oncology Group (COG) protocol. [If this test is ordered and the laboratory is informed that the patient is not on a COG protocol, this test will be canceled and automatically reordered by the laboratory as TALPF / T-Cell Acute Lymphoblastic Leukemia/Lymphoma \(ALL\), FISH, Pediatric Varies.](#)

At follow-up, conventional cytogenetic studies (CHRBM / Chromosome Analysis, Hematologic Disorders, Bone Marrow) and targeted T-ALL FISH probes can be evaluated based on the abnormalities identified in the diagnostic study. Order TALMF / T-Cell Acute Lymphoblastic Leukemia/Lymphoma (ALL), Specified FISH, Varies and request specific probes or abnormalities.

If the patient clinically relapses, a conventional chromosome study is useful to identify cytogenetic changes in the neoplastic clone or the possible emergence of a new therapy-related myeloid clone.

[For patients with T-cell lymphoma, order TLPDF / T-Cell Lymphoma, Diagnostic FISH, Varies.](#)

For testing paraffin-embedded tissue samples from patients with T-cell lymphoblastic lymphoma, order TLBLF / T-Lymphoblastic Leukemia/Lymphoma, FISH, Tissue.

Additional Testing Requirements

At diagnosis, conventional cytogenetic studies (COGBM / Chromosome Analysis, Hematologic Disorders, Children's Oncology Group Enrollment Testing, Bone Marrow) and this panel should be performed. If there is limited specimen available, only this test will be performed.

Shipping Instructions

Advise Express Mail or equivalent if not on courier service.

Necessary Information

1. [A reason for testing, a flow cytometry and/or a bone marrow pathology report, and a Children's Oncology Group \(COG\) registration number and protocol number should be submitted with each specimen.](#) The laboratory will not reject testing if this information is not provided, but appropriate testing and interpretation may be compromised or delayed. If this information is not provided, an appropriate indication for testing may be entered by Mayo Clinic Laboratories.
2. If the patient has received an opposite sex bone marrow transplant, note this information on the request.

Specimen Required

Submit only 1 of the following specimens:

Preferred

Specimen Type: Bone marrow

Container/Tube:

Preferred: Yellow top (ACD)

Acceptable: Green top (heparin) or lavender top (EDTA)

Specimen Volume: 2 to 3 mL

Collection Instructions:

1. It is preferable to send the first aspirate from the bone marrow collection.
2. Invert several times to mix bone marrow.

Acceptable

Specimen Type: Blood

Container/Tube:

Preferred: Yellow top (ACD)

Acceptable: Green top (heparin) or lavender top (EDTA)

Specimen Volume: 6 mL

Collection Instructions: Invert several times to mix blood.

Forms

[If not ordering electronically, complete, print, and send a Children's Oncology Group Test Request \(T829\)](#) with the specimen.

Reject Due To

All specimens will be evaluated at Mayo Clinic Laboratories for test suitability.

Specimen Minimum Volume

Blood: 2 mL

Bone Marrow: 1 mL

Specimen Stability Information

Specimen Type	Temperature	Time	Special Container
Varies	Ambient (preferred)		
	Refrigerated		

Clinical & Interpretive**Clinical Information**

In the United States, the incidence of acute lymphoblastic leukemia (ALL) is roughly 6000 new cases per year (as of 2019). ALL accounts for approximately 70% of all childhood leukemia cases (ages 0 to 19 years), making it the most common type of childhood cancer.

Approximately 85% of pediatric cases of ALL are B-cell lineage (B-ALL) and 15% are T-cell lineage (T-ALL). T-ALL is more common in adolescents than younger children and accounts for 25% of adult ALL. When occurring as a primary lymphoblastic lymphoma (LBL), approximately 90% are T-cell lineage versus only 10% B-cell lineage. T-LBL often present as a mediastinal mass in younger patients with or without concurrent bone marrow involvement.

Specific genetic abnormalities are identified in the majority of cases of T-ALL, although many of the classic abnormalities are "cryptic" by conventional chromosome studies and must be identified by fluorescence in situ hybridization (FISH) studies. Each of the genetic subgroups are important to detect and can be critical prognostic markers. One predictive marker, amplification of the *ABL1* gene region, has been identified in 5% of T-ALL, and these patients may be responsive to targeted tyrosine kinase inhibitors.

A combination of cytogenetic and FISH testing is currently recommended in all pediatric and adult patients to characterize the T-ALL clone for the prognostic genetic subgroups. A summary of the characteristic chromosome abnormalities identified in T-ALL are listed in the following table.

Table. Common Chromosome Abnormalities in T-cell Acute Lymphoblastic Leukemia

Cytogenetic change	Genes involved
del(1p33)	<i>TAL1/STIL</i>
t(5;14)(q35;q32)	<i>TLX3/BCL11B</i>
t(10;11)(p12;q14)	<i>MLLT10/PICALM</i>
Episomal amplification	<i>ABL1</i>
del(9p)	<i>CDKN2A(p16)</i>
t(11q23;var)	<i>MLL(KMT2A)</i>
t(4;11)(q21;q23)	<i>AFF1/MLL(KMT2A)</i>
t(6;11)(q27;q23)	<i>MLLT4(AFDN)/MLL(KMT2A)</i>
t(9;11)(p22;q23)	<i>MLLT3/MLL(KMT2A)</i>
t(10;11)(p12;q23)	<i>MLLT10)/MLL(KMT2A)</i>
t(11;19)(q23;p13.1)	<i>MLL(KMT2A)/ELL</i>
t(11;19)(q23;p13.3)	<i>MLL(KMT2A)/MLLT1</i>
t(7q34;var)	<i>TRB</i>
t(6;7)(q23;q34)	<i>MYB/TRB</i>
t(7;10)(q34;q24)	<i>TRB/TLX1</i>
t(7;11)(q34;p15)	<i>TRB/LMO1</i>
t(7;11)(q34;p13)	<i>TRB/LMO2</i>
t(14q11.2;var)	<i>TRAD</i>
t(8;14)(q24.1;q11.2)	<i>MYC/TRAD</i>
t(10;14)(q24;q11.2)	<i>TLX1/TRAD</i>
t(11;14)(p15;q11.2)	<i>LMO1/TRAD</i>
t(11;14)(p13;q11.2)	<i>LMO2/TRAD</i>
del(17p)	<i>TP53</i>

Reference Values

An interpretive report will be provided.

Interpretation

[A neoplastic clone is detected when the percent of cells with an abnormality exceeds the normal reference range for any given probe.](#)

The absence of an abnormal clone does not rule out the presence of neoplastic disorder.

Cautions

This test is not approved by the US Food and Drug Administration, and it is best used as an adjunct to existing clinical and pathologic information.

Fluorescence in situ hybridization (FISH) is not a substitute for conventional chromosome studies because the latter detects many chromosome abnormalities associated with other hematological disorders that would be missed by this FISH panel test.

Bone marrow is the preferred specimen type for this FISH test. If bone marrow is not available, a blood specimen may be used if there are malignant cells in the blood specimen (as verified by a hematopathologist).

Supportive Data

Each probe was independently tested and verified on unstimulated peripheral blood and bone marrow specimens. Normal cutoffs were calculated based on the results of 25 normal specimens. Each probe set was evaluated to confirm the probe set detected the abnormality it was designed to detect.

Clinical Reference

1. Swerdlow SH, Campo E, Harris NL, et al, eds: WHO Classification of Tumours of Haematopoietic and Lymphoid Tissues. IARC Press; 2017
2. Gesk S, Martin-Subero JI, Harder L, et al: Molecular cytogenetic detection of chromosomal breakpoints in T-cell receptor gene loci. *Leukemia*. 2003;17:738-745
3. Chin M, Mugishima H, Takamura M, et al: Hemophagocytic syndrome and hepatosplenic (gamma)(delta) T-cell lymphoma with isochromosome 7q and 8 trisomy. *J Pediatr Hematol Oncol*. 2004;26(6):375-378
4. Graux C, Cools J, Michaux L, et al: Cytogenetics and molecular genetics of T-cell acute lymphoblastic leukemia: from thymocyte to lymphoblast. *Leukemia*. 2006;20:1496-1510
5. Liu Y, Easton J, Shao Y, et al: The genomic landscape of pediatric and young adult T-lineage acute lymphoblastic leukemia. *Nat Genet*. 2017;49(8):1211-1218

Performance

Method Description

This test is performed using commercially available and laboratory-developed probes. Deletion of the *CDKN2A* locus on chromosome 9 and *TP53* on chromosome 17 are detected using enumeration strategy probes. Rearrangements involving *TAL1/STIL*, *TRB*, *MLL (KMT2A)*, and *TRAD* are detected using dual-color break-apart (BAP) strategy probes. Dual-color, dual-fusion fluorescence in situ hybridization (D-FISH) strategy probe sets are used to detect t(5;14), t(9;22), t(10;11), and in reflex testing when rearrangements of *MLL*, *TRB*, or *TRAD* genes are detected. Amplification of the *ABL1* (9q34) is detected using a D-FISH probe strategy. For enumeration and BAP strategy probe sets, 100 interphase nuclei are scored; 200 interphase nuclei are scored when D-FISH probes are used. All results are expressed as the percent abnormal nuclei. (Unpublished Mayo method)

PDF Report

No

Specimen Retention Time

4 weeks

Performing Laboratory Location

Rochester

Fees & Codes
Test Classification

This test was developed, and its performance characteristics determined by Mayo Clinic in a manner consistent with CLIA requirements. This test has not been cleared or approved by the US Food and Drug Administration.

CPT Code Information

88271 x2, 88275 x1, 88291 x1- FISH Probe, Analysis, Interpretation; 1 probe set

88271 x2, 88275x1 – FISH Probe, Analysis; each additional probe set (if appropriate)

LOINC® Information

Test ID	Test Order Name	Order LOINC Value
COGTF	COG, ALL (T-cell), FISH	In Process

Result ID	Reporting Name	LOINC®
602286	Result Summary	50397-9
602287	Interpretation	69965-2
602288	Result Table	93356-4

602289	Result	62356-1
GC016	Reason for Referral	42349-1
GC017	Specimen	31208-2
602291	Source	31208-2
602292	Method	85069-3
602293	Additional Information	48767-8
602294	Disclaimer	62364-5
602295	Released By	18771-6