

Overview

Method Name

Enzyme immunoassay (FEIA)

NY State Available

No

Specimen

Specimen Type

Serum

Specimen Required

Draw blood in a plain red-top tube(s), serum gel tube(s) is acceptable. Spin down and send 0.5 mL of serum refrigerated in a plastic vial.

Reject Due To

Hemolysis: NA

Thawing: Warm OK; Cold OK

Lipemia: NA

Icterus: NA

Other: NA

Specimen Minimum Volume

0.5 mL

Specimen Stability Information

Specimen Type	Temperature	Time	Special Container
Serum	Refrigerated (preferred)	28 days	
	Frozen	365 days	
	Ambient	7 days	

Clinical & Interpretive

Reference Values

Reference ranges have not been established for food-specific IgG tests. The clinical utility of food-specific IgG tests has not been established. These tests can be used in special clinical situations to select foods for evaluation by diet elimination and challenge in patients who have food-related complaints. It should be recognized that the presence of food-specific IgG alone cannot be taken as evidence of food allergy and only indicates immunologic sensitization by the food allergen in question. This test should only be ordered by physicians who recognize the limitations of the test.

Performance**PDF Report**

No

Performing Laboratory Location

Eurofins Viracor

Fees & Codes**Test Classification**

[This test was developed and its performance characteristics determined by Viracor Eurofins. It has not been cleared or approved by the U.S. Food and Drug Administration.](#)

CPT Code Information

86001