

Overview

Useful For

Confirming a diagnosis of bullous pemphigoid, cicatricial pemphigoid, pemphigoid gestationis and other variants of pemphigoid, all types of pemphigus, including paraneoplastic pemphigus (paraneoplastic multiorgan syndrome), dermatitis herpetiformis, linear IgA bullous dermatosis, chronic bullous disease of childhood, epidermolysis bullosa acquisita, porphyria cutanea tarda, bullous eruption of lupus erythematosus, and atypical or mixed forms of bullous disease, systemic lupus erythematosus, cutaneous lupus erythematosus, or other variants, vasculitis, lichen planus, and other inflammatory diseases

This test is **not useful** for diagnosis of malignancies involving the skin.

Testing Algorithm

For information see [Pathology Consultation Ordering Algorithm](#)

Special Instructions

- [Pathology Consultation Ordering Algorithm](#)

Method Name

Direct Immunofluorescence Assay (IFA)

NY State Available

Yes

Specimen

Specimen Type

Varies

Necessary Information

1. **Date of collection and location of biopsy site are required:** whether biopsy was obtained from sun-exposed versus unexposed skin, and whether it is from perilesional, involved, or uninvolved skin.
2. **Pathology Report/Clinical Notes are required** (include a brief history, pertinent laboratory results, suspected diagnosis, and reason for testing).

Specimen Required

Two or more biopsies from same site and sent in 1 specimen vial will be processed as 1 specimen. Two or more biopsies from different sites require separate specimen vials, however, they can be ordered together. Test performed on each site will be billed accordingly.

Transport Medium Method

Supplies: Michel's Transport Media for Immunofluorescent Testing on Tissue (T321)

Specimen Type: Tissue

Sources: Skin or 1 of the following mucosae: oral (oropharyngeal), nasal, genital, esophageal, conjunctival, laryngeal, or epiglottis

Container/Tube: Transport medium (Michel's, also called Zeus media)

Specimen Volume: 2-8 mm punch specimen, intact or bisected; excisional biopsy specimen intact or bisected

Collection Instructions:

1. Collect biopsy of uninvolved or involved skin. Refer to Recommended Biopsy Site Selection Based on Disease State below.
2. Immediately place specimen into a labeled vial of transport medium and seal tightly.

Snap-Frozen Method

Specimen Type: Tissue

Sources: Skin or 1 of the following mucosae: oral (oropharyngeal), nasal, genital, esophageal, conjunctival, laryngeal, or epiglottis

Container/Tube: Plastic vial

Specimen Volume: 2-8 mm punch specimen, intact or bisected; excisional biopsy specimen, intact or bisected

Collection Instructions:

1. Collect biopsy of uninvolved or involved skin. Refer to Recommended Biopsy Site Selection Based on Disease State below.
2. Immediately place specimen into liquid nitrogen and allow to freeze thoroughly (do not allow specimen to desiccate). If liquid nitrogen is not available, specimen may be frozen by placing it on a small square of aluminum foil on a block of dry ice. Liquid nitrogen is preferred.
3. Immediately wrap specimen carefully in aluminum foil. At no time should the specimen be allowed to thaw.
4. Place the wrapped specimen into the prelabeled plastic vial and seal tightly. Ship frozen.

Recommended Biopsy Site Selection Based on Disease State

1. **Pemphigus and pemphigoid groups** (including linear IgA bullous dermatosis and chronic bullous disease of childhood): Biopsy erythematous perilesional skin or mucosa. Avoid erosions, ulcers, and bullae while obtaining tissue adjacent to active lesions. Label as perilesional skin.
2. **Dermatitis herpetiformis:** Biopsy normal-appearing skin, 0.5-1 cm away from lesion. Label as perilesional skin.
3. **Lupus erythematosus:** Involved areas of skin such as erythematous or active borders are preferred biopsy sites to confirm the diagnosis of lupus erythematosus, either discoid or systemic. Label as involved skin. Avoid ulcers, old lesions, and facial lesions, if possible. Uninvolved, nonexposed skin is the preferred site to detect a lupus band as may be found in systemic lupus erythematosus. Should unexposed skin be desired, buttock or medial thigh is suggested. Label as uninvolved, nonexposed skin.
4. **Mixed connective tissue disease:** Biopsy as for lupus erythematosus except when sclerodermoid features are present. For sclerodermoid features, biopsy inflamed area. Label as involved or uninvolved, exposed or nonexposed skin.
5. **Vasculitis and urticaria:** The erythematous or active border of a new lesion is preferred. Avoid old lesions and ulcers. Label as involved skin. If appropriate, skin lesion is not present, diagnosis may sometimes be made from uninvolved skin.
6. **Porphyria cutanea tarda:** Biopsy involved skin. Avoid old lesions and ulcers. Label as involved skin.
7. **Lichen planus and lichenoid reactions:** Biopsy involved skin. Avoid old lesions and ulcers. Label as involved skin.

Specimen Minimum Volume

See Specimen Required

Reject Due To

Biopsy from lung, kidney, muscle, salivary gland, veins, synovial tissue, bronchial tissue, or bronchial lavage Biopsy in formalin fixation Frozen in alcohol Trumps media Glutaraldehyde	Reject
---	--------

Specimen Stability Information

Specimen Type	Temperature	Time	Special Container
Varies	Ambient (preferred)	30 days	
	Frozen	120 days	
	Refrigerated	30 days	

Clinical & Interpretive

Clinical Information

Skin or mucosal tissue from patients with autoimmune bullous diseases, connective tissue disease, vasculitis, lichen planus, and other inflammatory conditions often contains bound immunoglobulin, complement, or fibrinogen.

Biopsy specimens are examined for the presence of bound IgG, IgM, IgA, third component of complement (C3), fibrinogen, and IgG4.

Reference Values

An interpretive report will be provided.

Interpretation

A board-certified Dermatopathologist will review and interpret the test results in correlation with other clinical findings as provided.

Cautions

This test is an adjunctive test to be interpreted in the context of clinical information, histologic studies, and serologic studies as clinically indicated.

Clinical Reference

1. Jain S, Basavaraj V, Vimala MG: Utility of Direct Immunofluorescence Studies in Subclassification of Autoimmune Sub-Epidermal Bullous Diseases: A 2-Year Study in a Tertiary Care Hospital. *Turk Patoloji Derg.* 2016;32(2):91-98. doi: 10.5146/tjpath.2015.01345
2. Diercks GF, Pas HH, Jonkman MF: Immunofluorescence of Autoimmune Bullous Diseases. *Surg Pathol Clin.* 2017 Jun;10(2):505-512. doi: 10.1016/j.path.2017.01.011
3. Kershenovich R, Hodak E, Mimouni D: Diagnosis and classification of pemphigus and bullous pemphigoid. *Autoimmun Rev.* 2014 Apr-May;13(4-5):477-481. doi: 10.1016/j.autrev.2014.01.011
4. Buschman KE, Seraly M, Thong HY, Deng JS, Draviam RP, Abernethy JL: A predominant IgG4 subclass may be responsible for false-negative direct immunofluorescence in bullous pemphigoid. *J Cutan Pathol.* 2002 May;29(5):282-286. doi: 10.1034/j.1600-0560.2002.290504.x
5. Lamb PM, Patton T, Deng JS: The predominance of IgG4 in prodromal bullous pemphigoid. *Int J Dermatol.* 2008 Feb;47(2):150-153. doi: 10.1111/j.1365-4632.2008.03361.x

Performance**Method Description**

Frozen sections of biopsy specimens are brought to ambient temperature, air dried, washed with phosphate-buffered saline (PBS), and then layered with fluorescein isothiocyanate (FITC)-conjugated rabbit antihuman IgG, IgA, IgM, C3, fibrinogen, and IgG4. These slides are incubated in a moist chamber at ambient temperature. The sections are then washed with PBS, mounted in buffered glycerine, and viewed under a fluorescence microscope. (Mysorekar VV, Sumathy TK, Shyam Prasad AL: Role of direct immunofluorescence in dermatological disorders. *Indian Dermatol Online J.* 2015;6[3]:172-180. doi: 10.4103/2229-5178.156386)

PDF Report

No

Day(s) Performed

Monday through Friday

Report Available

2 to 3 days

Specimen Retention Time

Stained slides:14 days; Remaining biopsy tissue: 30 days

Performing Laboratory Location

Rochester

Fees & Codes

Fees

- Authorized users can sign in to [Test Prices](#) for detailed fee information.
- Clients without access to Test Prices can contact [Customer Service](#) 24 hours a day, seven days a week.
- Prospective clients should contact their account representative. For assistance, contact [Customer Service](#).

Test Classification

This test was developed using an analyte specific reagent. Its performance characteristics were determined by Mayo Clinic in a manner consistent with CLIA requirements. This test has not been cleared or approved by the US Food and Drug Administration.

CPT Code Information

Per biopsy site:

88346

88350 x 5

LOINC® Information

Test ID	Test Order Name	Order LOINC® Value
CIB	Cutaneous Direct IFA, Biopsy	In Process

Result ID	Test Result Name	Result LOINC® Value
71145	Interpretation	66121-5
71146	Participated in the Interpretation	No LOINC Needed
71147	Report electronically signed by	19139-5
71610	Addendum	35265-8
71855	Case Number	80398-1