

Overview

Useful For

Preferred screening test for detection of arsenic, cadmium, mercury and lead due to occupational exposure using random urine specimens

Testing Algorithm

If arsenic concentration is greater than or equal to 10 mcg/L, then speciation will be performed at an additional charge.

Special Instructions

- [Trace Metals Analysis Specimen Collection and Transport](#)

Profile Information

Test Id	Reporting Name	Available Separately	Always Performed
ASOU	Arsenic Occupational Exposure	Yes, (order ASUOE)	Yes
CDOU	Cadmium Occupational Exposure	Yes, (order CDUOE)	Yes
HGOU	Mercury Occupational Exposure	Yes, (order HGUOE)	Yes
PBOU	Lead Occupational Exposure	Yes, (order PBOUE)	Yes
CRETR	Creatinine, Random, U	No	Yes

Reflex Tests

Test Id	Reporting Name	Available Separately	Always Performed
SPAS	Arsenic Speciation, Random, U	Yes	No

Method Name

ASOU, CDOU, HGOU, PBOU: Inductively Coupled Plasma-Mass Spectrometry (ICP-MS)

CRETR: Enzymatic Colorimetric Assay

NY State Available

Yes

Specimen

Specimen Type

Urine

Specimen Required**Patient Preparation:**

-Patient should not eat seafood for a 48-hour period prior to start of collection.

-High concentrations of gadolinium and iodine are known to interfere with most metals tests. If either gadolinium- or iodine-containing contrast media has been administered, a specimen should not be collected for 96 hours.

Supplies: Urine Tubes, 10 mL (T068)**Collection Container/Tube:** Clean, plastic urine container with no metal cap or glued insert**Submission Container/Tube:** Plastic, 10-mL urine tube or clean, plastic aliquot container with no metal cap or glued insert**Specimen Volume:** 6 mL**Collection Instructions:**

1. Collect a random urine specimen.
2. See [Trace Metals Analysis Specimen Collection and Transport](#) in Special Instructions for complete instructions.

Reject Due To

All specimens will be evaluated at Mayo Clinic Laboratories for test suitability.

Specimen Minimum Volume

3 mL

Specimen Stability Information

Specimen Type	Temperature	Time	Special Container
Urine	Refrigerated (preferred)	7 days	
	Frozen	7 days	

Clinical & Interpretive**Clinical Information**

Arsenic (As), lead (Pb), cadmium (Cd), and mercury (Hg) are well-known toxins and toxic exposures are characterized by increased urinary excretion of these metals.

Arsenic is a naturally occurring element that is usually found in the environment combined with other elements such as oxygen, chlorine, and sulfur. Arsenic combined with these elements is called inorganic arsenic. Arsenic combined with carbon and hydrogen is referred to as organic arsenic. The organic forms (eg, arsenobetaine and arsenocholine) are relatively nontoxic, while the inorganic forms are toxic. The toxic inorganic forms are arsenite (As[3+]/As[III]) and arsenate (As[5+]/As[V]). Inorganic As(V) is readily reduced to inorganic As(III), which is then primarily broken down to the less toxic methylated metabolites, monomethylarsinic acid (MMA) and subsequently dimethylarsinic acid (DMA).

People are exposed to arsenic by eating food, drinking water, or breathing air. Of these, food is usually the largest source of arsenic. The predominant dietary source of arsenic is seafood, followed by rice/rice cereal, mushrooms, and poultry. While seafood contains the greatest amounts of arsenic, from fish and shellfish, this is mostly in an organic form of arsenic called arsenobetaine, which is much less harmful. Some seaweed may contain arsenic in the inorganic form, which is more toxic. In the United States, some areas also contain high natural levels of arsenic in rock, which can lead to elevated levels in the soil and drinking water. Occupational (eg, copper or lead smelting, wood treating, or pesticide application) exposure is another source where people may be introduced to elevated levels of arsenic. Lastly, hazardous waste sites may contain large quantities of arsenic and, if not disposed of properly, may get into the surrounding water, air, or soil.

A wide range of signs and symptoms may be seen in acute arsenic poisoning including headache, nausea, vomiting, diarrhea, abdominal pain, hypotension, fever, hemolysis, seizures, and mental status changes. Symptoms of chronic poisoning, also called arseniasis, are mostly insidious and nonspecific. The gastrointestinal tract, skin, and central nervous system are usually involved. Nausea, epigastric pain, colic abdominal pain, diarrhea, and paresthesias of the hands and feet can also occur.

Since arsenic is excreted predominantly by glomerular filtration, measurement of arsenic in urine is the most reliable means of detecting arsenic exposures within the last several days.

Lead toxicity primarily affects the gastrointestinal, neurologic, and hematopoietic systems.

Chronic exposure to cadmium causes accumulated renal damage.

The correlation between the levels of mercury (Hg) excretion in the urine and the clinical symptoms is considered poor.

Reference Values

ARSENIC:

Biological Exposure Indices (BEI): <35 mcg/L at end of work week

CADMIUM:

BEI: <5.0 mcg/g creatinine

MERCURY:

BEI: <35 mcg/g creatinine

LEAD:

BEI: <150 mcg/g creatinine

Interpretation**Arsenic:**

Mayo Clinic uses the American Conference of Governmental Industrial Hygienists (ACGIH) biological exposure index (BEI) as the reference value. The BEI is the sum of all the toxic species (inorganic arsenic plus methylated arsenic metabolites).

Physiologically, arsenic exists in a number of toxic and nontoxic forms. The total arsenic concentration reflects all the arsenic present in the sample regardless of species (eg, inorganic vs. methylated vs. organic arsenic). The measurement of urinary total arsenic levels is generally accepted as the most reliable indicator of recent arsenic exposure. However, if the total urine arsenic concentration is elevated, arsenic speciation must be performed to identify if it is the toxic forms (eg, inorganic and methylated arsenic forms) or the relatively nontoxic organic forms (eg, arsenobetaine and arsenocholine).

The inorganic toxic forms of arsenic (eg, As[III] and As[V]) are found in the urine shortly after ingestion, whereas the less toxic methylated forms, monomethylarsinic acid (MMA) and dimethylarsinic acid (DMA)) are the species that predominate longer than 24 hours after ingestion. In general, urinary As(III) and As(V) concentrations peak in the urine at approximately 10 hours and return to normal 20 to 30 hours after ingestion. Urinary MMA and DMA concentrations normally peak at approximately 40 to 60 hours and return to baseline 6 to 20 days after ingestion.

This test can determine if a patient has been exposed to above-average levels of arsenic. It cannot predict whether the arsenic levels in their body will affect their health.

Cadmium:

In chronic cadmium exposure, the kidneys are the primary target organ. Urine concentrations of cadmium can be useful to assess long-term exposure and determine cadmium body burden.

Cadmium excretion above 3.0 mcg/g creatinine indicates significant exposure to cadmium. For occupational testing, the Occupational Safety and Health Administration (OSHA) cadmium standard is below 3.0 mcg/g creatinine and the BEI is 5 mcg/g creatinine.

Mercury:

It had always been thought that urine was a more appropriate marker of inorganic mercury, because organic mercury represented only a small fraction of urinary mercury. Based on possible demethylation of methylmercury within the body, urine may represent a mixture of dietary methylmercury and inorganic mercury. Seafood consumption can contribute to urinary mercury levels (up to 30%),⁽¹⁾ which is consistent with the suggestion that due to demethylation processes in the human body, a certain proportion of urinary mercury can originate from dietary consumption of fish/seafood.⁽²⁾

Lead:

Measurements of urinary lead levels have been used to assess lead exposure. However, like lead blood, urinary lead excretion mainly reflects recent exposure and thus shares many of the same limitations for assessing Pb body burden or long-term exposure.^(3,4)

Urinary lead concentration increases exponentially with blood lead and can exhibit relatively high intra-individual variability, even at similar blood lead concentrations.^(5,6)

Cautions

Consumption of seafood before collection of a urine specimen for arsenic testing is likely to result in a report of an elevated concentration of arsenic found in the urine, which can be clinically misleading.

Nitric acid cannot be added to either the collection or aliquot container. Nitrate interferes with the extraction procedure that would need to take place in the event of a positive arsenic result.

Clinical Reference

1. Snoj Tratniid J, Falnoga I, Mazej D, et al: Results of the first national human biomonitoring in Slovenia: Trace elements in men and lactating women, predictors of exposure and reference values. *Int J Hyg Environ Health*. 2019;222(3):563-582
2. Sherman LS, Blum JD, Franzblau A, Basu N: New insights into biomarkers of human mercury exposure using naturally occurring mercury stable isotopes. *Environ Sci Technol*. 2013;47(7):3403-3409
3. Sakai T: Biomarkers of lead exposure. *Ind Health*. 2000;38(2):127-142
4. Skerfving S: Biological monitoring of exposure to inorganic lead. In: Clarkson TW, Friberg L, Nordberg GF, Sager PR, eds. *Biological Monitoring of Toxic Metals*. Rochester Series on Environmental Toxicity. Springer;1988:169-197
5. Gulson BL, Jameson CW, Mahaffey KR, et al: Relationships of lead in breast milk to lead in blood, urine, and diet of the infant and mother. *Environ Health Perspect*. 1998;106(10):667-674
6. Skerfving S, Ahlgren L, Christoffersson JO: Metabolism of inorganic lead in man. *Nutr Res* 1985;Suppl 1:601-607
7. Fillol CC, Dor F, Labat L, et al: Urinary arsenic concentrations and speciation in residents living in an area with naturally contaminated soils. *Sci Total Environ*. 2010 Feb 1;408(5):1190-1194
8. Caldwell KL, Jones RL, Verdon CP, Jarrett JM, Caudill SP, Osterloh JD: Levels of urinary total and speciated arsenic in the US population: National Health and Nutrition Examination Survey 2003-2004. *J Expo Sci Environ Epidemiol*. 2009 Jan;19(1):59-68
9. Lee R, Middleton D, Caldwell K, et al: A review of events that expose children to elemental mercury in the United States. *Environ Health Perspect*. 2009 Jun;117(6):871-878
10. Kosnett MJ, Wedeen RP, Rotherberg SJ, et al: Recommendations for medical management of adult lead exposure. *Environ Health Perspect*. 2007;115:463-471
11. de Burbane C, Buchet JP, Leroyer A, et al: Renal and neurologic effects of cadmium, lead, mercury, and arsenic in children: evidence of early effects and multiple interactions at environmental exposure levels. *Environ Health Perspect*. 2006;114:584-590
12. Agency for Toxic Substances and Disease Registry: Toxicological profile for arsenic. US Department of Health and Human Services. 2007. Available at www.atsdr.cdc.gov/ToxProfiles/tp2.pdf
13. Strathmann FG, Blum LM: Toxic elements. In: Rifai N, Horwath AR., Wittwer CT, eds. *Tietz Textbook of Clinical Chemistry and Molecular Diagnostics*. 6th ed. Elsevier; 2018:chap 42

Performance**Method Description**

Arsenic (As), cadmium (Cd), mercury (Hg), and lead (Pb) in urine are analyzed by inductively coupled plasma-mass spectrometry (ICP-MS) in kinetic energy discrimination (KED) mode using gallium (Ga), rhodium (Rh), and iridium (Ir) as internal standards and a 5% nitric acid salt matrix calibration.(Unpublished Mayo method)

PDF Report

No

Specimen Retention Time

14 days

Performing Laboratory Location

Rochester

Fees & Codes

Test Classification

This test was developed, and its performance characteristics determined by Mayo Clinic in a manner consistent with CLIA requirements. This test has not been cleared or approved by the US Food and Drug Administration.

CPT Code Information

82175

82300

83825

83655

82570

LOINC® Information

Test ID	Test Order Name	Order LOINC Value
HMUOE	Heavy Metal Occ Exp w/Reflex, U	29589-9

Result ID	Reporting Name	LOINC®
CRETR	Creatinine, Random, U	2161-8
608890	Arsenic Occupational Exposure	13463-5
608891	Total Arsenic Concentration	5586-3
608892	Cadmium Occupational Exposure	13471-8

608893	Mercury Occupational Exposure	13465-0
608894	Lead Occupational Exposure	13466-8