

## Overview

### Useful For

Detecting mercury toxicity due to occupational exposure in random urine specimens

### Special Instructions

- [Trace Metals Analysis Specimen Collection and Transport](#)

### Method Name

Only orderable as part of a profile. For more information see:

-HGUOE/ Mercury Occupational Exposure, Random, Urine

-HMUOE / Heavy Metal Occupational Exposure, with Reflex, Random, Urine

Triple Quadrupole Inductively Coupled Plasma-Mass Spectrometry (ICP-MS/MS)

### NY State Available

Yes

## Specimen

### Specimen Type

Urine

### Specimen Required

Only orderable as part of a profile. For more information see:

-HGUOE/ Mercury Occupational Exposure, Random, Urine

-HMUOE / Heavy Metal Occupational Exposure, with Reflex, Random, Urine

### Specimen Minimum Volume

3 mL

### Specimen Stability Information

Specimen Type	Temperature	Time	Special Container
Urine	Refrigerated (preferred)	7 days	
	Frozen	7 days	

## Clinical & Interpretive

---

**Clinical Information**

The correlation between the levels of mercury (Hg) excretion in the urine and the clinical symptoms is considered poor. However, urinary Hg is the most reliable way to assess exposure to inorganic Hg.

For more information, see HG / Mercury, Blood.

**Reference Values**

Only orderable as part of a profile. For more information see:

-HGUE/ Mercury Occupational Exposure, Random, Urine

-HMUE / Heavy Metal Occupational Exposure, with Reflex, Random, Urine

**Cautions**

To avoid contamination by dust, specimen should be collected away from the site of suspected exposure.

**Clinical Reference**

1. Snoj Tratniid J, Falnoga I, Mazej D, et al: Results of the first national human biomonitoring in Slovenia: Trace elements in men and lactating women, predictors of exposure and reference values. *Int J Hyg Environ Health.* 2019;222(3):563-582
2. Sherman LS, Blum JD, Franzblau A, Basu N: New insights into biomarkers of human mercury exposure using naturally occurring mercury stable isotopes. *Environ Sci Technol.* 2013;47(7):3403-3409
3. Lee R, Middleton D, Caldwell K, et al: A review of events that expose children to elemental mercury in the United States. *Environ Health Perspect.* 2009 Jun;117(6):871-878
4. Bjorkman L, Lundekvam BF, Laegreid T, et al: Mercury in human brain, blood, muscle and toenails in relation to exposure: an autopsy study. *Environ Health.* 2007 Oct 11;6:30

**Performance****Method Description**

Mercury (Hg) is analyzed by inductively coupled plasma-mass spectrometry (ICP-MS) in MS/MS mode using oxygen as a reactive gas to mass shift tungsten away from the mass of interest. Internal standard used is iridium for Hg. A salt matrix calibration is used.(Unpublished Mayo method)

**PDF Report**

No

**Day(s) Performed**

Monday through Friday

**Report Available**

1 day

**Specimen Retention Time**

14 days

---

**Performing Laboratory Location**

Rochester

**Fees & Codes****Fees**

- Authorized users can sign in to [Test Prices](#) for detailed fee information.
- Clients without access to Test Prices can contact [Customer Service](#) 24 hours a day, seven days a week.
- Prospective clients should contact their Regional Manager. For assistance, contact [Customer Service](#).

**Test Classification**

This test was developed, and its performance characteristics determined by Mayo Clinic in a manner consistent with CLIA requirements. This test has not been cleared or approved by the US Food and Drug Administration.

**LOINC® Information**

Test ID	Test Order Name	Order LOINC® Value
HGOU	Mercury Occupational Exposure	13465-0

  

Result ID	Test Result Name	Result LOINC® Value
608893	Mercury Occupational Exposure	13465-0