

Overview

Useful For

Aiding in the diagnosis of semaphorin 3B-associated membranous nephropathy

Testing Algorithm

For the initial technical component only immunohistochemical (IHC) stain performed, the appropriate bill-only test ID will be reflexed and charged (IHTOI). For each additional technical component only IHC stain performed, an additional bill-only test ID will be reflexed and charged (IHTOA).

Method Name

Immunohistochemistry (IHC)

NY State Available

Yes

Specimen

Specimen Type

TECHONLY

Ordering Guidance

This test includes only technical performance of the stain (no pathologist interpretation is performed). If diagnostic consultation by a pathologist is required, order PATHC / Pathology Consultation.

Shipping Instructions

Attach the green pathology address label and the pink Immunostain Technical Only label included in the kit to the outside of the transport container.

Specimen Required

Supplies: Immunostain Technical Only Envelope (T693)

Specimen Type: Tissue

Container/Tube: Immunostain Technical Only Envelope

Preferred: 2 unstained positively charged glass slide (25- x 75- x 1-mm) per test ordered; sections 4-microns thick

Acceptable: Formalin-fixed, paraffin-embedded (FFPE) tissue block

Digital Image Access

Mayo Clinic is preparing to migrate to a new slide scanning solution and will need to temporarily suspend digital imaging for tech-only IHC/ISH orders received after June 17. We will send a follow-up notification by September 1st indicating when digital imaging resumes. We appreciate your patience as we improve our digital imaging solution.

Forms

If not ordering electronically, complete, print, and send a [Immunohistochemical \(IHC\)/In Situ Hybridization \(ISH\) Stains Request](#) (T763) with the specimen.

Reject Due To

Wet/frozen tissue Cytology smears Nonformalin fixed tissue Nonparaffin embedded tissue Noncharged slides ProbeOn slides	Reject
--	--------

Specimen Stability Information

Specimen Type	Temperature	Time	Special Container
TECHONLY	Ambient (preferred)	0 hours	
	Refrigerated	0 hours	

Clinical & Interpretive

Clinical Information

Primary membranous nephropathy (MN) is an autoimmune disease of the kidney where antibodies target an antigen in the glomerular basement membrane resulting in kidney damage or failure. In up to 75% of primary membranous nephropathy cases PLA2R (phospholipase A2 receptor) and THSD7A (thrombospondin type-1 domain-containing 7A) are the target antigens. SEMA3B (semaphorin 3B)-associated MN is a very rare disease that accounts for only 1% to 3% of all MN specimens. SEMA3B-associated MN is observed mostly in pediatric patients and young adults.

Interpretation

This test does not include pathologist interpretation, only technical performance of the stain. If interpretation is required, order PATHC / Pathology Consultation for a full diagnostic evaluation or second opinion of the case.

The positive and negative controls are verified as showing appropriate immunoreactivity and documentation is retained at Mayo Clinic Rochester. If a control tissue is not included on the slide, a scanned image of the relevant quality control tissue is available upon request; call 855-516-8404.

Interpretation of this test should be performed in the context of the patient's clinical history and other diagnostic tests by a qualified pathologist.

Cautions

Age of a cut paraffin section can affect immunoreactivity. Stability thresholds vary widely among published literature and are antigen dependent. Best practice is for paraffin sections to be cut within 6 weeks.

Clinical Reference

1. Sethi S, Debiec H, Madden B, et al: [Semaphorin 3B-associated membranous nephropathy is a distinct type of disease predominantly present in pediatric patients. Kidney Int. 2020 Nov;98\(5\):1253-1264](#)
2. Sethi S: Membranous nephropathy: a single disease or a pattern of injury resulting from different diseases. Clin Kidney J. 2021 Mar 26;14(10):2166-2169
3. Sethi S: New 'antigens' in membranous nephropathy. J Am Soc Nephrol. 2021 Feb;32(2):268-278

Performance

Method Description

Immunohistochemistry on sections of paraffin-embedded tissue.(Unpublished Mayo method)

PDF Report

No

Day(s) Performed

Monday through Friday

Report Available

1 to 3 days

Specimen Retention Time

Until staining is complete.

Performing Laboratory Location

Rochester

Fees & Codes

Fees

- Authorized users can sign in to [Test Prices](#) for detailed fee information.
- Clients without access to Test Prices can contact [Customer Service](#) 24 hours a day, seven days a week.
- Prospective clients should contact their Regional Manager. For assistance, contact [Customer Service](#).

Test Classification

This test has been cleared, approved, or is exempt by the US Food and Drug Administration and is used per manufacturer's instructions. Performance characteristics were verified by Mayo Clinic in a manner consistent with CLIA requirements.

CPT Code Information

88342-TC, primary

88341-TC, if additional IHC

LOINC® Information

Test ID	Test Order Name	Order LOINC® Value
SEMA3	SEMA3B IHC, Tech Only	Order only;no result

Result ID	Test Result Name	Result LOINC® Value
617056	SEMA3B IHC, Tech Only	Bill only; no result