

## Overview

### Useful For

[Rapid testing for Powassan virus RNA \(lineage 1 and lineage 2\) using serum specimens](#)

An adjunctive test to serology for detection of early Powassan virus infection (ie, first few days after symptom onset). The Powassan virus polymerase chain reaction result may remain positive for a longer time in urine than in blood, serum, and cerebrospinal fluid (7 days or more).

This assay **should not be used for** screening asymptomatic individuals and should only be used to test patients with signs and symptoms of Powassan virus disease.

### Testing Algorithm

[For information, see Meningitis/Encephalitis Panel Algorithm](#)

### Method Name

Real-Time Polymerase Chain Reaction (PCR)

### NY State Available

Yes

## Specimen

### Specimen Type

Serum

### Additional Testing Requirements

This test is designed to be used during the first 7 days after symptom onset and should be ordered in conjunction with POWV / Powassan Virus, IgM, Enzyme-Linked Immunosorbent Assay, Serum.

### Specimen Required

#### Collection Container/Tube:

**Preferred:** 3.5-mL SST

**Acceptable:** Red top

**Submission Container:** Plastic vial

**Specimen Volume:** 0.5 mL

#### Collection Instructions:

1. Centrifuge and aliquot the serum into a plastic vial within 6 hours of collection.
2. Label specimen as serum

## Forms

If not ordering electronically, complete, print, and send a [Microbiology Test Request](#) (T732) with the specimen.

## Specimen Minimum Volume

0.3 mL

## Reject Due To

Gross hemolysis	Reject
Heat-inactivated specimen	Reject

## Specimen Stability Information

Specimen Type	Temperature	Time	Special Container
Serum	Refrigerated (preferred)	7 days	
	Frozen	7 days	

## Clinical & Interpretive

### Clinical Information

Powassan virus (POWV) is an emerging tick-borne virus, harbored by *Ixodes* species ticks, which are the same ticks that transmit Lyme disease (*Borrelia* spp.), *Babesia* spp., and *Anaplasma phagocytophilum*, among other pathogens. POWV is a member of the *Flavivirus* genus, which includes other arthropod-borne viruses (arboviruses) such as West Nile virus (WNV) and St. Louis encephalitis virus. Two lineages of POWV have been identified, sharing approximately 94% amino acid sequence identity, including Lineage 1, which is the prototypical POWV lineage transmitted by *Ixodes marxi* and *Ixodes cookei*, and Lineage 2, which includes deer tick virus and is transmitted by *Ixodes scapularis*. POWV is maintained in the environment in groundhogs, skunks, squirrels, and white footed mice. Unlike other tick-borne pathogens, following tick attachment to a host, POWV can be transmitted in as little as 15 minutes.

Following infection, the incubation period can last anywhere from 4 to 14 days, after which approximately 66% of patients will remain asymptomatic. Symptomatic patients may present with a nonspecific influenza-like illness, including high fever, fatigue, malaise, and myalgia. Approximately 30% of symptomatic patients will progress to develop neurologic manifestations, most commonly encephalitis. While some patients may recover, over 50% of individuals will have persistent neurologic sequelae. POWV has been associated with an overall mortality rate of 10%.

Although there is no targeted antiviral therapy and treatment is entirely supportive care, diagnosis is important for a number of reasons, including the ability to discontinue empiric antibiotics and to provide prognostic information for patients and families.

While limited data exist for POWV real-time reverse transcription polymerase chain reaction (RT-PCR) detection, data from testing for the related flavivirus, WNV, suggests that blood may be a more sensitive source than serum for

---

detecting viral RNA. Similarly, data from WNV and other flaviviruses suggest that viral RNA may be detected in urine for a longer period of time than in blood or serum. POWV RNA may be detected from cerebrospinal fluid in cases of neuroinvasive disease.

[Powassan infections are often diagnosed based on a patient's symptoms and exposure risk in conjunction with molecular and serologic testing. The use of RT-PCR can provide a rapid laboratory confirmation of POWV RNA early in infection, particularly during the first 7 days of illness when serologic testing is typically negative. After 7 days, POWV RT-PCR is less sensitive and serologic testing is the preferred diagnostic method. It is common for RT-PCR and serology to be used together for diagnosis of early infection, as they are complementary methods.](#)

### Reference Values

Negative

### Interpretation

A positive result indicates the presence of Powassan virus RNA and is consistent with early Powassan virus infection.

### Cautions

The sensitivity of the assay is dependent upon the time of illness onset in which the specimen is collected. Polymerase chain reaction testing has the greatest utility when used within the first few days of symptom onset.

A negative test does not exclude infection with Powassan virus. Therefore, the results obtained should be used in conjunction with clinical findings and serologic test results to make an accurate diagnosis.

This assay detects both viable and nonviable virus. Test performance depends on viral load in the specimen and may not correlate with cell culture performed on the same specimen.

### Supportive Data

[The following validation data supports the use of this assay for clinical testing.](#)

#### Accuracy/Diagnostic Sensitivity and Specificity:

Accuracy studies were performed by testing negative clinical specimens with whole viral genomic RNA for lineages 1 and 2 near the limit of detection (LOD) and yielded greater than or equal to 97% sensitivity and specificity.

#### Analytical Sensitivity/Limit of Detection:

The lower LOD of this assay is 1 to 52 target copies/mcL of RNA extract for EDTA whole blood and serum, 0.1 to 5.2 copies/mcL for urine, and 0.1 to 52 copies/mcL for cerebrospinal fluid.

#### Precision:

Inter-assay and intra-assay precisions are 100%.

#### Specificity:

A panel of 15 organisms that can be found in the specimen types acceptable for this assay, as well as closely-related viruses (eg, Dengue types 1-4, Japanese encephalitis virus, hepatitis E virus, Murray Valley encephalitis virus, St. Louis encephalitis virus, tick-borne encephalitis virus, yellow fever virus, Zika virus) and those that can cause a similar clinical syndrome were tested by this assay. No cross-reacting positive results were noted.

**Reportable Range:**

This is a qualitative assay, and the results are reported as either negative or positive for targeted Powassan virus.

**Clinical Reference**

1. Centers for Disease Control and Prevention (CDC), National Center for Emerging and Zoonotic Infectious Diseases, Division of Vector-Borne Diseases: Powassan Virus. For Healthcare Providers: Diagnostic testing. CDC; Updated June 23, 2021. Accessed April 26, 2022. Available at [www.cdc.gov/powassan/diagnostic-testing.html](http://www.cdc.gov/powassan/diagnostic-testing.html)
2. Piantadosi A, Rubin DB, McQuillen DP, et al: Emerging cases of Powassan virus encephalitis in New England: clinical presentation, imaging, and review of the literature. 2016 Clin Infect Dis. 2016 Mar 15;62(6):707-713

**Performance****Method Description**

[For this real-time reverse-transcription laboratory-developed polymerase chain reaction \(PCR\) assay, viral nucleic acid is extracted from specimens, followed by amplification and detection on the Roche LightCycler 480 instrument. This PCR assay has been optimized to detect a target sequence in the non-structural protein. Primers amplify a 179- base pair product. Powassan virus genomic RNA is first transcribed to complementary DNA \(cDNA\) by reverse transcriptase, followed by amplification of the cDNA product. The LightCycler 480 is an automated instrument that amplifies and monitors the development of target nucleic acid \(amplicon\) after each cycle of PCR. The detection of amplicon is based on fluorescence resonance energy transfer, which utilizes a hybridization probe with a donor fluorophore, fluorescein, at the 3' end and a second hybridization probe with an acceptor fluorophore, LC-Red 610, at the 5' end. When the target amplicon is present, the LC-Red 610 emits a measurable and quantifiable light signal at a specific wavelength. Presence of the specific organism nucleic acid may be confirmed by performing a melting curve analysis of the amplicon. Using features of the melting curve analysis, the assay primers and specific hybridization probes are able to detect Powassan virus lineage 1 and lineage 2.](#)(Unpublished Mayo method)

**PDF Report**

No

**Day(s) Performed**

Monday through Friday

**Report Available**

Same day/1 to 5 days

**Specimen Retention Time**

1 week

**Performing Laboratory Location**

Rochester

## Fees & Codes

### Fees

- Authorized users can sign in to [Test Prices](#) for detailed fee information.
- Clients without access to Test Prices can contact [Customer Service](#) 24 hours a day, seven days a week.
- Prospective clients should contact their account representative. For assistance, contact [Customer Service](#).

### Test Classification

This test was developed, and its performance characteristics determined by Mayo Clinic in a manner consistent with CLIA requirements. This test has not been cleared or approved by the US Food and Drug Administration.

### CPT Code Information

87798

### LOINC® Information

Test ID	Test Order Name	Order LOINC® Value
POWVS	Powassan Virus, PCR, Serum	55144-0

Result ID	Test Result Name	Result LOINC® Value
617490	Lineage 1	55144-0
618146	Lineage 2 (Deer Tick Virus)	55144-0
618150	Powassan Virus PCR Comment	77202-0