

## Overview

### Useful For

Identification of causative mutations when a deficient serum level of alpha-1-antitrypsin is not explained by routine testing, such as proteotyping, genotyping, or isoelectric focusing phenotyping.

Determining the specific allelic variant (full gene analysis) for prognosis and genetic counseling

### Testing Algorithm

See [Alpha-1 Antitrypsin-A Comprehensive Testing Algorithm](#) in Special Instructions.

### Special Instructions

- [Molecular Genetics: Congenital Inherited Diseases Patient Information](#)
- [Informed Consent for Genetic Testing](#)
- [Alpha 1 Antitrypsin-A Comprehensive Testing Algorithm](#)
- [Informed Consent for Genetic Testing \(Spanish\)](#)
- [Molecular Genetics: SERPINA1 Gene Patient Information](#)

### Method Name

Polymerase Chain Reaction (PCR) Amplification/DNA Sequencing

### NY State Available

Yes

## Specimen

### Specimen Type

Varies

### Shipping Instructions

Specimen preferred to arrive within 96 hours of collection.

### Specimen Required

**Patient Preparation:** A previous bone marrow transplant from an allogenic donor will interfere with testing. Call 800-533-1710 for instructions for testing patients who have received a bone marrow transplant.

**Specimen Type:** Whole blood

**Container/Tube:**

**Preferred:** Lavender top (EDTA) or yellow top (ACD)

**Acceptable:** Any anticoagulant

**Specimen Volume:** 3 mL

**Collection Instructions:**

1. Invert several times to mix blood.

2. Send specimen in original tube.

### Forms

1. **New York Clients-Informed consent is required.** Document on the request form or electronic order that a copy is on file. The following documents are available in Special Instructions:

-[Informed Consent for Genetic Testing](#) (T576)

-[Informed Consent for Genetic Testing-Spanish](#) (T826)

2. [Molecular Genetics: Congenital Inherited Diseases Patient Information](#) (T521) in Special Instructions and Forms.

3. If not ordering electronically, complete, print, and send a [Gastroenterology and Hepatology Client Test Request](#) (T728) with the specimen.

### Specimen Minimum Volume

1 mL

### Reject Due To

All specimens will be evaluated by Mayo Clinic Laboratories for test suitability.

### Specimen Stability Information

Specimen Type	Temperature	Time	Special Container
Varies	Refrigerated (preferred)		
	Ambient		
	Frozen		

## Clinical & Interpretive

### Clinical Information

Alpha-1-antitrypsin (A1A) is a protein that inhibits the enzyme neutrophil elastase. It is predominantly synthesized in the liver and secreted into the bloodstream. The inhibition function is especially important in the lungs because it protects against excess tissue degradation. Tissue degradation due to A1A deficiency is associated with an increased risk for early onset panlobular emphysema, which initially affects the lung bases (as opposed to smoking-related emphysema, which presents with upper lung field emphysema). Patients may become symptomatic in their 30s and 40s. The most frequent symptoms reported in a National Institute of Health study of 1,129 patients with severe deficiency (mean age 46 years) included cough (42%), wheezing (65%), and dyspnea with exertion (84%). Many patients were misdiagnosed as having asthma. It is estimated that approximately one-sixth of all lung transplants are for A1A deficiency.

Liver disease can also occur, particularly in children; it occurs much less commonly than emphysema in adults.

A1A deficiency is a relatively common disorder in the Northern European White population. The diagnosis of A1A deficiency is initially made by quantitation of protein levels in serum followed by phenotyping-determination of specific allelic variants by isoelectric focusing (IEF), genotyping-DNA based detection of specific mutations, or proteotyping-using liquid chromatography-tandem mass spectrometry (LC-MS/MS). While there are many different alleles in this gene, only 3 are common. The 3 major alleles include: M (full functioning, normal allele), S (associated with reduced levels of protein), and Z (disease-causing mutation associated with liver disease and premature emphysema). The S and Z alleles account for the majority of the abnormal alleles detected in affected patients. As a codominant disorder, both alleles are

---

expressed. An individual of SZ or S- null genotype may have a small increased risk for emphysema (but not liver disease) due to slightly reduced protein levels. On the other hand, an individual with the ZZ genotype is at greater risk for early onset liver disease and premature emphysema.

Smoking appears to hasten development of emphysema by 10 to 15 years. These individuals should be monitored closely for lung and liver function.

Historically, IEF phenotyping has been the primary method for characterizing variants, though in some cases the interpretation is difficult and prone to error. Serum quantitation is helpful in establishing a diagnosis but can be influenced by other factors. IEF phenotyping, LC-MS/MS proteotyping, and DNA-based genotyping are routinely used to test for deficiency alleles but can miss disease alleles other than the S and Z alleles. In patients suspected to have alpha-1 antitrypsin deficiency based on clinical findings or serum alpha-1-antitrypsin (AAT) levels, who do not have evidence of the SZ or ZZ genotype by routine methods, this gene analysis assay may provide useful information. Full sequencing of the *SERPINA1* coding region is performed for the detection of rare non-S or non-Z disease mutations.

See [Alpha-1-Antitrypsin-A Comprehensive Testing Algorithm](#) in Special Instructions.

### Reference Values

An interpretive report will be provided.

### Interpretation

All detected alterations are evaluated according to American College of Medical Genetics recommendations.(1) Variants are classified based on known, predicted, or possible pathogenicity and reported with interpretive comments detailing their potential or known significance.

### Cautions

A small percentage of individuals who are carriers or have a diagnosis of alpha-1 antitrypsin deficiency may have a mutation that is not identified by this method (eg, large genomic deletions, promoter mutations). The absence of a mutation, therefore, does not eliminate the possibility of positive carrier status or the diagnosis of alpha-1 antitrypsin deficiency. For testing of at risk family members, it is important to first document the presence of a *SERPINA1* gene mutation in an affected family member.

In some cases, DNA alterations of undetermined significance may be identified.

Test results should be interpreted in the context of clinical findings, family history, and other laboratory data. Errors in the interpretation of results may occur if information given is inaccurate or incomplete.

Rare polymorphisms exist that could lead to false-negative or false-positive results. If results obtained do not match the clinical findings, additional testing should be considered.

### Clinical Reference

1. Richards S, Aziz N, Bale S, et al: Standards and guidelines for the interpretation of sequence variants: a joint consensus recommendation of the American College of Medical Genetics and Genomics and the Association for Molecular Pathology. *Genet Med* 2015 May;17(5):405-424
2. Stoller JK, Aboussouan LS: Alpha-1-antitrypsin deficiency. *Lancet* 2005;365:2225-2236

3. McElvaney NG, Stoller JK, Buist AS, et al: Baseline characteristics of enrollees in the National Heart, Lung and Blood Institute Registry of alpha 1-antitrypsin deficiency. Alpha 1-Antitrypsin Deficiency Registry Study Group. Chest 1997;111:394-403
4. Snyder MR, Katzmann JA, Butz ML, et al: Diagnosis of alpha-1-antitrypsin deficiency: an algorithm of quantification, genotyping, and phenotyping. Clin Chem 2006;52:2236-2242
5. Graham RP, Dina MA, Howe SC, et al: SERPINA1 full-gene sequencing identifies rare mutations not detected in targeted mutation analysis. J Mol Diag 2015;17:689-694

## Performance

### Method Description

Bidirectional sequence analysis is performed to test for the presence of a mutation in all coding regions and intron and exon boundaries of the SERPINA1 gene.(Unpublished Mayo method)

### PDF Report

No

### Day(s) Performed

Varies

### Report Available

14 to 20 days

### Specimen Retention Time

Whole Blood: 2 weeks (if available), Extracted DNA: 3 months

### Performing Laboratory Location

Rochester

## Fees & Codes

### Fees

- Authorized users can sign in to [Test Prices](#) for detailed fee information.
- Clients without access to Test Prices can contact [Customer Service](#) 24 hours a day, seven days a week.
- Prospective clients should contact their account representative. For assistance, contact [Customer Service](#).

### Test Classification

This test was developed, and its performance characteristics determined by Mayo Clinic in a manner consistent with CLIA requirements. This test has not been cleared or approved by the US Food and Drug Administration.

### CPT Code Information

81479

---

**LOINC® Information**

Test ID	Test Order Name	Order LOINC® Value
SERPZ	SERPINA1 Gene, Full Gene Analysis	94222-7

Result ID	Test Result Name	Result LOINC® Value
113178	Result Summary	50397-9
113179	Result	82939-0
113180	Interpretation	69047-9
113181	Additional Information	48767-8
113182	Specimen	31208-2
113183	Source	31208-2
113184	Released By	18771-6