

Overview

Useful For

Monitoring trimipramine concentration during therapy

Evaluating potential trimipramine toxicity

May aid in evaluating patient compliance

Method Name

Liquid Chromatography-Tandem Mass Spectrometry (LC-MS/MS)

NY State Available

Yes

Specimen

Specimen Type

Serum Red

Specimen Required

Collection Container/Tube: Red top (Serum gel/SST are **not acceptable**)

Submission Container/Tube: Plastic vial

Specimen Volume: 1 mL

Collection Instructions:

1. Collect specimen immediately before next scheduled dose (minimum 12 hours after last dose).
2. Centrifuge and aliquot serum into plastic vial. **Serum must be separated from cells within 2 hours of collection.**

Forms

If not ordering electronically, complete, print, and send a [Therapeutics Test Request](#) (T831) with the specimen.

Specimen Minimum Volume

0.25 mL

Reject Due To

Gross hemolysis	Reject
Gross lipemia	Reject
Gross icterus	Reject

Specimen Stability Information

Specimen Type	Temperature	Time	Special Container
Serum Red	Refrigerated (preferred)	28 days	
	Frozen	28 days	
	Ambient	7 days	

Clinical & Interpretive

Clinical Information

Trimipramine is a tricyclic antidepressant with additional anxiety-reducing sedative activity. Daily dosages for adults range from 50 mg to 300 mg and are usually divided into 2 to 3 doses per day. Therapeutic ranges are based on serum samples collected at trough (ie, immediately before the next dose). Peak serum concentrations are typically achieved after 1 to 6 hours post dosage.

Common adverse effects include hypotension, tachycardia, constipation, dizziness, somnolence, and blurred vision. Risk of toxicity increases when concentrations exceed 500 ng/mL. Serious adverse effects include coma, seizures, and QRS prolongation with ventricular dysrhythmias.

Reference Values

Therapeutic concentration: 150-300 ng/mL

Note: Therapeutic ranges are for specimens collected at trough (ie, immediately before next scheduled dose). Levels may be elevated in non-trough specimens.

Interpretation

Most individuals display optimal response to trimipramine with serum levels of 150 to 300 ng/mL. Risk of toxicity is increased with trimipramine levels above 500 ng/mL.

Some individuals may respond well outside of this range or may display toxicity within the therapeutic range; thus, interpretation should include clinical evaluation.

Therapeutic ranges are based on specimens collected at trough (ie, immediately before the next dose).

Cautions

This test cannot be performed on whole blood. Serum must be separated from cells within 2 hours of collection; if serum is not removed within this time, tricyclic antidepressant levels may be falsely elevated due to drug release from red blood cells.

Specimens that are obtained from gel tubes are not acceptable because the drug can absorb on the gel and lead to falsely decreased concentrations.

Coadministration of fluvoxamine, moclobemide, or quinidine inhibits the metabolism and markedly increases the serum concentrations of trimipramine.

Clinical Reference

1. Wille SM, Cooreman SG, Neels HM, Lambert WE: Relevant issues in the monitoring and the toxicology of

antidepressants. Crit Rev Clin Lab Sci. 2008;45(1):25-89

2. Thanacoody HK, Thomas SHL: Antidepressant poisoning. Clin Med. 2003 Mar-Apr;3(2):114-118

3. Hiemke C, Baumann P, Bergemann N, et al: AGNP Consensus Guidelines for Therapeutic Drug Monitoring in Psychiatry: Update 2011. Pharmacopsychiatry. 2011 Sep;44(6):195-235

4. Burtis CA, Ashwood ER, Bruns DE, eds. Tietz Textbook of Clinical Chemistry and Molecular Diagnostics. 5th ed. Elsevier; 2012

Performance

Method Description

The tricyclic antidepressants are extracted from serum using a solvent crash to precipitate proteins. The supernatant is removed and analysis is by liquid chromatography-tandem mass spectrometry. (Unpublished Mayo method)

PDF Report

No

Day(s) Performed

Tuesday, Thursday, Sunday

Report Available

3 to 5 days

Specimen Retention Time

14 days

Performing Laboratory Location

Rochester

Fees & Codes

Fees

- Authorized users can sign in to [Test Prices](#) for detailed fee information.
- Clients without access to Test Prices can contact [Customer Service](#) 24 hours a day, seven days a week.
- Prospective clients should contact their Regional Manager. For assistance, contact [Customer Service](#).

Test Classification

This test was developed, and its performance characteristics determined by Mayo Clinic in a manner consistent with CLIA requirements. This test has not been cleared or approved by the US Food and Drug Administration.

CPT Code Information

80335

G0480 (if appropriate)

LOINC® Information

Test ID	Test Order Name	Order LOINC® Value
TRMP	Trimipramine, S	4083-2

Result ID	Test Result Name	Result LOINC® Value
64269	Trimipramine, S	4083-2