

Glycophorin A IHC, Tech Only

Overview

Useful For

Aiding in the identification of erythroid precursors in bone marrow

Testing Algorithm

For the initial technical component only immunohistochemical (IHC) stain performed, the appropriate bill-only test ID will be reflexed and charged (IHTOI). For each additional technical component only IHC stain performed, an additional bill-only test ID will be reflexed and charged (IHTOA).

Reflex Tests

Test Id	Reporting Name	Available Separately	Always Performed
IHTOI	IHC Initial, Tech Only	No	No
IHTOA	IHC Additional, Tech Only	No	No

Method Name

Immunohistochemistry (IHC)

NY State Available

Yes

Specimen

Specimen Type

TECHONLY

Ordering Guidance

This test includes only technical performance of the stain (no pathologist interpretation is performed). If diagnostic consultation by a pathologist is required order PATHC / Pathology Consultation.

Shipping Instructions

Attach the green pathology address label and the pink Immunostain Technical Only label included in the kit to the outside of the transport container.

Specimen Required

Supplies: Immunostain Technical Only Envelope (T693)

Specimen Type: Tissue



Glycophorin A IHC, Tech Only

Container/Tube: Immunostain Technical Only Envelope

Preferred: 2 Unstained positively charged glass slide (25- x 75- x 1-mm) per test ordered; sections 4 microns thick

Acceptable: Formalin-fixed, paraffin-embedded (FFPE) tissue block

Digital Image Access

- 1. Information on accessing digital images of IHC stains and the manual requisition form can be accessed through this website: https://news.mayocliniclabs.com/ihc-stains/
- 2. Clients ordering stains using a manual requisition form will not have access to digital images.
- 3. Clients wishing to access digital images must place the order for IHC stains electronically. Information regarding digital imaging can be accessed through this website: https://news.mayocliniclabs.com/ihc-stains/#FAQ

Forms

If not ordering electronically, complete, print, and send a <u>Immunohistochemical (IHC)/In Situ Hybridization (ISH) Stains</u>
Request (T763) with the specimen.

Reject Due To

Wet/frozen tissue Reject
Cytology smears
Nonformalin fixed tissue
Nonparaffin embedded tissue
Noncharged slides
ProbeOn slides

Specimen Stability Information

Specimen Type	Temperature	Time	Special Container
TECHONLY	Ambient (preferred)		
	Refrigerated		

Clinical & Interpretive

Clinical Information

Glycophorin A is expressed by erythroid precursors in the bone marrow and is also present on erythrocytes. Glycophorin A is useful to characterize erythroid cell development and aid in the diagnosis of erythroid leukemia.

Reference Values

NA



Glycophorin A IHC, Tech Only

Interpretation

This test does not include pathologist interpretation; only technical performance of the stain is performed. If an interpretation is required, order PATHC / Pathology Consultation for a full diagnostic evaluation or second opinion of the case.

The positive and negative controls are verified as showing appropriate immunoreactivity and documentation is retained at Mayo Clinic Rochester. If a control tissue is not included on the slide, a scanned image of the relevant quality control tissue is available upon request, call 855-516-8404.

Interpretation of this test should be performed in the context of the patient's clinical history and other diagnostic tests by a qualified pathologist.

Cautions

Age of a cut paraffin section can affect immunoreactivity. Stability thresholds vary widely among published literature and are antigen dependent. Best practice is for paraffin sections to be cut within 6 weeks.

Clinical Reference

- 1. Dong HY, Wilkes S, Yang H: CD71 is selectively and ubiquitously expressed at high levels in erythroid precursors of all maturation stages: a comparative immunochemical study with glycophorin A and hemoglobin A. Am J Surg Pathol. 2011 May;35(5):723-732
- 2. Rollins-Raval MA, Fuhrer K, Marafioti T, Roth CG: ALDH, CA I, and CD2AP: novel, diagnostically useful immunohistochemical markers to identify erythroid precursors in bone marrow biopsy specimens. Am J Clin Pathol. 2012 Jan;137(1):30-38
- 3. Liu W, Hasserjian RP, Hu Y, et al: Pure erythroid leukemia: a reassessment of the entity using the 2008 World Health Organization Classification. Mod Pathol. 2011 Mar;24(3):375-383

Performance

Method Description

Immunohistochemistry on sections of paraffin-embedded tissue.(Unpublished Mayo method)

PDF Report

Nο



Glycophorin A IHC, Tech Only

Specimen Retention Time

Until staining is complete.

Performing Laboratory Location

Rochester

Fees & Codes

Test Classification

This test has been cleared, approved, or is exempt by the US Food and Drug Administration and is used per manufacturer's instructions. Performance characteristics were verified by Mayo Clinic in a manner consistent with CLIA requirements.

CPT Code Information

88342-TC, primary

88341-TC, if additional IHC

LOINC® Information

GLYCF Glycophorin A IHC, Tech Only Order only;no result	Test ID	Test Order Name	Order LOINC Value
	GLYCF	Glycophorin A IHC, Tech Only	Order only;no result

Result ID	Reporting Name	LOINC®
70761	Glycophorin A IHC, Tech Only	Bill only; no result