

Overview

Useful For

Aiding in the identification of neoplastic cells

Reflex Tests

Test Id	Reporting Name	Available Separately	Always Performed
IHTOI	IHC Initial, Tech Only	No	No
IHTOA	IHC Additional, Tech Only	No	No

Testing Algorithm

[For the initial technical component only immunohistochemical \(IHC\) stain performed, the appropriate bill-only test ID will be reflexed and charged \(IHTOI\). For each additional technical component only IHC stain performed, an additional bill-only test ID will be reflexed and charged \(IHTOA\).](#)

Method Name

Immunohistochemistry (IHC)

NY State Available

Yes

Specimen

Specimen Type

TECHONLY

Ordering Guidance

This test includes only technical performance of the stain (no pathologist interpretation is performed). If diagnostic consultation by a pathologist is required order PATHC / Pathology Consultation.

Shipping Instructions

Attach the green pathology address label and the pink Immunostain Technical Only label included in the kit to the outside of the transport container.

Specimen Required

Supplies: Immunostain Technical Only Envelope (T693)

Specimen Type: Tissue

Container/Tube: Immunostain Technical Only Envelope

Preferred: 2 Unstained positively charged glass slide (25- x 75- x 1-mm) per test ordered; sections 4-microns thick

Acceptable: Formalin-fixed, paraffin-embedded (FFPE) tissue block

Digital Image Access

1. Information on accessing digital images of immunohistochemical (IHC) stains and the manual requisition form can be accessed through this website: <https://news.mayocliniclabs.com/ihc-stains/>
2. Clients ordering stains using a manual requisition form will not have access to digital images.
3. Clients wishing to access digital images must place the order for IHC stains electronically. Information regarding digital imaging can be accessed through this website: <https://news.mayocliniclabs.com/ihc-stains/#FAQ>

Forms

If not ordering electronically, complete, print, and send a [Immunohistochemical \(IHC\)/In Situ Hybridization \(ISH\) Stains Request](#) (T763) with the specimen.

Reject Due To

Wet/frozen tissue Cytology smears Nonformalin fixed tissue Nonparaffin embedded tissue Noncharged slides ProbeOn slides	Reject
--	--------

Specimen Stability Information

Specimen Type	Temperature	Time	Special Container
TECHONLY	Ambient (preferred)		
	Refrigerated		

Clinical & Interpretive

Clinical Information

p53 is a tumor-suppressor protein. Genetic events (variant and deletion) that affect both *P53* alleles can lead to loss of cell cycle control in the setting of DNA damage, resulting in genetic instability and neoplastic transformation. Altered p53 also has a prolonged half-life compared to wildtype p53 and, thus, accumulates in the nucleus and can be detected by immunohistochemistry. Abnormalities of the *P53* gene are one of the most common genetic changes associated with cancer and can be found in a wide variety of tumor types, where they are generally associated with a worse prognosis. The p53 protein can be readily detected in a subset of cancers of the [colon, stomach, bladder, breast, lung, and testes and in melanoma and lymphoma](#).

Interpretation

This test does not include pathologist interpretation, only technical performance of the stain. If interpretation is required, order PATHC / Pathology Consultation for a full diagnostic evaluation or second opinion of the case.

The positive and negative controls are verified as showing appropriate immunoreactivity and documentation is retained at Mayo Clinic Rochester. If a control tissue is not included on the slide, a scanned image of the relevant quality control tissue is available upon request, call 855-516-8404.

Interpretation of this test should be performed in the context of the patient's clinical history and other diagnostic tests by a qualified pathologist.

Cautions

Age of a cut paraffin section can affect immunoreactivity. Stability thresholds vary widely among published literature and are antigen dependent. Best practice is for paraffin sections to be cut within 6 weeks.

Clinical Reference

1. Bartley AN, Ross DW: Validation of p53 immunohistochemistry as a prognostic factor in breast cancer in clinical practice. *Arch Pathol Lab Med.* 2002 Apr;126(4):456-458
2. Camelo-Piragua S, Jansen M, Ganguly A, et al: A sensitive and specific diagnostic panel to distinguish diffuse astrocytoma from astrocytosis: chromosome 7 gain with mutant isocitrate dehydrogenase 1 and p53. *J NeuroPathol Exp Neurol.* 2011 Feb;70(2):110-115
3. Klemi PJ, Pylkkanen L, Kiilholma P, Kurvinen K, Joensuu H: p53 protein detected by immunohistochemistry as a prognostic factor in patients with epithelial ovarian carcinoma. *Cancer.* 1995 Oct 1;76(7):1201-1208
4. Mayall FG, Goddard H, Gibbs AR: p53 immunostaining in the distinction between benign and malignant mesothelial proliferations using formalin-fixed paraffin sections. *J Pathol.* 1992 Dec;168(4):377-381
5. van den Berg FM, Baas IO, Polak MM, Offerhaus GJ: Detection of p53 overexpression in routinely paraffin-embedded tissue of human carcinomas using a novel target unmasking fluid. *Am J Pathol.* 1993 Feb;142(2):381-385
6. Zarei S, Wang Y, Jenkins SM, Voss JS, Kerr SE, Bell DA: Clinicopathologic, immunohistochemical, and molecular characteristics of ovarian serous carcinoma with mixed morphologic features of high-grade and low-grade serous carcinoma. *Am J Surg Pathol.* 2020 Mar;44(3):316-328. doi: 10.1097/PAS.0000000000001419

Performance**Method Description**

Immunohistochemistry on sections of paraffin-embedded tissue.(Unpublished Mayo method)

PDF Report

No

Day(s) Performed

Monday through Friday

Report Available

1 to 3 days

Specimen Retention Time

Until staining is complete.

Performing Laboratory Location

Rochester

Fees & Codes**Fees**

- Authorized users can sign in to [Test Prices](#) for detailed fee information.
- Clients without access to Test Prices can contact [Customer Service](#) 24 hours a day, seven days a week.
- Prospective clients should contact their account representative. For assistance, contact [Customer Service](#).

Test Classification

This test was developed, and its performance characteristics determined by Mayo Clinic in a manner consistent with CLIA requirements. This test has not been cleared or approved by the US Food and Drug Administration.

CPT Code Information

88342-TC, primary

88341-TC, if additional IHC

LOINC® Information

Test ID	Test Order Name	Order LOINC® Value
P53	p53 IHC, Tech Only	Order only;no result

Result ID	Test Result Name	Result LOINC® Value
70840	p53 IHC, Tech Only	Bill only; no result