

Overview

Useful For

[Distinguishing gliomas from neurocytomas and ependymomas](#)

Testing Algorithm

[For the initial technical component only immunohistochemical \(IHC\) stain performed, the appropriate bill-only test ID will be reflexed and charged \(IHTOI\). For each additional technical component only IHC stain performed, an additional bill-only test ID will be reflexed and charged \(IHTOA\).](#)

Reflex Tests

Test Id	Reporting Name	Available Separately	Always Performed
IHTOI	IHC Initial, Tech Only	No	No
IHTOA	IHC Additional, Tech Only	No	No

Method Name

Immunohistochemistry

NY State Available

Yes

Specimen

Specimen Type

TECHONLY

Ordering Guidance

This test includes only technical performance of the stain (no pathologist interpretation is performed). If diagnostic consultation by a pathologist is required order PATHC / Pathology Consultation.

Shipping Instructions

Attach the green pathology address label and the pink Immunostain Technical Only label included in the kit to the outside of the transport container.

Specimen Required

Supplies: Immunostain Technical Only Envelope (T693)

Specimen Type: Tissue

Preferred: 2 Unstained positively charged glass slide (25- x 75- x 1-mm) per test ordered; sections 4-microns thick.

Acceptable: Formalin-fixed, paraffin-embedded (FFPE) tissue block

Digital Image Access

1. Information on accessing digital images of IHC stains and the manual requisition form can be accessed through this website: <https://news.mayocliniclabs.com/ihc-stains/>
2. Clients ordering stains using a manual requisition form will not have access to digital images.
3. Clients wishing to access digital images must place the order for IHC stains electronically. Information regarding digital imaging can be accessed through this website: <https://news.mayocliniclabs.com/ihc-stains/#FAQ>

Forms

If not ordering electronically, complete, print, and send a [Immunohistochemical \(IHC\)/In Situ Hybridization \(ISH\) Stains Request](#) (T763) with the specimen.

Reject Due To

Tissue/Other Wet/frozen tissue
 Cytology smears
 Nonformalin fixed tissue
 Nonparaffin embedded tissue
 Noncharged slides
 ProbeOn slides

Specimen Stability Information

Specimen Type	Temperature	Time	Special Container
TECHONLY	Ambient (preferred)		
	Refrigerated		

Clinical & Interpretive

Clinical Information

Oligodendrocyte transcription factor 2 (OLIG2) is a transcription factor that participates in oligodendrocyte and motor neuron differentiation. During embryogenesis OLIG2 promotes the growth of motor neuron progenitor cells. OLIG2 expression decreases upon further neuronal differentiation. OLIG2 is also involved in oligodendrocyte differentiation where expression remains present in mature glial cells. In gliomas, OLIG2 represses the p53 tumor suppressor pathway, thereby contributing to glioma progression.

Interpretation

This test includes only technical performance of the stain (no pathologist interpretation is performed). Mayo Clinic cannot provide an interpretation of tech only stains outside the context of a pathology consultation. If an interpretation is needed, refer to PATHC / Pathology Consultation for a full diagnostic evaluation or second opinion of the case. All material associated with the case is required. Additional specific stains may be requested as part of the pathology consultation, and will be performed as necessary at the discretion of the Mayo pathologist.

The positive and negative controls are verified as showing appropriate immunoreactivity and documentation is retained at Mayo Clinic Rochester. If a control tissue is not included on the slide, a scanned image of the relevant quality control tissue is available upon request. Contact 855-516-8404.

Interpretation of this test should be performed in the context of the patient's clinical history and other diagnostic tests by a qualified pathologist.

Cautions

Age of a cut paraffin section can affect immunoreactivity. Stability thresholds vary widely among published literature and are antigen-dependent. Best practice is for paraffin sections to be cut within 6 weeks.

Clinical Reference

1. Suzuki A, Nobusawa S, Natsume A, et al: Olig2 labeling index is correlated with histological and molecular classifications in low-grade diffuse gliomas. *J Neurooncol* 2014 Nov;120(2):283-291
2. Popova SN, Bergqvist M, Dimberg A, et al: Subtyping of gliomas of various WHO grades by the application of immunohistochemistry: *Histopathology* 2014 Feb;64(3):365-379
3. Ballester LY, Wang Z, Shandilya S, et al: Morphologic characteristics and immunohistochemical profile of diffuse intrinsic pontine gliomas. *Am J Surg Pathol* 2013 Sep;37(9):1357-1364

Performance**Method Description**

Immunohistochemistry on sections of paraffin-embedded tissue.(Unpublished Mayo method)

PDF Report

No

Specimen Retention Time

Until staining is complete.

Performing Laboratory Location

Rochester

Fees & Codes**Test Classification**

This test has been cleared, approved, or is exempt by the US Food and Drug Administration and is used per manufacturer's instructions. Performance characteristics were verified by Mayo Clinic in a manner consistent with CLIA requirements.

CPT Code Information

88342-TC, primary

88341-TC, if additional IHC

LOINC® Information

Test ID	Test Order Name	Order LOINC Value
OLIG2	OLIG2 IHC, Tech Only	Order only;no result

Result ID	Reporting Name	LOINC®
71361	OLIG2 IHC, Tech Only	Bill only; no result