

Overview

Useful For

Determining proliferation of tumor cells in paraffin-embedded tissue blocks from patients diagnosed with carcinoid or atypical carcinoid of the lung including metastases

Reflex Tests

Test Id	Reporting Name	Available Separately	Always Performed
KIPM	Ki67 Pulmonary IHC Manual	No	No

Testing Algorithm

Cases that are unable to be scanned for automated analysis will be changed to KIPM / Ki-67(MIB-1), Pulmonary, Quantitative Immunohistochemistry, Manual.

Method Name

Immunohistochemistry, Automated Quantitation, Hot-Spot Technique

NY State Available

Yes

Specimen

Specimen Type

Special

Shipping Instructions

Attach the green "Attention Pathology" address label (T498) to the outside of the transport container before putting into the courier mailer.

Necessary Information

- 1. Pathologist's name, address, and phone number are required.
- 2. Include accompanying pathology report stating the final diagnosis.

Specimen Required

Supplies: Pathology Packaging Kit (T554)

Specimen Type:

Preferred: Formalin-fixed, paraffin-embedded tissue block containing carcinoid/atypical carcinoid tumor of the lung including metastases.

Acceptable: 2 Unstained sections containing carcinoid/atypical carcinoid tumor of the lung including metastases on charged slides cut at 4 microns <1 month ago. Tissue on the slides should have been fixed in 10% neutral buffered formalin.

Container/Tube: Pathology Packaging Kit

Collection Instructions: Submit formalin-fixed, paraffin-embedded tissue block.

Additional Information: Paraffin block will be returned with the final report.

Forms

If not ordering electronically, complete, print, and send a [Immunohistochemical \(IHC\)/In Situ Hybridization \(ISH\) Stains Request](#) (T763) with the specimen.

Reject Due To

All specimens will be evaluated at Mayo Clinic Laboratories for test suitability.

Specimen Stability Information

Specimen Type	Temperature	Time	Special Container
Special	Ambient (preferred)		
	Refrigerated		

Clinical & Interpretive

Clinical Information

Ki-67(MIB-1 clone) is a monoclonal antibody that reacts with cells undergoing DNA synthesis by binding to the Ki-67 antigen, a marker known to be expressed only in proliferating cells. By measuring the amount of tumor cells expressing Ki-67, an estimate of DNA synthesis can be determined. Studies suggest that Ki-67(MIB-1) analysis of paraffin-embedded tissue specimens may provide useful prognostic information in various tumor types.

Reference Values

Varies by tumor type; values reported from 0% to 100%

Interpretation

Results will be reported as a percentage of tumor cells staining positive for Ki-67(MIB-1). Quantitative Ki-67(MIB-1) results should be interpreted within the clinical context for which the test was ordered.

The scoring method using Aiforia artificial intelligence for image analysis was developed and validated in the Biomarker and Image Analysis Laboratory, Department of Laboratory Medicine and Pathology, Mayo Clinic (see Method Description).

Cautions

The paraffin block analyzed must be representative of the patient's tumor.

Test results should be interpreted in the context of clinical findings and other laboratory data.

Clinical Reference

1. Boland JM, Kroneman TN, Jenkins SM, et al. Ki-67 Labeling index in pulmonary carcinoid tumors: Comparison between small biopsy and resection using tumor tracing and hot spot methods. Arch Pathol Lab Med. 2020;144(8):982-990
2. La Rosa S. Diagnostic, Prognostic, and Predictive Role of Ki67 Proliferative Index in Neuroendocrine and Endocrine Neoplasms: Past, Present, and Future. Endocr Pathol. 2023;34(1):79-97. doi:10.1007/s12022-023-09755-3

Performance

Method Description

A 4-micron thick section is cut from the paraffin block. The section is stained with an immunoperoxidase method using the monoclonal antibody Ki-67 (MIB-1 clone). This is the paraffin nuclear epitope to the Ki-67 antigen. Any nucleus that has an antigen-antibody complex will cause the bright-field, brown chromogen, diaminobenzidine (DAB), to precipitate onto it. All nuclei, both DAB-positive and -negative, are counterstained with diluted hematoxylin.

Ki-67(MIB-1)-stained slides are scanned using the Leica Aperio GT450 digital scanner. The captured digital image is analyzed using a validated AI (Artificial Intelligence) algorithm by Aiforia that calculates the percentage of positive staining tumor nuclei. The Aiforia data and corresponding slide are reviewed by a pathologist for final interpretation.(Unpublished Mayo method)

PDF Report

No

Day(s) Performed

Monday through Friday

Report Available

4 to 6 days

Specimen Retention Time

1 week after results are reported. Material made at Mayo Clinic may be retained at Mayo Clinic indefinitely.

Performing Laboratory Location

Mayo Clinic Laboratories - Rochester Main Campus

Fees & Codes

Fees

- Authorized users can sign in to [Test Prices](#) for detailed fee information.
- Clients without access to Test Prices can contact [Customer Service](#) 24 hours a day, seven days a week.
- Prospective clients should contact their account representative. For assistance, contact [Customer Service](#).

Test Classification

This test was developed and its performance characteristics determined by Mayo Clinic in a manner consistent with CLIA requirements. It has not been cleared or approved by the US Food and Drug Administration.

CPT Code Information

88361

LOINC® Information

Test ID	Test Order Name	Order LOINC® Value
KI67P	Ki67 Pulmonary IHC Automated	In Process

Result ID	Test Result Name	Result LOINC® Value
72132	Interpretation	29593-1
72133	Participated in the Interpretation	No LOINC Needed
72134	Report electronically signed by	19139-5
72135	Material Received	81178-6
72136	Disclaimer	62364-5
72137	Case Number	80398-1