



Test Definition: AMS

Amylase, Total, Serum

Overview

Useful For

Diagnosis and management of pancreatitis

Evaluation of pancreatic function

Method Name

Colorimetric Rate Reaction

NY State Available

No

Specimen

Specimen Type

Serum

Necessary Information

Patient's age and sex are required.

Specimen Required

Collection Container/Tube:

Preferred: Serum gel

Acceptable: Red top

Submission Container/Tube: Plastic vial

Specimen Volume: 0.5 mL

Collection Instructions:

1. Serum gel tubes should be centrifuged within 2 hours of collection.
2. Red-top tubes should be centrifuged, and the serum aliquoted into a plastic vial within 2 hours of collection.

Specimen Minimum Volume

0.25 mL

Reject Due To

Gross hemolysis	Reject
Gross lipemia	OK

Specimen Stability Information

Specimen Type	Temperature	Time	Special Container
Serum	Ambient (preferred)	7 days	
	Refrigerated	30 days	
	Frozen	30 days	

Clinical & Interpretive

Clinical Information

The amylase enzymes are a group of hydrolases that degrade complex carbohydrates into fragments. Amylase is produced primarily by the exocrine pancreas where the enzyme is synthesized by the acinar cells and then secreted into the intestinal tract by way of the pancreatic duct system. Amylases also are produced by the salivary glands, small intestine mucosa, ovaries, placenta, liver, and fallopian tubes. Pancreatic and salivary isoenzymes are found in serum.

Reference Values

0-30 days: 0-6 U/L

31-182 days: 1-17 U/L

183-365 days: 6-44 U/L

1-3 years: 8-79 U/L

4-17 years: 21-110 U/L

> or =18 years: 28-100 U/L

Interpretation

In acute pancreatitis, a transient rise in serum amylase activity occurs within 2 to 12 hours of onset; levels return to normal by the third or fourth day. A 4- to 6-fold elevation of amylase activity above the reference limit is usual with the maximal levels obtained in 12 to 72 hours. However, a significant number of subjects show lesser elevations and sometimes none. The magnitude of the elevation of serum enzyme activity is not related to the severity of pancreatic involvement. Normalization is not necessarily a sign of resolution.

In acute pancreatitis associated with hyperlipidemia, serum amylase activity may be spuriously normal; the amylasemia may be unmasked either by serial dilution of the serum or ultracentrifugation.

A significant amount of serum amylase is excreted in the urine and, therefore, elevation of serum activity is reflected in the rise of urinary amylase activity. Urine amylase, as compared to serum amylase, appears to be more frequently elevated, reaches higher levels, and persists for longer periods. However, the receiver operator curves (ROC) of various serum and urine amylase assays demonstrated that all urine assays had poorer diagnostic utility than all serum assays. In quiescent chronic pancreatitis, both serum and urine activities are usually subnormal.

Because it is produced by several organs, amylase is not a specific indicator of pancreatic function. Elevated levels also may be seen in a number of nonpancreatic disease processes including mumps, salivary duct obstruction, ectopic pregnancy, and intestinal obstruction/infarction.

Cautions

Amylase results may be elevated in patients with macroamylase. Macroamylase refers to a high-molecular weight form of amylase that is present in a patient's serum. Different causes of macroamylase have been suggested, the most

common being amylase complexed with an immunoglobulin. The large size of the macroamylase complex prevents its excretion in the urine. As a result, the serum amylase is usually elevated. This elevated amylase is not diagnostic for pancreatitis. By utilizing serum lipase and urinary amylase, the presence or absence of macroamylase may be determined.

Clinical Reference

1. Soldin SJ: Pediatric Reference Ranges. 2nd ed AACC Press; 1997
2. Rifai N, Horvath AR, Wittwer CT, eds: Tietz Textbook of Clinical Chemistry and Molecular Diagnostics. 6th ed. Elsevier; 2018
3. Swaroop VS, Chari ST, Clain JE: Acute pancreatitis. JAMA. 2004;291:2865-2868
4. Azzopardi E, Lloyd C, Teixeira SR, Conlan, RS, Whitaker, IS: Clinical applications of amylase: Novel perspectives. Surgery. 2016;160(1):26-37

Performance**Method Description**

The liquid Roche amylase method is an enzymatic colorimetric test using 4,6-ethyliden (G7)-p-nitrophenol (G1)-alpha, D-maltoheptaoside (ethylidene-G7PNP) as a substrate. Human salivary and pancreatic amylases convert the substrate at approximately the same rate. The alpha-amylase cleaves the substrate into G2, G3, and G4 PNP fragments. The G2 and G3 and G4 PNP fragments are further hydrolyzed by an alpha-glucosidase to yield p-nitrophenol and glucose. The rate of increase in absorbance at 415 nm (measuring the increase in p-nitrophenol) is proportional to amylase activity. (Package insert: Roche AMYL2 reagent. Roche Diagnostics; 12/2018)

PDF Report

No

Day(s) Performed

Monday through Saturday

Report Available

Same day/1 to 2 days

Specimen Retention Time

1 week

Performing Laboratory Location

Mayo Clinic Jacksonville Clinical Lab

Fees & Codes**Fees**

- Authorized users can sign in to [Test Prices](#) for detailed fee information.

-
- Clients without access to Test Prices can contact [Customer Service](#) 24 hours a day, seven days a week.
 - Prospective clients should contact their account representative. For assistance, contact [Customer Service](#).

Test Classification

This test has been cleared, approved, or is exempt by the US Food and Drug Administration and is used per manufacturer's instructions. Performance characteristics were verified by Mayo Clinic in a manner consistent with CLIA requirements.

CPT Code Information

82150

LOINC® Information

Test ID	Test Order Name	Order LOINC® Value
AMS	Amylase, Total, S	1798-8

Result ID	Test Result Name	Result LOINC® Value
AMS	Amylase, Total, S	1798-8