Thank you for using version 2025.4.1104.462 of Telerik Document Processing Libraries. Access to the latest updates and support requires a license. Learn more in the documentation below. Learn more about Telerik Document Processing Libraries licensing https://docs.telerik.com/devtools/document-processing/distribution-and-licensing/license-key

MAYO CLINIC LABORATORIES

Test Definition: C1Q

Complement C1q, Serum

Overview

Useful For

Assessment of an undetectable total complement level

Diagnosing congenital C1 (first component of complement) deficiency

Diagnosing acquired deficiency of C1 inhibitor

Method Name

Nephelometry

NY State Available

Yes

Specimen

Specimen Type

Serum

Specimen Required

Patient Preparation:

Fasting: 12 hours, preferred but not required

Supplies: Sarstedt Aliquot Tube, 5 mL (T914)

Collection Container/Tube:

Preferred: Serum gel

Acceptable: Red top

Submission Container/Tube: Plastic vial

Specimen Volume: 1 mL

Collection Instructions: Centrifuge and aliquot serum into a plastic vial

Specimen Minimum Volume

0.5 mL

Reject Due To

Gross hemolysis	OK
Gross lipemia	Reject
Gross icterus	OK

Specimen Stability Information

Specimen Type	Temperature	Time	Special Container
Serum	Refrigerated (preferred)	28 days	
	Ambient	21 days	
	Frozen	28 days	

Clinical & Interpretive

Clinical Information

The first component of complement (C1) is composed of 3 subunits designated as C1q, C1r, and C1s. C1q recognizes and binds to immunoglobulin complexed to antigen and initiates the complement cascade. Congenital deficiencies of any of the early complement components (C1, C2, C4) results in an inability to clear immune complexes. Inherited deficiency of C1 is rare.

Like the more common C2 deficiency, C1 deficiency is associated with increased incidence of immune complex disease (systemic lupus erythematosus, polymyositis, glomerulonephritis, and Henoch-Schonlein purpura). Low C1 levels have also been reported in patients with abnormal immunoglobulin levels (Bruton and common variable hypogammaglobulinemia and severe combined immunodeficiency). This is most likely due to increased catabolism.

The measurement of C1q is an indicator of the amount of C1 present.

Reference Values

12-22 mg/dL

Interpretation

An undetectable C1q in the presence of an absent total complement and normal C2, C3, and C4 suggests a congenital C1 (first component of complement) deficiency.

A low C1q in combination with a low C1 inhibitor and low C4 suggests an acquired C1 inhibitor deficiency.

Cautions

This is a different assay than C1q binding, which is an assay for circulating immune complexes.

Clinical Reference

1. Stegert M, Bock M, Trendelenburg M. Clinical presentation of human C1q deficiency: How much of a lupus? Mol Immunol. 2015;67(1):3-11
2. Tangye SG, Al-Herz W, Bousfiha A, et al. Human inborn errors of immunity: 2022 update on the classification from the International Union of Immunological Societies Expert Committee. J Clin Immunol. 2022;42(7):1473-1507
3. Beurskens FJ, van Schaarenburg RA, Trouw LA. C1q, antibodies and anti-C1q autoantibodies. Mol Immunol. 2015;68(1):6-13

Performance

Test Definition: C1Q

Complement C1q, Serum

Method Description
Nephelometry and anti-C1q antiserum are used to quantitate the C1q antigen level.(Instruction manual: Siemens Nephelometer II. Siemens, Inc; Version 2.3, 2008; Addendum to the Instruction Manual 2.3, 08/2017)

PDF Report
No

Day(s) Performed
Monday through Friday

Report Available
2 to 5 days

Specimen Retention Time
14 days

Performing Laboratory Location
Mayo Clinic Laboratories - Rochester Superior Drive

Fees & Codes

Fees
- Authorized users can sign in to Test Prices for detailed fee information.
- Clients without access to Test Prices can contact Customer Service 24 hours a day, seven days a week.
- Prospective clients should contact their account representative. For assistance, contact Customer Service.

Test Classification
This test was developed and its performance characteristics determined by Mayo Clinic in a manner consistent with CLIA requirements. It has not been cleared or approved by the US Food and Drug Administration.

CPT Code Information
86160

LOINC® Information

Test ID	Test Order Name	Order LOINC® Value
C1Q	Complement C1q, S	4478-4

Result ID	Test Result Name	Result LOINC® Value
C1Q	Complement C1q, S	4478-4