

Overview

Method Name

Enzymeimmunoassay(EIA)

NY State Available

Yes

Specimen

Specimen Type

Serum

Specimen Required**Specimen Type: Serum****Container/Tube: Red or SST****Specimen Volume: 2 mL****Collection Instructions:** Draw blood in a plain red-top tube(s). (Serum gel tube is acceptable.) Spin down and send 2 mL serum refrigerate in a plastic vial**Reject Due To**

Hem NA

olysis

Lipe NA

mia

Icter NA

us

Other Specimen that is too viscous to pipette. Tissue, biopsy, sputum, bronchial aspirate, FNA, bone marrow aspirate, stool or samples in transport media, fixative or Isolator tubes.

Specimen Minimum Volume

1.2 mL

Specimen Stability Information

Specimen Type	Temperature	Time	Special Container
Serum	Refrigerated (preferred)	14 days	
	Ambient	14 days	
	Frozen		

Clinical & Interpretive

Reference Values

Reference interval: None Detected

Reportable Range: Positive Results reported in ng/mL from 0.20 ng/mL to 20.00 ng/mL.

Positive Results above 20.00 ng/mL are reported as "Above the Limit of Quantification".

Cautions

Sodium hydroxide and sputolysin. Cross-reactivity occurs between blastomycosis and histoplasmosis, and in Paracoccidioidomycosis, penicillosis, coccidioidomycosis, aspergillosis, and sporotrichosis.

Performance**PDF Report**

No

Performing Laboratory Location

MiraVista Diagnostics

Fees & Codes**Test Classification**

This test was developed and its performance characteristics determined by MiraVista Diagnostics. It has not been cleared or approved by the FDA; however, FDA clearance or approval is not currently required for clinical use. The results are not intended to be used as the sole means for clinical diagnosis or patient management decisions.

CPT Code Information

87385