

Test Definition: CURCU

Copper/Creatinine Ratio, Random, Urine

Reporting Title: Copper/Creat Ratio, Random, U

Performing Location: Mayo Clinic Laboratories - Rochester Superior Drive

Specimen Requirements:

Patient Preparation: High concentrations of barium are known to interfere with this test. If barium-containing contrast media has been administered, **the specimen should not be collected for at least 96 hours.**

Supplies: Urine Tubes, 10 mL (T068)

Collection Container/Tube: Clean, plastic urine collection container with no metal cap or glued insert

Submission Container/Tube: Plastic, 10-mL urine tube or a clean, plastic aliquot container with no metal cap or glued insert

Specimen Volume: 3 mL

Collection Instructions:

1. Collect a random urine specimen.

2. See <u>Metals Analysis Specimen Collection and Transport</u> for complete instructions.

Forms:

Specimen Type	Temperature	Time	Special Container
Urine	Refrigerated (preferred)	28 days	
	Ambient	14 days	
	Frozen	28 days	

Result Codes:

Result ID	Reporting Name	Туре	Unit	LOINC®
CRETR	Creatinine, Random, U	Numeric	mg/dL	2161-8
615258	Copper/Creat Ratio, U	Numeric	mcg/g Cr	13829-7

LOINC[®] and CPT codes are provided by the performing laboratory.

Supplemental Report:

No

Components:

Test Id	Reporting Name	CPT Units	CPT Code	Always Performed	Available Separately
CURC	Copper/Creat Ratio, U	1	82525	Yes	No
CRETR	Creatinine, Random, U	1	82570	Yes	No

CPT Code Information:

82525 82570



Test Definition: CURCU

Copper/Creatinine Ratio, Random, Urine

Reference Values:

COPPER/CREATITINE: Males: 0-17 years: Not established > or =18 years: 9-43 mcg/g creatinine Females: 0-17 years: Not established > or =18 years: 7-72 mcg/g creatinine

CREATITINE:

> or =18 years old: 16-326 mg/dL
Reference values have not been established for patients who are younger than 18 years.