

**Reporting Title:** T4 (Thyroxine), Total Only, S**Performing Location:** Mayo Clinic Jacksonville Clinical Lab**Ordering Guidance:**

This test cannot be used in patients receiving treatment with lipid-lowering agents containing dextrothyroxine unless therapy is discontinued for 4 to 6 weeks to allow the physiological state to become reestablished prior to testing.

**Specimen Requirements:****Patient Preparation:** For 4 to 6 weeks before specimen collection, patient should not be taking lipid-lowering medications containing dextrothyroxine.**Supplies:** Sarstedt Aliquot Tube, 5 mL (T914)**Collection Container/Tube:****Preferred:** Serum gel**Acceptable:** Red top**Submission Container/Tube:** Plastic vial**Specimen Volume:** 1 mL**Collection Instructions:** Centrifuge and aliquot serum into a plastic vial.**Forms:**

If not ordering electronically, complete, print, and send a [Renal Diagnostics Test Request](#) (T830) with the specimen.

Specimen Type	Temperature	Time	Special Container
Serum	Refrigerated (preferred)	7 days	
	Frozen	30 days	

**Result Codes:**

Result ID	Reporting Name	Type	Unit	LOINC®
T4	T4 (Thyroxine), Total Only, S	Numeric	mcg/dL	83119-8

LOINC® and CPT codes are provided by the performing laboratory.

**Supplemental Report:**

No

**CPT Code Information:**

84436

**Reference Values:**

Pediatric

0-5 days: 5.0-18.5 mcg/dL

6 days-2 months: 5.4-17.0 mcg/dL

3-11 months: 5.7-16.0 mcg/dL

1-5 years: 6.0-14.7 mcg/dL

6-10 years: 6.0-13.8 mcg/dL

11-19 years: 5.9-13.2 mcg/dL

Adult (> or =20 years): 4.5-11.7 mcg/dL

For International System of Units (SI) conversion for Reference Values, see

[www.mayocliniclabs.com/order-tests/si-unit-conversion.html](http://www.mayocliniclabs.com/order-tests/si-unit-conversion.html)