

Reporting Title: TSH, Sensitive, S
Performing Location: Mayo Clinic Jacksonville Clinical Lab

Ordering Guidance:
This is a standalone test for sensitive thyrotropin (s-TSH; formerly thyroid-stimulating hormone).

If a cascade approach is preferred, order THSCM / Thyroid Function Cascade, Serum, which utilizes a cascaded testing procedure to efficiently evaluate and monitor functional thyroid status. Serum s-TSH is the first-line test and when the s-TSH result is abnormal, appropriate follow-up tests will automatically be performed.

Specimen Requirements:
Collection Container/Tube:
Preferred: Serum gel
Acceptable: Red top
Submission Container/Tube: Plastic vial
Specimen Volume: 0.6 mL

- Collection Instructions:**
- Serum gel tubes should be centrifuged within 2 hours of collection.
 - Red-top tubes should be centrifuged, and the serum aliquoted into a plastic vial within 2 hours of collection.

Forms:
If not ordering electronically, complete, print, and send a [Renal Diagnostics Test Request](#) (T830) with the specimen.

Specimen Type	Temperature	Time	Special Container
Serum	Refrigerated (preferred)	7 days	
	Ambient	7 days	
	Frozen	30 days	

Result Codes:

Result ID	Reporting Name	Type	Unit	LOINC®
STSH	TSH, Sensitive, S	Numeric	mIU/L	11579-0

LOINC® and CPT codes are provided by the performing laboratory.

Supplemental Report:
No

CPT Code Information:
84443

Reference Values:
0-5 days: 0.7-15.2 mIU/L

Test Definition: STSH

Thyroid-Stimulating Hormone-Sensitive
(s-TSH), Serum

-
- 6 days-2 months: 0.7-11.0 mIU/L
 - 3-11 months: 0.7-8.4 mIU/L
 - 1-5 years: 0.7-6.0 mIU/L
 - 6-10 years: 0.6-4.8 mIU/L
 - 11-19 years: 0.5-4.3 mIU/L
 - > or =20 years: 0.3-4.2 mIU/L

For SI unit Reference Values, see <https://www.mayocliniclabs.com/order-tests/si-unit-conversion.html>