

Reporting Title: von Willebrand Factor Ag, P

Performing Location: Mayo Clinic Jacksonville Clinical Lab

Ordering Guidance:

For optimum clinical utility and diagnostic efficiency, this test's results generally must be used together with the results of the von Willebrand factor ristocetin cofactor activity and factor VIII coagulant activity tests. The diagnosis of von Willebrand disease requires a combination of clinical and laboratory information. For a streamlined approach to testing, a panel of tests with reflexive testing and interpretive reporting is recommended. See AVWPR / von Willebrand Disease Profile, Plasma.

Additional Testing Requirements:

VWACT / von Willebrand Factor Activity, Plasma and F8A / Coagulation Factor VIII Activity Assay, Plasma are recommended in conjunction with this test (von Willebrand antigen).

Specimen Requirements:

Specimen Type: Platelet-poor plasma

Collection Container/Tube: Light-blue top (3.2% sodium citrate)

Submission Container/Tube: Plastic vial

Specimen Volume: 1 mL

Collection Instructions:

1. For complete instructions, see [Coagulation Guidelines for Specimen Handling and Processing](#).
2. Centrifuge, transfer all plasma into a plastic vial, and centrifuge plasma again.
3. Aliquot plasma into a plastic vial, leaving 0.25 mL in the bottom of centrifuged vial.
4. Freeze plasma immediately (no longer than 4 hours after collection) at -20 degrees C or, ideally, less than or equal to -40 degrees C.

Additional Information:

1. Double-centrifuged specimen is critical for accurate results as platelet contamination may cause spurious results.
2. Each coagulation assay requested should have its own vial.

Forms:

If not ordering electronically, complete, print, and send a [Coagulation Test Request](#) (T753) with the specimen.

Specimen Type	Temperature	Time	Special Container
Plasma Na Cit	Frozen	14 days	

Result Codes:

Result ID	Reporting Name	Type	Unit	LOINC®
VWAG	von Willebrand Factor Ag, P	Numeric	%	27816-8

LOINC® and CPT codes are provided by the performing laboratory.

Supplemental Report:

No

CPT Code Information:

85246

Reference Values:

55-200%

Note: Individuals of blood group "O" may have lower plasma von Willebrand factor (VWF) antigen than those of other ABO blood groups, such that apparently normal individuals of blood group "O" may have plasma VWF antigen as low as 40% to 50%, whereas the lower limit of the reference range for individuals of other blood groups may be 60% to 70%. Children: Neonates, infants, and children have normal or mildly increased plasma VWF antigen, with respect to the adult reference range.