

# **Test Definition: MPO**

Myeloperoxidase Antibodies, IgG, Serum

Reporting Title: Myeloperoxidase Ab, S

Performing Location: Mayo Clinic Laboratories - Rochester Superior Drive

#### **Additional Testing Requirements:**

To distinguish between microscopic polyangiitis and other forms of anti-neutrophil cytoplasmic antibody-associated vasculitis, also order PR3 / Proteinase 3 Antibodies, IgG, Serum and ANCA / Cytoplasmic Neutrophil Antibodies, Serum; alternatively, VASC / Antineutrophil Cytoplasmic Antibodies Vasculitis Panel, Serum may be ordered instead, which initially tests for myeloperoxidase and proteinase 3 antibodies, with reflex to ANCA when appropriate.

## **Specimen Requirements:**

Supplies: Sarstedt Aliquot Tube 5 mL (T914)

**Collection Container/Tube:** 

**Preferred:** Serum gel **Acceptable:** Red top

Submission Container/Tube: Plastic vial

Specimen Volume: 0.5 mL

Collection Instructions: Centrifuge and aliquot serum into a plastic vial.

#### Forms:

If not ordering electronically, complete, print, and send a Renal Diagnostics Test Request (T830) with the specimen.

Specimen Type	Temperature	Time	Special Container
Serum	Refrigerated (preferred)	21 days	
	Frozen	21 days	

### **Result Codes:**

Result ID	Reporting Name	Туре	Unit	LOINC®
MPO	Myeloperoxidase Ab, S	Numeric	U	48404-8

LOINC® and CPT codes are provided by the performing laboratory.

### **Supplemental Report:**

No

#### **CPT Code Information:**

83516

### **Reference Values:**

<0.4 U (negative)

0.4-0.9 U (equivocal)

> or =1.0 U (positive)

Reference values apply to all ages.