

Overview

Useful For

Investigation of patients with possible Wilson disease

Testing Algorithm

See [Wilson Disease Testing Algorithm](#) in Special Instructions.

Special Instructions

- [Wilson Disease Testing Algorithm](#)

Method Name

Nephelometric Assay

NY State Available

Yes

Specimen

Specimen Type

Serum

Specimen Required

Patient Preparation: Patient should be fasting: 4 hours preferred, nonfasting acceptable.

Collection Container/Tube:

Preferred: Serum gel

Acceptable: Red top

Submission Container/Tube: Plastic vial

Specimen Volume: 1 mL

Forms

[If not ordering electronically, complete, print, and send 1 of the following forms with the specimen:](#)

-[Benign Hematology Test Request](#) (T755).

-[General Request](#) (T239)

-[Gastroenterology and Hepatology Client Test Request](#) (T728)

Specimen Minimum Volume

0.5 mL

Reject Due To

Gross hemolysis	Reject
-----------------	--------

Gross lipemia	Reject
Gross icterus	Reject

Specimen Stability Information

Specimen Type	Temperature	Time	Special Container
Serum	Refrigerated (preferred)	7 days	
	Frozen	30 days	

Clinical and Interpretive

Clinical Information

Ceruloplasmin is a positive acute-phase reactant and a copper-binding protein that accounts for over 95% of serum copper in normal adults. Ceruloplasmin is measured primarily to assist with a diagnosis of Wilson disease. Other indications include Menkes disease, dietary copper insufficiency, and risk of cardiovascular disease.

Wilson disease is a rare inherited disorder of copper transport that results in low serum copper and ceruloplasmin and accumulation of copper in various tissues. The pathological accumulation of copper in the liver, brain, cornea, and kidney causes cirrhosis, neuropsychiatric symptoms, Kayser-Fleischer rings, and hematuria/proteinuria, respectively. See [Wilson Disease Testing Algorithm](#) in Special Instructions for appropriate use of clinical findings, serum biomarkers, genetic tests, and tissue biopsies when working up suspected cases.

Menkes disease is an X-linked disorder in which dietary copper is absorbed from the gastrointestinal tract but cannot be transported, so copper is not available to the liver for incorporation into ceruloplasmin.

Dietary ceruloplasmin deficiency may be due to inadequate dietary copper intake, long-term parenteral nutrition without copper supplementation, malabsorption, penicillamine therapy, or a combination of these.

Reference Values

Males:

0-8 weeks: 7.4-23.7 mg/dL

9 weeks-5 months: 13.5-32.9 mg/dL

6-11 months: 13.7-38.9 mg/dL

12 months-7 years: 21.7-43.3 mg/dL

8-13 years: 20.5-40.2 mg/dL

14-17 years: 17.0-34.8 mg/dL

> or =18 years: 19.0-31.0 mg/dL

Females:

0-8 weeks: 7.4-23.7 mg/dL

9 weeks-5 months: 13.5-32.9 mg/dL

6-11 months: 13.7-38.9 mg/dL

12 months-7 years: 21.7-43.3 mg/dL

8-13 years: 20.5-40.2 mg/dL

14-17 years: 20.8-43.2 mg/dL

> or = 18 years: 20.0-51.0 mg/dL

Interpretation

Low concentrations of ceruloplasmin are consistent with Wilson disease and warrant further investigation.

Values vary considerably from patient to patient and may be in the normal range in some patients with Wilson disease (indicating a different primary defect).

Ceruloplasmin is a positive acute-phase reactant. Increases in serum ceruloplasmin have been reported during pregnancy, in women taking oral contraceptives, in hepatitis, pneumonia, tuberculosis, rheumatoid arthritis, myocardial infarction, various forms of anemia, and many obscure neurological disorders.

Cautions

[Ceruloplasmin is a positive acute-phase reactant; therefore, levels are elevated in cases of inflammation \(as in chronic hepatitis or active infection\). Consequently, ceruloplasmin levels are not always extremely low in patients with Wilson disease.](#)

Birth control pills and pregnancy increase ceruloplasmin levels.

Clinical Reference

1. Tang WH, Wu Y, Hartiala J, et al: Clinical and genetic association of serum ceruloplasmin with cardiovascular risk. *Arterioscler Thromb Vasc Biol* 2012;32:516-522
2. Dadu RT, Dodge R, Nambi V, et al: Ceruloplasmin and heart failure in the Atherosclerosis Risk in Communities study. *Circ Heart Fail* 2013 Sep 1;6(5):936-943
3. Cox DW, Tumer Z, Roberts EA: Copper transport disorders: Wilson's disease and Menkes disease. *Inborn Metabolic Disease*. Edited by J Fernandes, JM Sandubray, F VandenBerghe. Berlin, Heidelberg, New York, Springer-Verlag, 2000, pp 385-391
4. Sontakke AN, More U: Changes in serum ceruloplasmin levels with commonly used methods of contraception. *Indian J Clin Biochem*. 2004 Jan;19(1):102-104

Performance

Method Description

Human ceruloplasmin forms a precipitate with a specific antiserum which is then measured nephelometrically. (Package insert: Ceruloplasmin, Siemens, Newark, DE 08/2018)

PDF Report

No

Day(s) and Time(s) Test Performed

Monday through Saturday; Continuously

Analytic Time

1 day (not reported on Sunday)

Maximum Laboratory Time

3 days

Specimen Retention Time

7 days

Performing Laboratory Location

Rochester

Fees and Codes**Fees**

- Authorized users can sign in to [Test Prices](#) for detailed fee information.
- Clients without access to Test Prices can contact [Customer Service](#) 24 hours a day, seven days a week.
- Prospective clients should contact their Regional Manager. For assistance, contact [Customer Service](#).

Test Classification

This test has been cleared or approved by the U.S. Food and Drug Administration and is used per manufacturer's instructions. Performance characteristics were verified by Mayo Clinic in a manner consistent with CLIA requirements.

CPT Code Information

82390

LOINC® Information

Test ID	Test Order Name	Order LOINC Value
CERS	Ceruloplasmin, S	2064-4

Result ID	Test Result Name	Result LOINC Value
CERS	Ceruloplasmin, S	2064-4