

Overview
Useful For

Only orderable as part of a profile. See WNC / West Nile Virus Antibody, IgG and IgM, Spinal Fluid.

Aids in diagnosis of central nervous system infection with West Nile virus

Method Name

Only orderable as part of a profile. See WNC / West Nile Virus Antibody, IgG and IgM, Spinal Fluid.

NY State Available

Yes

Specimen
Specimen Type

CSF

Specimen Required

Only orderable as part of a profile. See WNC / West Nile Virus Antibody, IgG and IgM, Spinal Fluid.

Specimen Minimum Volume

0.8 mL

Reject Due To

Gross hemolysis	Reject
-----------------	--------

Specimen Stability Information

Specimen Type	Temperature	Time	Special Container
CSF	Refrigerated (preferred)	7 days	
	Frozen	30 days	

Clinical and Interpretive
Clinical Information

West Nile virus (WNV) is a mosquito-borne flavivirus (single-stranded RNA) that primarily infects birds but can also infect humans and horses. WNV was first isolated in 1937 from an infected person in the West Nile district of Uganda. Until the viral infection was recognized in 1999 in birds in New York City, WNV was found only in the Eastern Hemisphere, with wide distribution in Africa, Asia, the Middle East, and Europe.(1-3) Most recently, in 2012, a total of 5,674 cases of WNV were reported to the CDC, among which 2,873 (51%) were classified as neuroinvasive disease (eg, meningitis or encephalitis) and 286 (5%) cases resulted in death.(2)

Most people who are infected with WNV will not develop clinical signs of illness. It is estimated that about 20% of those who become infected will develop West Nile fever with mild symptoms, including fever, headache, myalgia,

and occasionally a skin rash on the trunk of the body. Case fatality rates among patients hospitalized during recent outbreaks have ranged from 4% to 14%. Advanced age is the most important risk factor for death, and patients older than 70 years of age are at particularly high risk.(1)

Laboratory diagnosis is best achieved by demonstration of specific IgG and IgM class antibodies in serum specimens. PCR (LCWNV / West Nile Virus, Molecular Detection, PCR) can detect WNV RNA in specimens from patients with recent WNV infection (ie, 3-5 days following infection) when specific antibodies to the virus are not yet present. However, the likelihood of detection is relatively low as the sensitivity of PCR detection is approximately 55% in cerebrospinal fluid and approximately 10% in blood, from patients with known WNV infection.

Reference Values

Only orderable as part of a profile. See WNC / West Nile Virus Antibody, IgG and IgM, Spinal Fluid.

Interpretation

A positive result may indicate recent or past central nervous system (CNS) infection with West Nile virus (WNV). Clinical correlation is necessary.

This assay is unable to distinguish between intrathecal antibody synthesis and serum antibodies introduced into the cerebrospinal fluid at the time of lumbar puncture or from a breakdown in the blood-brain barrier. Positive results should be interpreted with other laboratory and clinical data prior to a diagnosis of CNS infection.

Cautions

Test results should be used in conjunction with clinical evaluation, exposure history and other available diagnostic procedures.

The significance of negative test results in immunosuppressed patients is uncertain.

False-negative results due to competition by high levels of IgG, while theoretically possible, have not been observed.

False-positive results may occur in patients infected with other flaviviruses, including dengue virus, St. Louis virus, and Zika virus and in persons previously infected with West Nile virus (WNV).

Because closely related arboviruses exhibit serologic cross-reactivity, it sometimes may be epidemiologically important to attempt to pinpoint the infecting virus by conducting plaque reduction neutralization tests (PRNT) using an appropriate battery of closely related viruses. Such testing is available through the CDC and select public health laboratories.

WNV antibody results for cerebrospinal fluid (CSF) should be interpreted with caution. Complicating factors include low antibody levels found in CSF, passive transfer of antibody from blood, and contamination via a traumatic lumbar puncture.

Clinical Reference

1. [Petersen LR, Marafin AA: West Nile Virus: a primer for the clinician. Ann Intern Med 2002;137:173-179](#)

2. MMWR: West Nile Virus and Other Arboviral Diseases-United States, 2012. 2013;62(25):513-517

3. Brinton MA: The molecular biology of West Nile Virus: a new invader of the western hemisphere. Ann Rev Microbiol 2002;56:371-402

4. Centers for Disease Control and Prevention (CDC). Provisional Surveillance Summary of the West Nile Virus epidemic. United States, January-November 2002. MMWR Morb Mortal Wkly Rep 2002;51(50):1129-1133

5. Centers for Disease Control and Prevention (CDC). Investigations of West Nile Virus infections in recipients of blood transfusions. MMWR Morb Mortal Wkly Rep 2002;51(43):973-974

Performance

Method Description

Polystyrene microwells are coated with recombinant West Nile virus (WNV) antigen. Diluted serum specimens and controls are incubated in the wells to allow specific antibody present in the specimens to react with the antigen. Nonspecific reactants are removed by washing, and peroxidase-conjugated antihuman IgG is added and reacts with specific IgG. Excess conjugate is removed by washing. Enzyme substrate and chromogen are added, and the color is allowed to develop. After adding the Stop reagent, the resultant color change is quantified by a spectrophotometric reading of optical density (OD). Specimen OD readings are compared with reference cutoff readings to determine results. (Package insert: Flavivirus [West Nile] ELISA IgG. Focus Technologies, Cypress, CA 10/16/2012)

PDF Report

No

Day(s) Performed

Monday, Wednesday, Friday; 9 a.m.

Report Available

Same day/1 day

Performing Laboratory Location

Rochester

Fees and Codes

Fees

- Authorized users can sign in to [Test Prices](#) for detailed fee information.
- Clients without access to Test Prices can contact [Customer Service](#) 24 hours a day, seven days a week.
- Prospective clients should contact their Regional Manager. For assistance, contact [Customer Service](#).

Test Classification

This test has been modified from the manufacturer's instructions. Its performance characteristics were determined by Mayo Clinic in a manner consistent with CLIA requirements. This test has not been cleared or approved by the US Food and Drug Administration.

CPT Code Information

86789

LOINC® Information

Test ID	Test Order Name	Order LOINC Value
WNGC	West Nile Virus Ab, IgG, CSF	41236-1

Result ID	Test Result Name	Result LOINC Value
WNGC	West Nile Virus Ab, IgG, CSF	77953-8