

Overview

Useful For

Screening for recent or past exposure to *Mycoplasma pneumoniae*

Profile Information

Test ID	Reporting Name	Available Separately	Always Performed
MYCOG	M. pneumoniae Ab, IgG, S	No	Yes
MYCOM	M. pneumoniae Ab, IgM, S	No	Yes
MYCON	M. pneumoniae Ab Interpretation	No	Yes

Reflex Tests

Test ID	Reporting Name	Available Separately	Always Performed
MMYCO	M. pneumoniae Ab, IgM, S by IFA	No	No

Testing Algorithm

If *Mycoplasma pneumoniae*, IgM is reactive or equivocal, then *M pneumoniae* IgM by IFA will be performed at an additional charge.

Method Name

[MYCOG, MYCOM: Enzyme Immunoassay \(EIA\)](#)

MMYCO: Indirect Immunofluorescence Assay (IFA)

NY State Available

Yes

Specimen

Specimen Type

Serum

Advisory Information

Detection of IgM or IgG class antibodies to *Mycoplasma pneumoniae* provides exposure information. The preferred method of diagnosis of acute *M pneumoniae* infection is by molecular detection: MPRP / *Mycoplasma pneumoniae*, Molecular Detection, PCR.

Specimen Required

[Collection Container/Tube:](#)

Preferred: Serum gel

Acceptable: Red top

Submission Container/Tube: Plastic vial

Specimen Volume: 1 mL

Forms

If not ordering electronically, complete, print, and send a [Microbiology Test Request](#) (T244) with the specimen.

Specimen Minimum Volume

0.5 mL

Reject Due To

Gross hemolysis	Reject
Gross lipemia	Reject
Other	Heat inactivated specimen

Specimen Stability Information

Specimen Type	Temperature	Time	Special Container
Serum	Refrigerated (preferred)	14 days	
	Frozen	14 days	

Clinical and Interpretive

Clinical Information

Mycoplasma pneumoniae is a small bacterium transmitted via organism-containing droplets. It is a cause of upper respiratory infection, pharyngitis, and tracheobronchitis, particularly in children, and has been associated with approximately 20% of cases of community-acquired pneumonia. Central nervous system and cardiac manifestations are probably the most frequent extrapulmonary complications of infections due to *M pneumoniae*. The disease is usually self-limited, although severe disease has been reported in immunocompromised patients.

Identification of *M pneumoniae* by culture-based methods is time consuming and insensitive. Serology-based assays for *M pneumoniae* have several drawbacks. The development of IgM antibodies takes approximately 1 week and the IgM response in adults may be variable or it may be decreased in immunosuppressed individuals. Confirmation of the disease is dependent on the observation of a 4-fold rise in IgG antibody titers between acute and convalescent specimens, several weeks following the initial onset of illness, providing clinical utility only for retrospective testing. Real-time PCR offers a rapid and sensitive option for detection of *M pneumoniae* DNA from clinical specimens allows for diagnosis of acute or current infection.

Reference Values

IgG: negative

IgM: negative

IgM by IFA: negative

Interpretation

IgG ELISA Result	IgM ELISA Result	Interpretation
Positive	Negative	Results suggest past exposure.
Positive	Reactive	Prior exposure to <i>M pneumoniae</i> detected. Confirmatory testing for IgM to <i>M pneumonia</i> will be performed by an immunofluorescence assay.
	Equivocal	
Negative	Negative	No antibodies to <i>M pneumoniae</i> detected. Acute infection cannot be ruled out as antibody levels may be below the limit of detection. If clinically indicated, a second serum should be submitted in 14 to 21 days.
	Reactive	
Negative	Equivocal	No prior exposure to <i>Mycoplasma pneumoniae</i> . Confirmatory testing for IgM to <i>M pneumonia</i> will be performed by an immunofluorescence assay..
	Reactive	
Equivocal	Negative	Recommend follow-up testing in 10 to 14 days if clinically indicated.
	Reactive	Confirmatory testing for IgM to <i>M pneumonia</i> will be performed by an immunofluorescence assay.
	Equivocal	

Cautions

A diagnosis of *Mycoplasma pneumoniae* infection should not be solely based on results of serologic testing for this agent. Test results should be interpreted in conjunction with clinical evaluation and the results of other diagnostic procedures (eg, molecular detection).

The continued presence or absence of antibodies cannot be used to determine the success or failure of therapy.

Testing should not be performed as a screening procedure for the general population. Testing should only be done when clinical evidence suggests the diagnosis of *M pneumoniae*-associated disease.

The performance of this test has not been established on neonates and immunocompromised patients.

Performance of the IgM assay has not been tested with specimens known to be positive for antibodies to organisms that are known to be associated with lower respiratory illness (ie, influenza A and B, cytomegalovirus, *Chlamydomphila pneumoniae*, parainfluenza), and closely related serovars known to cross-react with *M pneumoniae*, such as *M genitalium* and *M hominis*, as well as various *Ureaplasma* species. Cross-reactivity studies with such organisms have not been performed with this assay.

The IgG removal system included with the IgM test system has been shown to functionally remove the IgG from specimens containing total IgG levels ranging from 300 to 600 mg/mL. The effectiveness of this removal system at IgG levels exceeding 600 mg/mL has not been established.

Clinical Reference

Smith T: *Mycoplasma pneumoniae* infections: diagnosis based on immunofluorescence titer of IgG and IgM antibodies. Mayo Clin Proc 1986;61:830-831

Performance

Method Description

IgG:

Diluted sera are incubated in antigen-coated microwells. Any antigen-specific antibody in the samples will bind to the immobilized antigen. The plate is washed to remove unbound antibody and other serum components. Peroxidase conjugated goat-antihuman IgG is added to the wells and incubated. The conjugate will react with the IgG antibody/antigen on the solid phase. The wells are washed to remove unreacted conjugate. The microwells containing immobilized conjugate are incubated with peroxidase substrate solution. Hydrolysis of the substrate by peroxidase produces a color change. After a period of time the reaction is stopped by the addition of diluted acid, and the color changes are measured photometrically. The color intensity of the solution depends on the antibody concentration in the serum sample. (Package insert: *M pneumoniae* IgG Test System. Zeus Scientific Inc., Branchburg, NJ. Revision Date 3/22/2016)

IgM EIA:

Test sera are diluted with the sample diluent provided. The sample diluent contains antihuman IgG that precipitates and removes IgG and rheumatoid factor from the sample, leaving IgM free to react with immobilized antigen. Diluted sera are incubated in antigen-coated microwells. Any antigen-specific antibody in the samples will bind to the immobilized antigen. The plate is washed to remove unbound antibody and other serum components. Peroxidase-conjugated goat-antihuman IgM (chain specific) is added to the wells and incubated. The conjugate will react with the IgM antibody/antigen on the solid phase. The wells are washed to remove unbound conjugate. The microwells containing immobilized conjugate are incubated with peroxidase substrate solution. Hydrolysis of the substrate by peroxidase produces a color change. After a period of time the reaction is stopped by the addition of diluted acid, and the color changes are measured photometrically. The color intensity of the solution depends on the antibody concentration in the serum sample. (Package insert: *M pneumoniae* IgM Test System. Zeus Scientific, Inc., Branchburg, NJ. Revision Date 9/22/2016)

IgM Immunofluorescence Assay (IFA):

Mycoplasma pneumoniae antigenic substrate is fixed onto microscope slide wells. Serum that has been pretreated to remove IgG antibodies is incubated with the substrate. If IgM antibody to *M pneumoniae* is present, it will bind to the substrate. Fluorescein-labeled antihuman-IgM conjugate is added to the slide wells and the slide is incubated. If antibody is present, it can be observed as a characteristic positive, bright, apple-green fluorescent reaction when the slide is read on a fluorescence microscope. (Package insert: *Mycoplasma pneumoniae* IgM IFA Antibody Test System. Zeus Scientific, Inc., Raritan, NJ 2004)

PDF Report

No

Day(s) and Time(s) Test Performed

Monday through Friday; 9 a.m.

Analytic Time

Same day/1 day

Maximum Laboratory Time

3 days

Specimen Retention Time

14 days

Performing Laboratory Location

Rochester

Fees and Codes**Fees**

- Authorized users can sign in to [Test Prices](#) for detailed fee information.
- Clients without access to Test Prices can contact [Customer Service](#) 24 hours a day, seven days a week.
- Prospective clients should contact their Regional Manager. For assistance, contact [Customer Service](#).

Test Classification

This test has been modified from the manufacturer's instructions. Its performance characteristics were determined by Mayo Clinic in a manner consistent with CLIA requirements. This test has not been cleared or approved by the U.S. Food and Drug Administration.

CPT Code Information86738 x 2-*Mycoplasma pneumoniae* by EIA86738-*Mycoplasma pneumoniae* by indirect IFA (if appropriate)**LOINC® Information**

Test ID	Test Order Name	Order LOINC Value
MYCO	M. pneumoniae Ab, IgG and IgM, S	58733-7

Result ID	Test Result Name	Result LOINC Value
MYCOG	M. pneumoniae Ab, IgG, S	45224-3
MYCOM	M. pneumoniae Ab, IgM, S	5257-1
MYCON	M. pneumoniae Ab Interpretation	69048-7