

Overview

Useful For

Identifying amphetamines (and methamphetamines), opiates, as well as metabolites of cocaine and marijuana in meconium specimens

Reflex Tests

Test ID	Reporting Name	Available Separately	Always Performed
AMPHM	Amphetamines, Confirmation, M	Yes	No
COKEM	Cocaine and Metabolites, Confirm, M	Yes	No
OPATM	Opiate Confirmation, M	Yes	No
THCM	Carboxy-THC Confirmation, M	Yes	No

Testing Algorithm

Testing begins with immunoassay screen. Positives are confirmed and quantitated by liquid-chromatography/tandem-mass spectrometry (LC-MS/MS) at an additional charge.

Method Name

Enzyme-Linked Immunosorbent Assay (ELISA)

NY State Available

Yes

Specimen

Specimen Type

Meconium

Ordering Guidance

For chain-of-custody testing, order DSM4X / Drugs of Abuse Screen 4, Chain of Custody, Meconium.

Specimen Required

Supplies: Stool container, Small (Random), 4 oz (T288)

Container/Tube: Stool container (T288)

Specimen Volume: 1 g (approximately 1 teaspoon)

Collection Instructions: Collect entire random meconium specimen.

Forms

If not ordering electronically, complete, print, and send a [Therapeutics Test Request](#) (T831) with the specimen.

Specimen Minimum Volume

0.45 g (approximately 0.5 teaspoon)

Reject Due To

Other	Grossly bloody reject, Pink OK Stool Diapers
-------	--

Specimen Stability Information

Specimen Type	Temperature	Time	Special Container
Meconium	Frozen (preferred)	14 days	
	Refrigerated	24 hours	

Clinical and Interpretive
Clinical Information

Illicit drug use during pregnancy is a major social and medical issue. Drug abuse during pregnancy is associated with significant perinatal complications, which include a high incidence of stillbirths, meconium-stained fluid, premature rupture of the membranes, maternal hemorrhage (abruption placenta or placenta praevia), and fetal distress.(1) In the neonate, the mortality rate, as well as morbidity (eg, asphyxia, prematurity, low birthweight, hyaline membrane distress, infections, aspiration pneumonia, cerebral infarction, abnormal heart rate and breathing problems, drug withdrawal) are increased.(1)

The disposition of drug in meconium is not well understood. The proposed mechanism is that the fetus excretes drug into bile and amniotic fluid. Drug accumulates in meconium either by direct deposit from bile or through swallowing of amniotic fluid.(2) The first evidence of meconium in the fetal intestine appears at approximately the tenth to twelfth week of gestation, and slowly moves into the colon by the sixteenth week of gestation.(3) Therefore, the presence of drugs in meconium has been proposed to be indicative of in utero drug exposure during the final 4 to 5 months of pregnancy, a longer historical measure than is possible by urinalysis.(2)

Reference Values

Negative

Positives are reported with a quantitative LC-MS/MS result.

Cutoff concentrations

Amphetamines by ELISA: 100 ng/g

Methamphetamine by ELISA: 100 ng/g

Benzoylcegonine (cocaine metabolite) by ELISA: 100 ng/g

Opiates by ELISA: 100 ng/g

Tetrahydrocannabinol carboxylic acid (marijuana metabolite) by ELISA: 20 ng/g

Interpretation

A positive result indicates that the baby was exposed to the drugs indicated.

Cautions

No significant cautionary statements

Clinical Reference

1. Ostrea EM Jr: Understanding drug testing in the neonate and the role of meconium analysis. J Perinat Neonatal Nurs 2001 Mar;14(4):61-82; quiz 105-106
2. Ostrea EM Jr, Brady MJ, Parks PM, et al: Drug screening of meconium in infants of drug-dependent mothers: an alternative to urine testing. J Pediatr 1989 Sep;115(3):474-477
3. Ahanya SN, Lakshmanan J, Morgan BL, Ross MG: Meconium passage in utero mechanisms, consequences, and management. Obstet Gynecol Surv 2005 Jan;60(1):45-56; quiz 73-74

Performance**Method Description**

Meconium specimen is homogenized in a buffer. The homogenate is analyzed by enzyme-linked immunosorbent assay (ELISA) to detect the presence of drug. If drug presence is indicated by a positive result of ELISA, liquid chromatography-tandem mass spectrometry analysis is performed to verify the presence of the drug. (Unpublished Mayo method)

PDF Report

No

Day(s) Performed

Monday through Saturday

Report Available

1 to 3 days

Specimen Retention Time

2 weeks

Performing Laboratory Location

Rochester

Fees and Codes**Fees**

- Authorized users can sign in to [Test Prices](#) for detailed fee information.
- Clients without access to Test Prices can contact [Customer Service](#) 24 hours a day, seven days a week.
- Prospective clients should contact their Regional Manager. For assistance, contact [Customer Service](#).

Test Classification

This test was developed, and its performance characteristics determined by Mayo Clinic in a manner consistent with CLIA requirements. This test has not been cleared or approved by the US Food and Drug Administration.

CPT Code Information

80307

See individual reflex tests for appropriate CPT codes

LOINC® Information

Test ID	Test Order Name	Order LOINC Value
DASM4	Drugs of Abuse Screen, Meconium 4	49046-6

Result ID	Test Result Name	Result LOINC Value
32078	Amphetamine	26895-3
32080	Methamphetamine	27289-8
32082	Cocaine	26956-3
32084	Opiate	29158-3
32086	Tetrahydrocannabinol	26893-8
32087	Chain of Custody	77202-0