

---

**Overview****Useful For**

Monitoring serum concentration of lamotrigine

Assessing compliance

Adjusting lamotrigine dose in patients receiving other anticonvulsant drugs which interact pharmacokinetically with lamotrigine

**Method Name**

Liquid Chromatography-Tandem Mass Spectrometry (LC-MS/MS)

**NY State Available**

Yes

**Specimen****Specimen Type**

Serum

**Specimen Required****Collection Container/Tube:**

**Preferred:** Red top

**Acceptable:** Serum gel

**Submission Container/Tube:** Plastic vial

**Specimen Volume:** 1 mL

**Collection Instructions:**

1. Drawblood immediately before next scheduled dose.
2. For sustained-release formulations ONLY,draw blood aminimum of 12 hours after last dose.
3. [Centrifuge within 2 hours of collection.](#)
4. Aliquot serum into plastic vial.

**Forms**

If not ordering electronically, complete, print, and send 1 of the following forms with the specimen:

-[Neurology Specialty Testing Client Test Request](#) (T732)

-[General Request](#) (T239)

---

[-Therapeutics Test Request](#) (T831)

### Specimen Minimum Volume

0.5 mL

### Reject Due To

Gross hemolysis	OK
Gross lipemia	OK
Gross icterus	OK

### Specimen Stability Information

Specimen Type	Temperature	Time	Special Container
Serum	Refrigerated (preferred)	28 days	
	Ambient	28 days	
	Frozen	28 days	

## Clinical and Interpretive

### Clinical Information

Lamotrigine (Lamictal) is approved for therapy of bipolar I disorder and a wide variety of seizure disorders including Lennox-Gastaut syndrome, primary generalized tonic-clonic seizures, and partial seizures. Its many off-label uses include treatment of migraine, trigeminal neuralgia, and treatment-refractory depression. Lamotrigine inhibits glutamate release (an excitatory amino acid) and voltage-sensitive sodium channels to stabilize neuronal membranes; it also weakly inhibits the 5-HT<sub>3</sub> (serotonin) receptor.

Lamotrigine oral bioavailability is very high (approximately 98%). The drug is metabolized by glucuronic acid conjugation to inactive metabolites. The half-life is 25 to 33 hours in adults but decreases with concurrent use of phenytoin or carbamazepine (13-14 hours) and increases with concomitant valproic acid therapy (59-70 hours), renal dysfunction, or hepatic impairment. The therapeutic range is relatively wide, 2.5 to 15 mcg/mL for most individuals. Common adverse effects are dizziness, ataxia, blurred or double vision, nausea, or vomiting.

### Reference Values

Patients receiving therapeutic doses usually have lamotrigine concentrations of 2.5-15.0 mcg/mL.

### Interpretation

The serum concentration should be interpreted in the context of the patient's clinical response and may provide useful information in patients showing poor response, noncompliance, or adverse effects, particularly when lamotrigine is co-administered with other anticonvulsant drugs.

While most patients show response to the drug when the trough concentration is in the range of 2.5 to 15.0 mcg/mL, and show signs of toxicity when the peak serum concentration is greater than 20 mcg/mL, some patients can tolerate peak concentrations as high as 70 mcg/mL.

### Cautions

Serum separator tube acceptable but serum should be removed from gel within 24 hours of collection.

---

## Clinical Reference

- [1. Johannessen SI, Battino D, Berry DJ, et al: Therapeutic drug monitoring of the newer antiepileptic drugs. Ther Drug Monit. 2003 Jun;25\(3\):347-363](#)
- Johannessen SI, Landmark CJ: Value of therapeutic drug monitoring in epilepsy. Expert Rev Neurother. 2008 Jun;8(6):929-939
- Johannessen SI, Tomson T: Pharmacokinetic variability of newer antiepileptic drugs: when is monitoring needed? Clin Pharmacokinet. 2006;45(11):1061-1075
- Physician's Desk Reference. 71st ed. Thomson PDR; 2017
- Hardman JG, Limbird LE, Gilman AG, eds. Goodman and Gilman's The Pharmacological Basis of Therapeutics. 10th ed. McGraw-Hill Book Company; 2001
- Hiemke C, Bergemann N, Clement HW, et al: Consensus guidelines for therapeutic drug monitoring in neuropsychopharmacology: Update 2017. Pharmacopsychiatry. 2018 Jan;51(1-02):9-62

## Performance

### Method Description

Samples are diluted and extracted online extraction by high turbulence liquid chromatography, with detection by tandem mass spectrometry.(Unpublished Mayo method)

### PDF Report

No

### Day(s) Performed

Monday through Sunday

### Report Available

Same day/1 to 2 days

### Specimen Retention Time

14 days

### Performing Laboratory Location

Rochester

## Fees and Codes

### Fees

- Authorized users can sign in to [Test Prices](#) for detailed fee information.
- Clients without access to Test Prices can contact [Customer Service](#) 24 hours a day, seven days a week.
- Prospective clients should contact their Regional Manager. For assistance, contact [Customer Service](#).

### Test Classification

This test was developed, and its performance characteristics determined by Mayo Clinic in a manner consistent with CLIA requirements. This test has not been cleared or approved by the US Food and Drug Administration.

**CPT Code Information**

80175

**LOINC® Information**

Test ID	Test Order Name	Order LOINC Value
LAMO	Lamotrigine, S	6948-4

Result ID	Test Result Name	Result LOINC Value
80999	Lamotrigine, S	6948-4