

Overview

Useful For

Monitoring therapy to ensure drug absorption, clearance, or compliance

Method Name

Liquid-Chromatography Mass Spectrometry (LC-MS/MS)

NY State Available

Yes

Specimen

Specimen Type

Serum Red

Specimen Required

Collection Container/Tube: Red top (gel tubes/SST are **not** acceptable)

Submission Container/Tube: Plastic vial

Specimen Volume: 1 mL

Collection Instructions:

1. Serum for a peak level should be collected 60 minutes after dose.
2. Centrifuge within 2 hours of collection.

Forms

If not ordering electronically, complete, print, and send a [Therapeutics Test Request](#) (T831) with the specimen.

Specimen Minimum Volume

0.2 mL

Reject Due To

Gross hemolysis	OK
Gross lipemia	OK
Gross icterus	OK

Specimen Stability Information

Specimen Type	Temperature	Time	Special Container
Serum Red	Refrigerated (preferred)	28 days	
	Ambient	28 days	
	Frozen	28 days	

Clinical and Interpretive

Clinical Information

Sulfamethoxazole is a sulfonamide antibiotic that is administered in conjunction with another antibacterial, trimethoprim. These agents are used to treat a variety of infections including methicillin-resistant *Staphylococcus aureus* and for prophylaxis in immunosuppressed patients, such as HIV-positive individuals. Therapeutic drug monitoring is not commonly performed unless there are concerns about adequate absorption, clearance, or compliance. Monitoring of sulfamethoxazole is indicated only when prolonged (>3 months) therapy is required.

Sulfamethoxazole is absorbed readily after oral administration, with peak serum concentration occurring 2 to 3 hours after an oral dose. Its average elimination half-life is 6 to 10 hours. Toxicity includes crystalluria with resultant calculi and renal disease. Toxicity is due to a high concentration of acetylated, relatively insoluble forms of the drug. Excess fluid should be taken with sulfamethoxazole to avoid formation of urine sulfonamide crystals.

Reference Values

>50 mcg/mL

Interpretation

Serum drug concentrations should be interpreted with respect to the minimum inhibitory concentration (MIC) of targeted organisms. Most patients will display peak steady-state serum concentrations greater than 50 mcg/mL when collected at least 1 hour after an oral dose. Targets concentrations may be higher, depending on the intent of therapy.

For *Pneumocystis carinii* pneumonia (PCP pneumonia), peak concentrations: 100-150 mcg/mL

Toxicity: >200 mcg/mL

Toxicity (formation of urinary crystals) associated with sulfamethoxazole occurs with prolonged exposure to serum concentrations greater than 125 mcg/mL.

Cautions

Specimens collected in serum gel tubes are not acceptable, as the drug can absorb on the gel and lead to falsely decreased concentrations.

Clinical Reference

1. Hughes WT, Feldman S, Chaudhary SC, Ossi MJ, et al: Comparison of pentamidine isethionate and trimethoprim-sulfamethoxazole in the treatment of *Pneumocystis carinii* pneumonia. *J Pediatr* 1978;92(2):285-291
2. Dao BD, Barreto JN, Wolf RC, Dierkhising RA, et al: Serum peak sulfamethoxazole concentrations demonstrate difficulty in achieving a target range: a retrospective cohort study. *Curr Ther Res Clin Exp* 2014;76:104-109
3. Young T, Oliphant C, Araoyinbo I, Volmink J: Co-trimoxazole prophylaxis in HIV: the evidence. *S Afr Med J* 2008;98(4):258-259
4. Avdic E, Cosgrove SE: Management and control strategies for community-associated methicillin-resistant *Staphylococcus aureus*. *Expert Opin Pharmacother* 2008;9(9):1463-1479
5. Kamme C, Melander A, Nilsson N: Serum and saliva concentrations of sulfamethoxazole and trimethoprim in

adults in children: relation between saliva concentrations and in vitro activity against nasopharyngeal pathogens. Scand J Infect Dis 1983;15:107-113

6. Goodman, Gilman's: The Pharmacological Basis of Therapeutics. 13th edition. Edited by LL Brunton, R Hilal-Dandan, BC Knollmann. McGraw-Hill Publishing, 2018

Performance

Method Description

Samples are extracted with analyte detection by mass spectrometry.(Unpublished Mayo method)

PDF Report

No

Day(s) and Time(s) Test Performed

Monday, Thursday; 1 p.m.

Analytic Time

2 days

Maximum Laboratory Time

5 days

Specimen Retention Time

14 days

Performing Laboratory Location

Rochester

Fees and Codes

Fees

- Authorized users can sign in to [Test Prices](#) for detailed fee information.
- Clients without access to Test Prices can contact [Customer Service](#) 24 hours a day, seven days a week.
- Prospective clients should contact their Regional Manager. For assistance, contact [Customer Service](#).

Test Classification

This test was developed and its performance characteristics determined by Mayo Clinic in a manner consistent with CLIA requirements. This test has not been cleared or approved by the U.S. Food and Drug Administration.

CPT Code Information

80299

LOINC® Information

Test ID	Test Order Name	Order LOINC Value
SFZ	Sulfamethoxazole, S	10342-4

Result ID	Test Result Name	Result LOINC Value
8238	Sulfamethoxazole, S	10342-4