

Overview

Useful For

Detecting lead exposure using hair specimens

Special Instructions

- [Collecting Hair and Nails for Metals Testing](#)

Method Name

InductivelyCoupledPlasma-MassSpectrometry(ICP-MS)

NY State Available

No

Specimen

Specimen Type

Hair

Necessary Information

Indicate source of hair (axillary, head, or pubic), if known

Specimen Required

Supplies: Hair and Nails Collection Kit (T565)

Specimen Volume: 0.2 g

Collection Instructions: Prepare and transport specimen per the instructions in the kit or see [Collecting Hair and Nails for Metals Testing](#) in Special Instructions.

Specimen Minimum Volume

0.05 g

Reject Due To

All specimens will be evaluated at Mayo Clinic Laboratories for test suitability.

Specimen Stability Information

Specimen Type	Temperature	Time	Special Container
Hair	Ambient (preferred)		
	Frozen		
	Refrigerated		

Clinical and Interpretive

Clinical Information

Hair analysis for lead can be used to corroborate blood analysis or to document past lead exposure. If the hair is collected and segmented in a time sequence (based on length from root), the approximate time of exposure can be assessed.

Reference Values

<4.0 mcg/g of hair

Reference values apply to all ages.

Interpretation

Normal hair lead content is below 4.0 mcg/g. While hair lead content above 10.0 mcg/g may indicate significant lead exposure, hair is also subject to potential external contamination with environmental lead and contaminants in artificial hair treatments (e.g. dyeing, bleaching, or permanents). Ultimately, the hair lead content needs to be interpreted in addition to the overall clinical scenario including symptoms, physical findings, and other diagnostic results when determining further actions.

Cautions

Blood lead analysis has the strongest correlation with toxicity.

Clinical Reference

1. Strumylaite L, Ryselis S, Kregzdyte R: Content of lead in human hair from people exposed to lead. *Int J Hyg Environ Health*. 2004;207:345-351
2. Barbosa F, Tanus-Santos J, Gerlach R, Parsons P: A Critical review of biomarkers used for monitoring human exposure to lead: advantages, limitations, and future needs. *Environ Health Perspect*. 2005;113:1669-1674
3. Sanna E, Liguori A, Palmes L, et al: Blood and hair lead levels in boys and girls living in two Sardinian towns at different risks of lead pollution. *Ecotoxicol Environ Saf*. 2003;55:293-299
4. DiPietro ES, Phillips DL, Paschal DC, Neese JW. Determination of trace elements in human hair. *Biol Trace Elem Res* 1989;22:83-100
5. Strathmann FG, Blum LM: Toxic elements. In: Nader R, Horwath AR, Wittwer CT, eds. *Tietz Textbook of Clinical Chemistry and Molecular Diagnostics*. 6th ed. Elsevier; 2018:chap 42

Performance

Method Description

Lead in hair is analyzed by inductively coupled plasma-mass spectrometry (ICP-MS) in kinetic energy discrimination (KED) mode using gallium, iridium, and lutetium as internal standards, and a salt matrix calibration.(Unpublished Mayo method)

PDF Report

No

Day(s) and Time(s) Test Performed

Tuesday; 3 p.m.

Analytic Time

2 days

Maximum Laboratory Time

7 days

Specimen Retention Time

14 days

Performing Laboratory Location

Rochester

Fees and Codes**Fees**

- Authorized users can sign in to [Test Prices](#) for detailed fee information.
- Clients without access to Test Prices can contact [Customer Service](#) 24 hours a day, seven days a week.
- Prospective clients should contact their Regional Manager. For assistance, contact [Customer Service](#).

Test Classification

This test was developed and its performance characteristics determined by Mayo Clinic in a manner consistent with CLIA requirements. This test has not been cleared or approved by the U.S. Food and Drug Administration.

CPT Code Information

83655

LOINC® Information

Test ID	Test Order Name	Order LOINC Value
PBHA	Lead, Hair	5673-9

Result ID	Test Result Name	Result LOINC Value
31898	Lead, Hair	5673-9
PBHSC	Specimen Source	31208-2