

Overview

Useful For

Monitoring toxicity in overdose cases

Method Name

Photometric

NY State Available

Yes

Specimen

Specimen Type

Serum

Specimen Required

Collection Container/Tube:

Preferred: Serum gel

Acceptable: Red top

Submission Container/Tube: Plastic vial

Specimen Volume: 0.5 mL

Collection Instructions:

1. Serum gel tubes should be centrifuged within 2 hours of collection.
2. Red-top tubes should be centrifuged, and the serum aliquoted into a plastic vial within 2 hours of collection.

Specimen Minimum Volume

0.25 mL

Reject Due To

Gross hemolysis	Reject
-----------------	--------

Specimen Stability Information

Specimen Type	Temperature	Time	Special Container
Serum	Refrigerated (preferred)	7 days	
	Ambient	24 hours	
	Frozen	180 days	

Clinical & Interpretive

Clinical Information

Acetaminophen (found in Anacin-3, Comtrex, Contac, Datril, Dristan, Excedrin, Nyquil, Sinutab, Tempera, Tylenol, Vanquish, and many others) is an analgesic, antipyretic drug lacking significant anti-inflammatory activity. It is metabolized by the liver with a normal elimination half-life of less than 4 hours. In normal therapeutic doses, a minor metabolite, possessing electrophilic alkylating activity, readily reacts with glutathione in the liver to yield a detoxified product. In overdose situations, liver glutathione is consumed and the toxic metabolite (postulated metabolite: benzoquinone) reacts with cellular proteins resulting in hepatotoxicity, characterized by centrilobular necrosis and possible death if untreated. *N*-acetylcysteine can substitute for glutathione and serves as an antidote.

Serum concentration and half-life are the only way to assess degree of intoxication in early stages since other liver function studies (eg, bilirubin, liver function enzymes) will not show clinically significant increases until after tissue damage has occurred, at which point therapy is ineffective.

Reference Values

Therapeutic: 10-30 mcg/mL

Toxic: >150 mcg/mL 4 hours after dose

Interpretation

The normal half-life is less than 4 hours, while the toxic half-life is greater than 4 hours.

The toxic level is dependent on half-life. When the half-life is 4 hours, hepatotoxicity generally will not occur unless the concentration is above 150 mcg/mL. The level at which toxicity occurs decreases with increasing half-life until it is encountered at values as low as 50 mcg/mL when the half-life reaches 12 hours.

For half-life determination, draw 2 specimens at least 4 hours apart and note the exact time of each draw. Half-life can be calculated from the concentrations and the time interval.

Cautions

First specimen should be drawn no sooner than 2 hours postingestion.

Clinical Reference

Rumack BH, Peterson RG: Acetaminophen overdose: incidence, diagnosis, and management in 416 patients. *Pediatrics* Nov 1978;62:898-903

Performance

Method Description

The assay is based on a homogeneous enzyme immunoassay technique used for the quantitative analysis of acetaminophen in human serum or plasma. The assay is based on competition between drug in the sample and drug labeled with the enzyme glucose-6-phosphate dehydrogenase (G6PDH) for antibody binding sites. Enzyme activity decreases upon binding to the antibody, so the drug concentration in the sample can be measured in terms of enzyme

activity. Active enzyme converts oxidized nicotinamide adenine dinucleotide (NAD+) to NADH, resulting in an absorbance change that is measured spectrophotometrically. Endogenous serum G6PDH does not interfere because the coenzyme functions only with the bacterial (*Leuconostoc mesenteroides*) enzyme employed in the assay.(Package insert: Roche Acetaminophen Gen. 2 reagent, Roche Diagnostic Corp, Indianapolis, IN, 04/2016)

PDF Report

No

Day(s) Performed

Monday through Sunday

Report Available

Same day/1 day

Specimen Retention Time

1 week

Performing Laboratory Location

Mayo Clinic Laboratories - Rochester Main Campus

Fees & Codes**Fees**

- Authorized users can sign in to [Test Prices](#) for detailed fee information.
- Clients without access to Test Prices can contact [Customer Service](#) 24 hours a day, seven days a week.
- Prospective clients should contact their account representative. For assistance, contact [Customer Service](#).

Test Classification

This test has been cleared, approved, or is exempt by the US Food and Drug Administration and is used per manufacturer's instructions. Performance characteristics were verified by Mayo Clinic in a manner consistent with CLIA requirements.

CPT Code Information

80143

LOINC® Information

Test ID	Test Order Name	Order LOINC® Value
ACMA	Acetaminophen, S	3298-7
Result ID	Test Result Name	Result LOINC® Value
ACMA	Acetaminophen, S	3298-7