



Test Definition: FGF1F

FGFR1 (8p11.2) Amplification, FISH, Tissue

Overview

Useful For

Identifying patients with squamous cell carcinoma of the lung, breast, esophagus, thymus, and other locations who may benefit from treatment with directed kinase inhibitors

Reflex Tests

Test Id	Reporting Name	Available Separately	Always Performed
_PBCT	Probe, +2	No, (Bill Only)	No
_PADD	Probe, +1	No, (Bill Only)	No
_PB02	Probe, +2	No, (Bill Only)	No
_PB03	Probe, +3	No, (Bill Only)	No
_IL25	Interphases, <25	No, (Bill Only)	No
_I099	Interphases, 25-99	No, (Bill Only)	No
_I300	Interphases, >=100	No, (Bill Only)	No

Testing Algorithm

This test includes a charge for the probe application, analysis, and professional interpretation of results for one probe set (2 individual fluorescence in situ hybridization probes). No analysis charges will be incurred if an insufficient number of representative cells are available for analysis.

Appropriate ancillary probes may be performed at consultant discretion to render comprehensive assessment. Any additional probes will have the results included within the final report and will be performed at an additional charge.

Method Name

Fluorescence In Situ Hybridization (FISH)

NY State Available

Yes

Specimen

Specimen Type

Tissue

Ordering Guidance

This test does not include a pathology consultation. If a pathology consultation is requested, order PATHC / Pathology Consultation, and appropriate testing will be added at the discretion of the pathologist and performed at an additional charge.

Multiple oncology (cancer) gene panels are also available. For more information see [Hematology, Oncology, and Hereditary Test Selection Guide](#).

Shipping Instructions

Advise Express Mail or equivalent if not on courier service.

Necessary Information

1. A pathology report is required for testing to be performed. If not provided, appropriate testing and/or interpretation may be compromised or delayed. Acceptable pathology reports include working drafts, preliminary pathology, or surgical pathology reports.

2. The following information must be included in the report provided.

-Patient name

-Block number - **must be on all blocks, slides, and paperwork**

-Date of collection

-Tissue source

3. A reason for testing must be provided. If this information is not provided, an appropriate indication for testing may be entered by Mayo Clinic Laboratories.

Specimen Required

Submit only 1 of the following specimens:

Preferred:

Specimen Type: Tissue block (fresh tissue is **not acceptable**)

Collection Instructions:

1. Submit a formalin-fixed, paraffin-embedded tumor tissue block.

2. Blocks prepared with alternative fixation methods (eg, Prefer, Bouin's) will be attempted but are less favorable for successful results. Provide fixation method used.

Additional Information:

1. Paraffin-embedded specimens can be from any anatomic location (skin, soft tissue, lymph node, etc.).

2. Decalcified paraffin-embedded specimens will have testing attempted; however, the success rate is approximately 50%. **Testing may be canceled** if sufficient tumor tissue is not present.

3. **Submitted fresh tissue specimens will be canceled upon receipt.** If only fresh tissue is available, embed in paraffin prior to sending.

Acceptable:

Specimen Type: Tissue slides

Slides: 1 Hematoxylin and eosin-stained and 4 unstained

Collection Instructions: Submit 1 slide stained with hematoxylin and eosin and 4 consecutive unstained, positively charged, unbaked slides with 5 micron-thick sections of the tumor tissue.

Forms

If not ordering electronically, complete, print, and send an [Oncology Test Request](#) (T729) with the specimen.

Specimen Minimum Volume

Slides: 1 Hematoxylin and eosin stained and 2 unstained

Reject Due To

All specimens will be evaluated at Mayo Clinic Laboratories for test suitability.

Specimen Stability Information

Specimen Type	Temperature	Time	Special Container
Tissue	Ambient (preferred)		
	Refrigerated		

Clinical & Interpretive**Clinical Information**

Fibroblast growth factor receptor 1 (*FGFR1*) is a receptor tyrosine kinase. *FGFR1* overexpression or amplification in squamous cell carcinoma and other neoplastic processes and is associated with tumor growth. Studies have shown overexpression or amplification of *FGFR1* to be sensitive to targeted kinase inhibitors, and *FGFR1* inhibitors may be a promising therapeutic option. Therefore, patients with *FGFR1* amplification may be candidates for kinase inhibitor therapies.

Reference Values

An interpretive report will be provided.

Interpretation

FGFR1 will be clinically interpreted as positive, negative, or equivocal.

The *FGFR1* locus is reported as amplified (positive) when the *FGFR1*:D8Z2 ratio is greater than 2.0 or an average of 6 or more copies of the *FGFR1* probe are observed per tumor nucleus.

A tumor with an *FGFR1*:D8Z2 ratio less than or equal to 2.0 and having an average of less than 6 copies of *FGFR1* per tumor nucleus is considered negative for amplification of the *FGFR1* locus.

A negative result does not exclude the presence of a neoplastic disorder.

Cautions

This test is not approved by the U.S. Food and Drug Administration, and it is best used as an adjunct to existing clinical and pathologic information.

This fluorescence in situ hybridization (FISH) assay does not rule out other chromosome abnormalities.

Fixatives other than formalin (eg, Prefer, Bouin's) may not be successful for FISH assays. Non-formalin fixed specimens will not be rejected.

Paraffin-embedded tissues that have been decalcified may not be successful for FISH analysis. The success rate of FISH studies on decalcified tissue is approximately 50%, but FISH will be attempted if sufficient tumor is present for analysis.

FISH studies will be attempted if sufficient tumor is present for analysis. The pathologist reviewing the hematoxylin and

eosin-stained slide may find it necessary to cancel testing if insufficient tissue/tumor is available for testing.

If no FISH signals or a lack of sufficient tumor tissue are observed post-hybridization, the case will be released indicating a lack of FISH results.

Clinical Reference

1. Schultheis AM, Bos M, Schmitz K, et al. Fibroblast growth factor receptor 1 (FGFR1) amplification is a potential therapeutic target in small-cell lung cancer. *Mod Pathol.* 2014;27(2):214-212
2. Heist RS, Mino-Kenudson M, Sequist LV, et al. FGFR1 amplification in squamous cell carcinoma of the lung. *J Thorac Oncol.* 2012;7(12):1775-1780
3. Weis J, Sos ML, Seidel D, et al. Frequent and focal FGFR1 amplification associates with therapeutically tractable FGFR1 dependency in squamous cell lung cancer. *Sci Transl Med.* 2010;2(62):62ra93
4. Craddock KJ, Ludkovski O, Sykes J, et al. Prognostic value of fibroblast growth factor receptor 1 gene locus amplification in resected lung squamous cell carcinoma. *J Thorac Oncol.* 2013;8(11):1371-1377
5. Schildhaus HU, Heukamp LC, Merkelbach-Bruse S, et al. Definition of a fluorescence in-situ hybridization score identifies high- and low-level FGFR1 amplification types in squamous cell lung cancer. *Mod Pathol.* 2012;25(11):1473-1480
6. Drago JZ, Formisano L, Juric D, et al. FGFR1 amplification mediates endocrine resistance but retains TORC sensitivity in metastatic hormone receptor–positive (HR+) breast cancer. *Clinical Cancer Research.* 2019;25(21):6443-51
7. Andre F, Bachelot T, Campone M, et al. Targeting FGFR with dovitinib (TKI258): preclinical and clinical data in breast cancer. *Clin Cancer Res.* 2013;19(13):3693-702

Performance**Method Description**

This test is performed using a commercially available FGFR1 FISH probe and a chromosome 8 centromere probe (D8Z2). Paraffin-embedded tissue samples are cut at 5 microns and mounted on positively charged glass slides. The selection of tissue and the identification of target areas on the hematoxylin and eosin (H and E)-stained slide are performed by a pathologist. Using the H and E-stained slide as a reference, target areas are etched with a diamond-tipped engraving tool on the back of the unstained slide to be assayed. The probe set is hybridized to the appropriate target areas, and 2 technologists each independently analyze 30 interphase nuclei (60 total) with the results expressed as a ratio of *FGFR1*:D8Z2 signals. (Unpublished Mayo method)

PDF Report

No

Day(s) Performed

Monday through Friday

Report Available

7 to 10 days

Specimen Retention Time

Slides and H and E used for analysis are retained by the laboratory in accordance with regulatory requirements. Client

provided paraffin blocks and extra unstained slides will be returned after testing is complete.

Performing Laboratory Location

Mayo Clinic Laboratories - Rochester Main Campus

Fees & Codes

Fees

- Authorized users can sign in to [Test Prices](#) for detailed fee information.
- Clients without access to Test Prices can contact [Customer Service](#) 24 hours a day, seven days a week.
- Prospective clients should contact their account representative. For assistance, contact [Customer Service](#).

Test Classification

This test was developed and its performance characteristics determined by Mayo Clinic in a manner consistent with CLIA requirements. It has not been cleared or approved by the US Food and Drug Administration.

CPT Code Information

- 88271x2, 88291-DNA probe, each (first probe set), Interpretation and report
- 88271x2-DNA probe, each; each additional probe set (if appropriate)
- 88271x1-DNA probe, each; coverage for sets containing 3 probes (if appropriate)
- 88271x2-DNA probe, each; coverage for sets containing 4 probes (if appropriate)
- 88271x3-DNA probe, each; coverage for sets containing 5 probes (if appropriate)
- 88274 w/modifier 52-Interphase in situ hybridization, <25 cells, each probe set (if appropriate)
- 88274-Interphase in situ hybridization, 25 to 99 cells, each probe set (if appropriate)

LOINC® Information

Test ID	Test Order Name	Order LOINC® Value
FGF1F	FGFR1 (8p11.2) Amp, FISH, Ts	78915-6

Result ID	Test Result Name	Result LOINC® Value
55213	Result Summary	50397-9
55214	Interpretation	69965-2
55216	Result	78915-6
CG939	Reason for Referral	42349-1
55217	Specimen	31208-2
55218	Source	31208-2
55219	Tissue ID	80398-1
55220	Method	85069-3
55221	Additional Information	48767-8
55222	Disclaimer	62364-5
55225	Released By	18771-6