



Test Definition: VWACT

von Willebrand Factor Activity, Plasma

Overview

Useful For

Diagnosis of von Willebrand disease (VWD) and differentiation of VWD subtypes or differentiation of VWD from hemophilia A

Monitoring therapeutic efficacy of treatment with DDAVP (desmopressin) or von Willebrand factor concentrates in patients with VWD

Reflex Tests

Test Id	Reporting Name	Available Separately	Always Performed
RIST	Ristocetin Cofactor, P	No	No

Testing Algorithm

If the von Willebrand factor activity is less than 55%, then the von Willebrand factor ristocetin cofactor activity assay will be performed at an additional charge.

Special Instructions

- [Coagulation Guidelines for Specimen Handling and Processing](#)

Method Name

Latex Immunoassay

NY State Available

No

Specimen

Specimen Type

Plasma Na Cit

Ordering Guidance

This activity assay is most effective when it is combined with measurement of von Willebrand factor antigen and factor VIII coagulant activity, preferably as a panel of tests with reflexive testing and interpretive reporting. See AVWPR / von Willebrand Disease Profile, Plasma.

Additional Testing Requirements

F8A / Coagulation Factor VIII Activity Assay, Plasma and VWAG / von Willebrand Factor Antigen, Plasma are recommended in conjunction with this test (von Willebrand activity).

Specimen Required**Specimen Type:** Platelet-poor plasma**Collection Container/Tube:** Light-blue top (3.2% sodium citrate)**Submission Container/Tube:** Plastic vial (polypropylene preferred)**Specimen Volume:** 2 mL Platelet-poor plasma in 2 vials, each containing 1 mL**Collection Instructions:**

1. Specimen must be collected prior to factor replacement therapy.
2. For complete instructions, see [Coagulation Guidelines for Specimen Handling and Processing](#).
3. Centrifuge, transfer all plasma into a polypropylene vial, and centrifuge plasma again.
4. Aliquot plasma (1-2 mL per aliquot) into 2 separate polypropylene vials leaving 0.25 mL in the bottom of centrifuged vial.
5. Immediately freeze plasma (no longer than 4 hours after collection) at -20 degrees C or, ideally, < or =-40 degrees C.
6. Send specimens in the same shipping container.

Additional Information:

1. Double-centrifuged specimen is critical for accurate results as platelet contamination may cause spurious results.
2. Each coagulation assay requested should have its own vial.

Forms

If not ordering electronically, complete, print, and send a [Coagulation Test Request](#) (T753) with the specimen.

Specimen Minimum Volume

Platelet-poor plasma: 0.5 mL

Reject Due To

Gross hemolysis	Reject
Gross lipemia	Reject
Gross icterus	Reject

Specimen Stability Information

Specimen Type	Temperature	Time	Special Container
Plasma Na Cit	Frozen	14 days	

Clinical & Interpretive**Clinical Information**

von Willebrand factor (VWF) is a multimeric adhesive glycoprotein that is important for platelet-platelet and platelet-vessel hemostatic interactions. In addition, plasma VWF serves as a carrier protein for coagulation factor VIII, stabilizing its procoagulant activity. VWF circulates in the blood in 2 distinct compartments; plasma VWF mainly reflects VWF synthesis and release from vascular endothelial cells and platelet VWF (about 10% of the blood VWF) reflects VWF synthesis by bone marrow megakaryocytes with storage primarily in the alpha granules of circulating platelets. VWF antigen measurement assesses the mass of plasma VWF protein but does not reflect VWF functions or platelet VWF. The

major function of VWF (mediating platelet-platelet or platelet-vessel interaction) is most commonly assessed by measurement of plasma ristocetin cofactor activity.

Patients with congenital severe type 3 von Willebrand disease (VWD) have markedly decreased or immeasurably low VWF antigen in the plasma (and in the platelets), and plasma VWF activity is very low or nondetectable. Patients with types 2A and 2B variants of VWD (with abnormal plasma VWF function and multimeric structure) may have normal or decreased plasma VWF antigen but typically have decreased plasma VWF activity and decreased higher molecular weight VWF multimers in the plasma. Patients with type 2M or type 2N VWD have normal levels of antigen, but either decreased VWF activity not caused by absence of higher molecular weight VWF multimers (type 2M VWD) or decreased factor VIII coagulant activity (type 2N VWD). Patients with type 1 VWD (with decreased but normally functioning plasma VWF) have concordantly decreased plasma VWF antigen and activity. Patients with acquired von Willebrand syndrome may have either normal or decreased plasma VWF antigen and decreased VWF activity.

Note: VWF activity measurement is most effective when it is combined with measurement of VWF antigen and factor VIII coagulant activity, preferably as a panel of tests with reflexive testing and interpretive reporting (eg, AVWPR / von Willebrand Disease Profile, Plasma).

Reference Values

55-200%

Normal, full-term infants may have mildly increased levels which reach adult levels by 90 days postnatal. Healthy, premature infants (30-36 weeks gestation) may have increased levels that reach adult levels by 180 days.

Note: Individuals of blood group "O" may have lower plasma von Willebrand factor (VWF) activity than those of other ABO blood groups, such that apparently normal individuals of blood group "O" may have plasma VWF activity as low as 40% to 50%, whereas the lower limit of the reference range for individuals of other blood groups may be 60% to 70%.

Interpretation

von Willebrand factor (VWF) activity is reduced in parallel with VWF antigen in von Willebrand disease (VWD), except in types 2A, 2B, and 2M, and some cases of acquired von Willebrand syndrome (AVWS) in which the VWF activity is disproportionately decreased relative to the level of VWF antigen.

The VWF activity may be decreased in congenital VWD or AVWS that may be associated with a variety of disorders including monoclonal gammopathies, lymphoproliferative disorders, autoimmune disorders, hypothyroidism, severe aortic stenosis, left ventricular assist device, and arteriovenous malformation.

The VWF activity may be increased in association with pregnancy or estrogen use (including oral contraceptives), acute (acute-phase reactant) or chronic inflammation, exercise or stress, liver disease, vasculitis, and thrombotic thrombocytopenic purpura/hemolytic uremic syndrome (TTP/HUS). Such increases in VWF activity may obscure the laboratory diagnosis of mild VWD.

Cautions

Measurement of von Willebrand factor (VWF) activity alone has limited diagnostic value. The diagnosis of von Willebrand disease requires a combination of clinical and laboratory information. VWF activity assay results generally must be used together with assays of VWF antigen and factor VIII coagulant activity for optimum clinical utility and diagnostic efficiency.

Results may be affected by:

- Unfractionated heparin >4.0 U/mL; may cause an overestimation of the test result
- Hemoglobin >70 mg/dL; may cause the result to be underestimated
- Bilirubin >4.2 mg/dL; may cause the result to be underestimated
- Triglycerides >1020 mg/dL; may cause the result to be underestimated
- Rheumatoid factor >200 IU/mL; may cause an overestimation of the test result

Specimens from patients who have received preparation of mouse monoclonal antibody for diagnosis or therapy may contain human antimouse antibody (HAMA). The presence of HAMA may cause an overestimation of results in immunoassays that utilize mouse monoclonal antibodies. This assay contains a blocking agent against HAMA to minimize this interference.

Clinical Reference

1. Montgomery RR. Structure and function of von Willebrand factor. In: Colman RW, Hirsh J, Marder VJ, et al, eds. Hemostasis and Thrombosis: Basic Principles and Clinical Practice. 4th ed. Lippincott Williams and Wilkins; 2001:249-274
2. Sadler JE, Lillicrap DL. von Willebrand disease: diagnosis, classification, and treatment. In: Marder VJ, Aird WC, Bennett JS, Schulman S, White II GC, eds. Hemostasis and Thrombosis: Basic Principles and Clinical Practice. 6th ed. Lippincott Williams and Wilkins; 2013:670-683
3. Falavero EJ and Lippi G. eds. Hemostasis and Thrombosis, Methods and Protocols. Humana Press; 2017
4. Salem RO, Van Cott EM. A new automated screening assay for the diagnosis of von Willebrand Disease. Am J Clin Pathol. 2007;127(5):730-735

Performance

Method Description

This is a latex particle-enhanced immunoassay to quantify von Willebrand factor (VWF) activity in plasma. The activity of VWF is determined by measuring the increase of turbidity produced by the agglutination of the latex reagent. A specific anti-VWF monoclonal antibody adsorbed onto the latex reagent, directed against the platelet-binding site of VWF (glycoprotein Ib receptor), reacts with the VWF of patient plasma. The degree of agglutination is directly proportional to the activity of VWF in the sample and is determined by measuring the decrease of transmitted light caused by the aggregates. (Package insert: HemosIL von Willebrand Factor Activity. Instrumentation Laboratory; 09/2006)

PDF Report

No

Day(s) Performed

Monday through Friday

Report Available

1 to 3 days

Specimen Retention Time

14 days

Performing Laboratory Location

Mayo Clinic Jacksonville Clinical Lab

Fees & Codes**Fees**

- Authorized users can sign in to [Test Prices](#) for detailed fee information.
- Clients without access to Test Prices can contact [Customer Service](#) 24 hours a day, seven days a week.
- Prospective clients should contact their account representative. For assistance, contact [Customer Service](#).

Test Classification

This test has been modified from the manufacturer's instructions. Its performance characteristics were determined by Mayo Clinic in a manner consistent with CLIA requirements. This test has not been cleared or approved by the US Food and Drug Administration.

CPT Code Information

85397

LOINC® Information

Test ID	Test Order Name	Order LOINC® Value
VWACT	von Willebrand Factor Activity, P	68324-3

Result ID	Test Result Name	Result LOINC® Value
VWACT	von Willebrand Factor Activity, P	68324-3