

## Overview

### Useful For

Determining whether *Escherichia coli* O157:H7 may be the cause of diarrhea

Reflexive testing for Shiga toxin and/or *E coli* O157:H7 nucleic acid amplification test-positive feces

This test is generally **not useful for** patients hospitalized more than 3 days because the yield from specimens from these patients is very low, as is the likelihood of identifying a pathogen that has not been detected previously.

### Reflex Tests

Test Id	Reporting Name	Available Separately	Always Performed
GID	Bacteria Identification	No, (Bill Only)	No
ISAE	Aerobe Ident by Sequencing	No, (Bill Only)	No
REFID	Additional Identification Procedure	No, (Bill Only)	No
EC	Serologic Agglut Method 2 Ident	No, (Bill Only)	No
RMALD	Ident by MALDI-TOF mass spec	No, (Bill Only)	No

### Testing Algorithm

When this test is ordered, the reflex tests may be performed at an additional charge.

For more information see [Laboratory Testing for Infectious Causes of Diarrhea](#).

### Special Instructions

- [Laboratory Testing for Infectious Causes of Diarrhea](#)

### Highlights

This test provides evidence of the presence of the bacterium, *Escherichia coli* O157:H7, in feces in a viable state, and provides an isolate for submission to a health department if needed. Minnesota healthcare professionals are required to report all confirmed or suspected cases of *E coli* O157:H7 and other Shiga toxin-producing *E coli* to the Minnesota Department of Health. Mayo Clinic Laboratories clients should refer to their local health departments regarding public health submission of *E coli* O157:H7 and other Shiga toxin-producing *E coli* isolates.

### Method Name

Conventional Culture

### NY State Available

Yes

## Specimen

### Specimen Type

Fecal

### Additional Testing Requirements

In some cases, local public health requirements may impact Mayo Clinic Laboratories clients, requiring, for example, submission of isolates to public health laboratories. Clients should familiarize themselves with local requirements and are responsible for submitting isolates to appropriate public health laboratories. Clients can obtain isolates of *Escherichia coli* O157:H7 species recovered from specimens submitted to Mayo Clinic Laboratories by calling 800-533-1710 as soon as possible after reporting (to ensure viability of the bacterium).

### Shipping Instructions

Specimen must arrive within 96 hours of collection.

### Necessary Information

Specimen source is required.

### Specimen Required

**Patient Preparation:** Patient should **not** use barium or bismuth for 7 to 10 days before specimen collection.

**Supplies:** Culture and Sensitivity Stool Transport Vial (T058)

**Container/Tube:** Commercially available transport system specific for recovery of enteric pathogens from fecal specimens (15 mL of non-nutritive transport medium containing phenol red as a pH indicator, either Cary-Blair or Para-Pak C and S)

**Specimen Volume:** Representative portion of fecal specimen

### Collection Instructions:

1. Collect fresh feces and submit 1 gram or 5 mL in container with transport medium.
2. Place feces in preservative within 2 hours of collection.
3. Place vial in a sealed plastic bag.

### Specimen Minimum Volume

1 mL

### Reject Due To

Unpreserved feces ECOFIX preservative Formalin or PVA fixative	Reject
--	--------

### Specimen Stability Information

Specimen Type	Temperature	Time	Special Container
Fecal	Ambient (preferred)	4 days	
	Refrigerated	4 days	

## Clinical & Interpretive

### Clinical Information

Diarrhea may be caused by a number of agents, including bacteria, viruses, parasites, and chemicals; these agents may result in similar symptoms. A thorough patient history covering symptoms, severity and duration of illness, age, travel history, food consumption, history of recent antibiotic use, and illnesses in the family or other contacts will help the healthcare professional determine the appropriate testing to be performed.

Shiga toxin-producing *Escherichia coli* (STEC) are *E. coli* strains capable of producing Shiga toxin, which can result in diarrhea that can be bloody. The incubation period between exposure and symptom onset is 1 to 9 days.

Hemolytic-uremic syndrome (HUS) is a systemic complication of STEC infection and is characterized by kidney failure, microangiopathic hemolytic anemia, and nonimmune thrombocytopenia. HUS complicates approximately 15% of STEC infections in children younger than 10 years and 6% to 9% overall.

Treatment of STEC infection consists of supportive care. Antibiotic therapy is generally not beneficial in patients with STEC infection and has been associated with development of HUS in some studies. Thus, when STEC is clinically suspected, antibiotics should be withheld. Antiperistaltic agents also increase the risk of systemic complications and should be avoided.

### Reference Values

No growth of *Escherichia coli* O157:H7

### Interpretation

The growth of *Escherichia coli* O157:H7 identifies a potential cause of diarrhea.

### Cautions

The yield of *Escherichia coli* O157:H7 is reduced when specimens are delayed in transit to the laboratory (>2 hours from collection for unpreserved specimens).

Check local public health requirements, which may require submission of isolates to a public health laboratory.

Primary testing for Shiga toxin-producing *E. coli* using Shiga toxin polymerase chain reaction and not specifically just for *E. coli* O157:H7 is recommended because roughly half of Shiga toxin-producing *E. coli* are not O157:H7.

Susceptibilities should **not** be performed on *E. coli* O157:H7 since antibiotics are not used for treatment. Any healthcare professional contemplating a request for susceptibility testing on *E. coli* O157:H7 should consult with the Laboratory Section Director for guidance.

### Clinical Reference

1. DuPont HL. Persistent diarrhea: A clinical review. *JAMA*. 2016;315(24):2712-2723. doi:10.1001/jama.2016.7833

2. Page AV, Liles WC. Enterohemorrhagic *Escherichia coli* infections and the hemolytic-uremic syndrome. *Med Clin North Am.* 2013;97(4):681-695

3. Nelson JM, Griffin PM, Jones TF, et al. Antimicrobial and antimotility agent use in persons with shiga toxin-producing *Escherichia coli* O157 infection in FoodNet Sites. *Clin Infect Dis.* 2011;52(9):1130-1130

## Performance

### Method Description

The fecal specimen is inoculated onto sorbitol MacConkey agar. After incubation, suspect colonies are identified using one or a combination of the following techniques: matrix-assisted laser desorption/ionization time-of-flight (MALDI-TOF) mass spectrometry, conventional biochemical tests, carbon source utilization, serologic methods, or nucleic acid sequencing of the 16S ribosomal RNA gene. Isolates are reported as *Escherichia coli* O157:H7 or *Escherichia coli* O157, unable to detect H7 antigen.(Pillai DR, Griener T. Culture for *Campylobacter* and related organisms. In: Leber AL, Church DL, eds. *Clinical Microbiology Procedures Handbook*. 4th ed. ASM Press; 2016:Section 3.8.2)

### PDF Report

No

### Day(s) Performed

Monday through Friday

### Report Available

2 to 4 days

### Specimen Retention Time

7 days

### Performing Laboratory Location

Mayo Clinic Laboratories - Rochester Main Campus

## Fees & Codes

### Fees

- Authorized users can sign in to [Test Prices](#) for detailed fee information.
- Clients without access to Test Prices can contact [Customer Service](#) 24 hours a day, seven days a week.
- Prospective clients should contact their account representative. For assistance, contact [Customer Service](#).

### Test Classification

This test has been cleared, approved, or is exempt by the US Food and Drug Administration and is used per manufacturer's instructions. Performance characteristics were verified by Mayo Clinic in a manner consistent with CLIA requirements.

**CPT Code Information**

87046-*Escherichia coli* O157:H7 Culture, Stool-with isolation and preliminary examination  
87077-Bacteria Identification (if appropriate)  
87153-Aerobe Ident by Sequencing (if appropriate)  
87077-Additional Identification Procedure (if appropriate)  
87147-Serologic Agglut Method 2 Ident (if appropriate)  
87077-Ident by MALDI-TOF mass spec (if appropriate)

**LOINC® Information**

Test ID	Test Order Name	Order LOINC® Value
E157C	E. coli O157:H7 Culture, F	10851-4

Result ID	Test Result Name	Result LOINC® Value
E157C	E. coli O157:H7 Culture, F	10851-4