

### Overview

#### Useful For

Follow up for abnormal biochemical results suggestive of 2-hydroxyglutaric aciduria

Establishing a molecular diagnosis for patients with 2-hydroxyglutaric aciduria

Identifying variants within genes known to be associated with 2-hydroxyglutaric aciduria, allowing for predictive testing of at-risk family members

#### Reflex Tests

Test Id	Reporting Name	Available Separately	Always Performed
_STR1	Comp Analysis using STR (Bill only)	No, (Bill only)	No
_STR2	Add'l comp analysis w/STR (Bill Only)	No, (Bill only)	No
CULFB	Fibroblast Culture for Genetic Test	Yes	No
CULAF	Amniotic Fluid Culture/Genetic Test	Yes	No
MATCC	Maternal Cell Contamination, B	Yes	No

#### Genetics Test Information

This test utilizes next-generation sequencing to detect single nucleotide and copy number variants in 4 genes associated with 2-hydroxyglutaric aciduria: *D2HGDH*, *IDH2*, *L2HGDH*, *SLC25A1*. See [Targeted Genes and Methodology Details for 2-Hydroxyglutaric Aciduria Gene Panel](#) and Method Description for additional details.

Identification of a disease-causing variant may assist with diagnosis, prognosis, clinical management, familial screening, and genetic counseling for 2-hydroxyglutaric aciduria.

#### Testing Algorithm

Prenatal specimens:

If an amniotic fluid specimen or cultured amniocytes are received, an amniotic fluid culture will be performed at an additional charge.

If a chorionic villi specimen or cultured chorionic villi are received, a fibroblast culture will be performed at an additional charge.

For any prenatal specimen that is received, maternal cell contamination testing will be performed at an additional charge.

Skin biopsy or cultured fibroblast specimens:

For skin biopsy or cultured fibroblast specimens, fibroblast culture testing will be performed at an additional charge. If

---

viable cells are not obtained, the client will be notified.

Cord blood:

For cord blood specimens that have an accompanying maternal blood specimen, maternal cell contamination studies will be performed at an additional charge.

**Special Instructions**

- [Molecular Genetics: Biochemical Disorders Patient Information](#)
- [Informed Consent for Genetic Testing](#)
- [Blood Spot Collection Card-Spanish Instructions](#)
- [Blood Spot Collection Card-Chinese Instructions](#)
- [Informed Consent for Genetic Testing \(Spanish\)](#)
- [Blood Spot Collection Instructions](#)
- [Targeted Genes and Methodology Details for 2-Hydroxyglutaric Aciduria Gene Panel](#)

**Method Name**

Sequence Capture and Targeted Next-Generation Sequencing (NGS) followed by Polymerase Chain Reaction (PCR) and Sanger Sequencing

**NY State Available**

Yes

**Specimen****Specimen Type**

Varies

**Ordering Guidance**

The recommended first-tier test for 2-hydroxyglutaric aciduria is urine organic acids; order OAU / Organic Acids Screen, Random, Urine.

Customization of this panel and single gene analysis for any gene present on this panel is available. For more information see CGPH / Custom Gene Panel, Hereditary, Next-Generation Sequencing, Varies. To modify this panel via CGPH, use the Inborn Errors of Metabolism disease state for step 1 on the [Custom Gene Ordering Tool](#).

**Additional Testing Requirements**

All prenatal specimens must be accompanied by a maternal blood specimen; order MATCC / Maternal Cell Contamination, Molecular Analysis, Varies on the maternal specimen, as this must be a different order number than the prenatal specimen.

**Specimen Required**

**Patient Preparation:** A previous hematopoietic stem cell transplant from an allogenic donor will interfere with testing. For information about testing patients who have received a hematopoietic stem cell transplant, call 800-533-1710.

**Submit only 1 of the following specimens:**

**Specimen Type:** Whole blood

**Container/Tube:**

**Preferred:** Lavender top (EDTA) or yellow top (ACD)

**Acceptable:** Green top (sodium heparin)

**Specimen Volume:** 3 mL

**Collection Instructions:**

1. Invert several times to mix blood.
2. Send whole blood specimen in original tube. **Do not aliquot.**
3. Whole blood collected postnatal from an umbilical cord is also acceptable. See Additional Information

**Specimen Stability Information:** Ambient (preferred) 4 days/Refrigerated 4 days/Frozen 4 days

**Additional Information:**

1. Specimens are preferred to be received within 4 days of collection. Extraction will be attempted for specimens received after 4 days, and DNA yield will be evaluated to determine if testing may proceed.
2. To ensure minimum volume and concentration of DNA are met, the requested volume must be submitted. Testing may be canceled if DNA requirements are inadequate.
3. For postnatal umbilical cord whole blood specimens, maternal cell contamination studies are recommended to ensure test results reflect that of the patient tested. A maternal blood specimen is required to complete maternal cell contamination studies. Order MATCC / Maternal Cell Contamination, Molecular Analysis, Varies on both the cord blood and maternal blood specimens under separate order numbers.

**Specimen Type:** Saliva

**Patient Preparation:** Patient should not eat, drink, smoke, or chew gum 30 minutes prior to collection.

**Supplies:**

DNA Saliva Kit High Yield (T1007)

Saliva Collection Kit (T786)

**Container/Tube:**

**Preferred:** High-yield DNA saliva kit

**Acceptable:** Saliva swab

**Specimen Volume:** 1 Tube if using T1007 or 2 swabs if using T786

**Collection Instructions:** Collect and send specimen per kit instructions.

**Specimen Stability Information:** Ambient (preferred) 30 days/Refrigerated 30 days

**Additional information:** Saliva specimens are acceptable but not recommended. Due to lower quantity/quality of DNA yielded from saliva, some aspects of the test may not perform as well as DNA extracted from a whole blood sample. When applicable, specific gene regions that were unable to be interrogated will be noted in the report. Alternatively, additional specimen may be required to complete testing.

**Specimen Type:** Blood spot

**Supplies:** Card-Blood Spot Collection (Filter Paper) (T493)

**Container/Tube:**

**Preferred:** Collection card (Whatman Protein Saver 903 Paper)

**Acceptable:** PerkinElmer 226 filter paper or blood spot collection card

**Specimen Volume:** 2 to 5 Blood spots

**Collection Instructions:**

---

1. An alternative blood collection option for a patient older than 1 year is a fingerstick. For detailed instructions, see . [How to Collect a Dried Blood Spot Sample](#).

2. Let blood dry on the filter paper at ambient temperature in a horizontal position for a minimum of 3 hours.

3. Do not expose specimen to heat or direct sunlight.

4. Do not stack wet specimens.

5. Keep specimen dry.

**Send:** Ambient (preferred)/Refrigerated

**Additional Information:**

1. Blood spot specimens are acceptable but not recommended. Multiple extractions will be required to obtain sufficient yield for supplemental analysis, and there is significant risk for test failure due to insufficient DNA.

2. Due to lower concentration of DNA yielded from blood spot, some aspects of the test may not perform as well as DNA extracted from a whole blood sample. When applicable, specific gene regions that were unable to be interrogated will be noted in the report. Alternatively, additional specimen may be required to complete testing.

3. For collection instructions, see [Blood Spot Collection Instructions](#)

4. For collection instructions in Spanish, see [Blood Spot Collection Card-Spanish Instructions](#) (T777)

5. For collection instructions in Chinese, see [Blood Spot Collection Card-Chinese Instructions](#) (T800)

**Specimen Type:** Skin biopsy

**Supplies:** Fibroblast Biopsy Transport Media (T115)

**Container/Tube:** Sterile container with any standard cell culture media (eg, minimal essential media, RPMI 1640). The solution should be supplemented with 1% penicillin and streptomycin.

**Specimen Volume:** 4-mm punch

**Specimen Stability Information:** Ambient (preferred) <24 hours/Refrigerated <24 hours

**Additional Information:**

1. Specimens are preferred to be received within 24 hours of collection. Culture and extraction will be attempted for specimens received after 24 hours and will be evaluated to determine if testing may proceed.

2. A separate culture charge will be assessed under CULFB / Fibroblast Culture for Biochemical or Molecular Testing. An additional 3 to 4 weeks are required to culture fibroblasts before genetic testing can occur.

**Specimen Type:** Cultured fibroblasts

**Source:** Skin or tissue

**Container/Tube:** T-25 flask

**Specimen Volume:** 2 Flasks

**Collection Instructions:** Submit confluent cultured fibroblast cells from a biopsy.

**Specimen Stability Information:** Ambient (preferred) <24 hours/Refrigerated <24 hours

**Additional Information:**

1. Specimens are preferred to be received within 24 hours of collection. Culture and extraction will be attempted for specimens received after 24 hours and will be evaluated to determine if testing may proceed.

2. A separate culture charge will be assessed under CULFB / Fibroblast Culture for Biochemical or Molecular Testing. An additional 3 to 4 weeks are required to culture fibroblasts before genetic testing can occur.

**Specimen Type:** Extracted DNA

**Container/Tube:**

**Preferred:** Screw Cap Micro Tube, 2mL with skirted conical base

**Acceptable:** Matrix tube, 1mL

**Collection Instructions:**

1. The preferred volume is at least 100 mL at a concentration of 75 ng/mL.
2. Include concentration and volume on tube.

**Specimen Stability Information:** Frozen (preferred) 1 year/Ambient/Refrigerated

**Additional Information:** DNA must be extracted in a CLIA-certified laboratory or equivalent and must be extracted from a specimen type listed as acceptable for this test (including applicable anticoagulants). Our laboratory has experience with Chemagic, Puregene, Autopure, MagnaPure, and EZ1 extraction platforms and cannot guarantee that all extraction methods are compatible with this test. If testing fails, one repeat will be attempted, and if unsuccessful, the test will be reported as failed and a charge will be applied. If applicable, specific gene regions that were unable to be interrogated due to DNA quality will be noted in the report.

**PRENATAL SPECIMENS**

**Due to its complexity, consultation with the laboratory is required** for all prenatal testing; call 800-533-1710 to speak to a genetic counselor.

**Specimen Type:** Amniotic fluid

**Container/Tube:** Amniotic fluid container

**Specimen Volume:** 20 mL

**Specimen Stability Information:** Ambient (preferred) <24 hours/Refrigerated <24 hours

**Additional Information:** Specimen will only be tested after culture.

1. Specimens are preferred to be received within 24 hours of collection. Culture and extraction will be attempted for specimens received after 24 hours and will be evaluated to determine if testing may proceed.
2. A separate culture charge will be assessed under CULAF / Culture for Genetic Testing, Amniotic Fluid. An additional 2 to 3 weeks are required to culture amniotic fluid before genetic testing can occur.
3. **All prenatal specimens must be accompanied by a maternal blood specimen;** order MATCC / Maternal Cell Contamination, Molecular Analysis, Varies on the maternal specimen.

**Specimen Type:** Confluent cultured amniocytes

This does not include cultured chorionic villi.

**Container/Tube:** T-25 flask

**Specimen Volume:** 2 Flasks

**Collection Instructions:** Submit confluent cultured cells from another laboratory

**Specimen Stability Information:** Ambient (preferred) <24 hours/Refrigerated <24 hours

**Additional Information:**

1. Specimens are preferred to be received within 24 hours of collection. Culture and extraction will be attempted for specimens received after 24 hours and will be evaluated to determine if testing may proceed.
2. A separate culture charge will be assessed under CULFB / Fibroblast Culture for Biochemical or Molecular Testing.
3. **All prenatal specimens must be accompanied by a maternal blood specimen;** order MATCC / Maternal Cell Contamination, Molecular Analysis, Varies on the maternal specimen.

**Specimen Type:** Chorionic villi

**Container/Tube:** 15-mL tube containing 15 mL of transport media

**Specimen Volume:** 20 mg

**Specimen Stability Information:** Ambient (preferred) <24 hours/Refrigerated <24 hours

**Additional Information:** Specimen will only be tested after culture.

1. Specimens are preferred to be received within 24 hours of collection. Culture and extraction will be attempted for specimens received after 24 hours and will be evaluated to determine if testing may proceed.
2. A separate culture charge will be assessed under CULFB / Fibroblast Culture for Biochemical or Molecular Testing. An additional 3 to 4 weeks are required to culture fibroblasts before genetic testing can occur.
3. **All prenatal specimens must be accompanied by a maternal blood specimen;** order MATCC / Maternal Cell Contamination, Molecular Analysis, Varies on the maternal specimen.

**Specimen Type:** Cultured chorionic villi

**Container/Tube:** T-25 flasks

**Specimen Volume:** 2 full flasks

**Collection Instructions:** Submit confluent cultured cells from another laboratory

**Specimen Stability Information:** Ambient (preferred) <24 hours/Refrigerated <24 hours

**Additional Information:**

1. Specimens are preferred to be received within 24 hours of collection. Culture and extraction will be attempted for specimens received after 24 hours and will be evaluated to determine if testing may proceed.
2. A separate culture charge will be assessed under CULFB / Fibroblast Culture for Biochemical or Molecular Testing.
3. **All prenatal specimens must be accompanied by a maternal blood specimen;** order MATCC / Maternal Cell Contamination, Molecular Analysis, Varies on the maternal specimen

## Forms

1. **New York Clients-Informed consent is required.** Document on the request form or electronic order that a copy is on file. The following documents are available:
  - [Informed Consent for Genetic Testing](#) (T576)
  - [Informed Consent for Genetic Testing \(Spanish\)](#) (T826)
2. [Molecular Genetics: Biochemical Disorders Patient Information](#) (T527)
3. If not ordering electronically, complete, print, and send a [Biochemical Genetics Test Request](#) (T798) with the specimen.

## Specimen Minimum Volume

See Specimen Required

## Reject Due To

All specimens will be evaluated at Mayo Clinic Laboratories for test suitability.

## Specimen Stability Information

Specimen Type	Temperature	Time	Special Container
Varies	Varies		

## Clinical & Interpretive

### Clinical Information

The 2-hydroxyglutaric aciduria disorders are a group of cerebral organic acidurias that present biochemically with an elevation of 2-hydroxyglutaric acid (2-HG) in the urine. There are two enantiomers or forms of 2-hydroxyglutaric acid,

---

the D-form and the L-form. Depending on the genetic defect, individuals may have an elevation of one or both forms of 2-HG. Routine organic acid analysis (OAU / Organic Acids Screen, Random, Urine), while able to detect 2-HG, is unable to distinguish between the two enantiomers; however, they can be separated with more specialized biochemical testing.(1)

L-2-hydroxyglutaric aciduria (L-2-HGA) is caused by defects in *L2HGDH* and is characterized by progressive cerebellar ataxia and intellectual disability, seizures, and macrocephaly beginning in infancy or early childhood. Symptoms worsen over time leading to severe disability by early adulthood. Magnetic resonance imaging findings include subcortical leukoencephalopathy, generalized cerebellar and cerebral atrophy, and atrophy of the corpus callosum.(1)

D-2-hydroxyglutaric aciduria (D-2-HGA) is characterized by elevated levels of D-2-hydroxyglutaric acid and typically manifests with developmental delay, seizures, and hypotonia, though can vary widely from asymptomatic to severe. There are 2 types of D-2-HGA depending on the genetic cause. D-2-HGA can either be autosomal recessive, resulting from variants in *D2HGDH* causing reduced enzymatic activity (type I) or autosomal dominant gain-of-function variants in *IDH2* causing overproduction of D-2-HG (type II).(2,3)

D,L-2-hydroxyglutaric aciduria is the most severe of the 3 and caused by defects in *SLC25A1*, which encodes the mitochondrial citrate carrier. It is characterized by neonatal-onset encephalopathy with severe muscular weakness, intractable seizures, respiratory distress, and lack of psychomotor development resulting in early death. Because of the genetic heterogeneity of the 2-hydroxyglutaric acidurias and the specialized biochemical testing needed to distinguish among the conditions, this genetic panel, which incorporates *D2HGDH*, *L2HGDH*, *IDH2*, and *SLC25A1*, is an efficient way to diagnose these conditions.(4)

### Reference Values

An interpretive report will be provided.

### Interpretation

All detected alterations are evaluated according to American College of Medical Genetics and Genomics recommendations.(5) Variants are classified based on known, predicted, or possible pathogenicity and reported with interpretive comments detailing their potential or known significance.

### Cautions

Clinical Correlation:

Test results should be interpreted in the context of clinical findings, family history, and other laboratory data.

Misinterpretation of results may occur if the information provided is inaccurate or incomplete.

If testing was performed because of a clinically significant family history, it is often useful to first test an affected family member. Detection of at least one reportable variant in an affected family member would allow for more informative testing of at-risk individuals.

To discuss the availability of additional testing options or for assistance in the interpretation of these results, contact the Mayo Clinic Laboratories genetic counselors at 800-533-1710.

Technical Limitations:

Next-generation sequencing may not detect all types of genomic variants. In rare cases, false-negative or false-positive results may occur. The depth of coverage may be variable for some target regions; assay performance below the minimum acceptable criteria or for failed regions will be noted. Given these limitations, negative results do not rule out

the diagnosis of a genetic disorder. If a specific clinical disorder is suspected, evaluation by alternative methods can be considered.

There may be regions of genes that cannot be effectively evaluated by sequencing or deletion and duplication analysis as a result of technical limitations of the assay, including regions of homology, high guanine-cytosine (GC) content, and repetitive sequences. Confirmation of select reportable variants will be performed by alternate methodologies based on internal laboratory criteria.

This test is validated to detect 95% of deletions up to 75 base pairs (bp) and insertions up to 47 bp. Deletions-insertions (delins) of 40 or more bp, including mobile element insertions, may be less reliably detected than smaller indels.

This analysis targets single and multi-exon deletions/duplications; however, in some instances single exon resolution cannot be achieved due to isolated reduction in sequence coverage or inherent genomic complexity. Balanced structural rearrangements (such as translocations and inversions) may not be detected.

Deletion/duplication events that extend past the genes included on the panel may occur. In these instances, genes included in the ordered test are provided on the report and interpreted, and genomic breakpoints are reported if they are confirmed. However, copy number variants for genes not listed in the Method Description are typically not reported or interpreted for haploinsufficiency/triplosensitivity. CMACB / Chromosomal Microarray, Congenital, Blood; WESPR / Panel to Whole Exome Sequencing Reflex Test, Varies; or WGSDX / Whole Genome Sequencing for Hereditary Disorders, Varies is recommended for a full interpretation of deletions/duplications predicted to extend past the genes included on the panel.

This test is not designed to detect low levels of mosaicism or to differentiate between somatic and germline variants. If there is a possibility that any detected variant is somatic, additional testing may be necessary to clarify the significance of results.

Genes may be added or removed based on updated clinical relevance. For the most up to date list of genes included in this test And detailed information regarding gene specific performance and technical limitations, see Method Description or contact a laboratory genetic counselor.

If the patient has had an allogeneic hematopoietic stem cell transplant or a recent non-leukocyte reduced blood transfusion, results may be inaccurate due to the presence of donor DNA. Call Mayo Clinic Laboratories for instructions for testing patients who have received a bone marrow transplant.

#### Reclassification of Variants:

Currently, it is not standard practice for the laboratory to systematically review previously classified variants on a regular basis. The laboratory encourages healthcare professionals to contact the laboratory at any time to learn how the classification of a particular variant may have changed over time.

#### Variant Evaluation:

Evaluation and categorization of variants are performed using published American College of Medical Genetics and Genomics and the Association for Molecular Pathology recommendations as a guideline. Other gene-specific guidelines may also be considered. Variants are classified based on known, predicted, or possible pathogenicity and reported with interpretive comments detailing their potential or known significance. Variants classified as benign or likely benign are

---

not reported.

Multiple in silico evaluation tools may be used to assist in the interpretation of these results. The accuracy of predictions made by in silico evaluation tools is highly dependent upon the data available for a given gene, and periodic updates to these tools may cause predictions to change over time. Results from in silico evaluation tools should be interpreted with caution and professional clinical judgment.

### Clinical Reference

1. Kranendijk M, Struys EA, Salomons GS, Van der Knaap MS, Jakobs C. Progress in understanding 2-hydroxyglutaric acidurias. *J Inher Metab Dis*. 2012;35(4):571-587. doi:10.1007/s10545-012-9462-5
2. Kranendijk M, Struys EA, van Schaftingen E, et al. IDH2 mutations in patients with D-2-hydroxyglutaric aciduria. *Science*. 2010;330(6002):336. doi:10.1126/science.1192632
3. Pop A, Struys EA, Jansen EEW, et al. D-2-hydroxyglutaric aciduria Type I: functional analysis of D2HGDH missense variants. *Hum Mutat*. 2019;40(7):975-982. doi:10.1002/humu.23751
4. Nota B, Struys EA, Pop A, et al. Deficiency in SLC25A1, encoding the mitochondrial citrate carrier, causes combined D-2- and L-2-hydroxyglutaric aciduria. *Am J Hum Genet*. 2013;92(4):627-631. doi:10.1016/j.ajhg.2013.03.009
5. Richards S, Aziz N, Bale S, et al. Standards and guidelines for the interpretation of sequence variants: a joint consensus recommendation of the American College of Medical Genetics and Genomics and the Association for Molecular Pathology. *Genet Med*. 2015;17(5):405-424

### Performance

#### Method Description

Next-generation sequencing (NGS) and/or Sanger sequencing is performed to test for the presence of variants in coding regions and intron/exon boundaries of the genes analyzed, as well as some other regions that have known disease-causing variants. The human genome reference GRCh37/hg19 build was used for sequence read alignment. At least 99% of the bases are covered at a read depth over 30X. Sensitivity is estimated at above 99% for single nucleotide variants, above 94% for deletions-insertions (delins) less than 40 base pairs (bp), above 95% for deletions up to 75 bp and insertions up to 47 bp. NGS and/or a polymerase chain reaction (PCR)-based quantitative method is performed to test for the presence of deletions and duplications in the genes analyzed.

There may be regions of genes that cannot be effectively evaluated by sequencing or deletion and duplication analysis as a result of technical limitations of the assay, including regions of homology, high guanine-cytosine (GC) content, and repetitive sequences. See [Targeted Genes and Methodology Details for 2-Hydroxyglutaric Aciduria Gene Panel](#) for details regarding the targeted genes analyzed for this test and the specific gene regions not routinely covered. (Unpublished Mayo method)

Genes analyzed: *D2HGDH*, *IDH2*, *L2HGDH*, *SLC25A1*

#### PDF Report

Supplemental

#### Day(s) Performed

Varies

## Report Available

28 to 42 days

## Specimen Retention Time

Whole blood: 28 days (if available); Saliva: 30 days (if available); Extracted DNA: 3 months; Blood Spots: 1 year (if available)

## Performing Laboratory Location

Mayo Clinic Laboratories - Rochester Main Campus

## Fees & Codes

### Fees

- Authorized users can sign in to [Test Prices](#) for detailed fee information.
- Clients without access to Test Prices can contact [Customer Service](#) 24 hours a day, seven days a week.
- Prospective clients should contact their account representative. For assistance, contact [Customer Service](#).

### Test Classification

This test was developed and its performance characteristics determined by Mayo Clinic in a manner consistent with CLIA requirements. It has not been cleared or approved by the US Food and Drug Administration.

### CPT Code Information

81479

88233-Tissue culture, skin, solid tissue biopsy (if appropriate)

88240-Cryopreservation (if appropriate)

### LOINC® Information

Test ID	Test Order Name	Order LOINC® Value
2OHGP	2-OH Glutaric Aciduria Gene Panel	105341-2

Result ID	Test Result Name	Result LOINC® Value
608764	Test Description	62364-5
608765	Specimen	31208-2
608766	Source	31208-2
608767	Result Summary	50397-9
608768	Result	82939-0
608769	Interpretation	69047-9
608770	Resources	99622-3
608771	Additional Information	48767-8
608772	Method	85069-3
608773	Genes Analyzed	48018-6
608774	Disclaimer	62364-5

---

608775	Released By	18771-6
--------	-------------	---------