

# **Test Definition: PERAM**

Perampanel, Serum

## **Overview**

### **Useful For**

Monitoring serum concentration of perampanel, in specific clinical conditions (ie, severe kidney impairment, mild to moderate hepatic impairment, and end-stage kidney disease)

Assessing compliance

Assessing potential toxicity

## **Method Name**

Liquid Chromatography Tandem Mass Spectrometry (LC-MS/MS)

#### **NY State Available**

Yes

## Specimen

## Specimen Type

Serum Red

### **Specimen Required**

Supplies: Sarstedt Aliquot Tube, 5 mL (T914)

**Collection Container/Tube:** 

Preferred: Red top (serum gel/SST is not acceptable)

Acceptable: None

Submission Container/Tube: Plastic vial

**Specimen Volume:** 1 mL **Collection Instructions:** 

- 1. Draw blood immediately before next scheduled dose.
- 2. For sustained-release formulations ONLY, draw blood a minimum of 12 hours after last dose.
- 3. Within 2 hours of collection, centrifuge and aliquot serum into a plastic vial.

#### **Forms**

If not ordering electronically, complete, print, and send 1 of the following forms with the specimen:

- -<u>Therapeutics Test Request</u> (T831)
- -Neurology Specialty Testing Client Test Request (T732)

## **Specimen Minimum Volume**

0.5 mL

# **Reject Due To**



# **Test Definition: PERAM**

Perampanel, Serum

Gross	OK
hemolysis	
Gross lipemia	OK
Gross icterus	OK

# **Specimen Stability Information**

Specimen Type	Temperature	Time	Special Container
Serum Red	Refrigerated (preferred)	28 days	
	Ambient	28 days	
	Frozen	28 days	

## Clinical & Interpretive

#### **Clinical Information**

Perampanel (Fycompa) is approved for adjunctive therapy to treat primary generalized tonic-clonic seizures in patients aged 12 years and older as well as the treatment of partial-onset seizures with or without secondarily generalized seizures in patients with epilepsy aged 4 years and older.

# **Reference Values**

180-980 ng/mL

#### Interpretation

The serum concentration should be interpreted in the context of the patient's clinical response and may provide useful information in patients showing poor response or adverse effects, particularly when perampanel is coadministered with other anticonvulsant drugs.

Most individuals display optimal response to perampanel with serum levels of 180 to 980 ng/mL. Some individuals may respond well outside of this range or may display toxicity within the therapeutic range; thus, interpretation should include clinical evaluation. Toxic levels have not been well established. Therapeutic ranges are based on specimen collected at trough (ie, immediately before the next dose).

#### **Cautions**

This test cannot be performed on whole blood.

### **Clinical Reference**

- 1. Reimers A, Berg JA, Burns ML, Brodtkorb E, Johannessen SI, Johannessen Landmark C. Reference ranges for antiepileptic drugs revisited: a practical approach to establish national guidelines. Drug Des Devel Ther. 2018;12:271-280. doi:10.2147/DDDT.S154388
- 2. Hiemke C, Bergemann N, Clement HW, et al. Consensus guidelines for therapeutic drug monitoring in neuropsychopharmacology: Update 2017. Pharmacopsychiatry. 2018;51(1-02):9-62. doi:10.1055/s-0043-116492
- 3. Milone MC, Shaw LM. Therapeutic drugs and their management. In: Rifai N, Chiu RWK, Young I, Burnham CAD, Wittwer CT, eds. Tietz Textbook of Laboratory Medicine. 7th ed. Elsevier; 2023:420-453



# **Test Definition: PERAM**

Perampanel, Serum

#### **Performance**

## **Method Description**

The serum sample is diluted in acetonitrile internal standard. The protein precipitate is centrifuged, and a portion of the supernatant is diluted with mobile phase for detection by a tandem mass spectrometer. (Unpublished Mayo method)

# **PDF Report**

No

# Day(s) Performed

Tuesday, Friday

#### Report Available

2 to 5 days

## **Specimen Retention Time**

14 days

# **Performing Laboratory Location**

Mayo Clinic Laboratories - Rochester Superior Drive

## **Fees & Codes**

#### **Fees**

- Authorized users can sign in to <u>Test Prices</u> for detailed fee information.
- Clients without access to Test Prices can contact <u>Customer Service</u> 24 hours a day, seven days a week.
- Prospective clients should contact their account representative. For assistance, contact <u>Customer Service</u>.

# **Test Classification**

This test was developed and its performance characteristics determined by Mayo Clinic in a manner consistent with CLIA requirements. It has not been cleared or approved by the US Food and Drug Administration.

#### **CPT Code Information**

80299

# **LOINC®** Information

Test ID	Test Order Name	Order LOINC® Value
PERAM	Perampanel, S	88895-8

Result ID	Test Result Name	Result LOINC® Value
609438	Perampanel, S	88895-8