

Overview

Useful For

Assessing the potential for early onset of nephropathy in diabetic patients using random urine specimens

Profile Information

Test Id	Reporting Name	Available Separately	Always Performed
RALB1	Albumin, Random, U	No	Yes
CRE2	Creatinine	No	Yes
A_CR	Albumin/Creatinine Ratio	No	Yes

Method Name

RALB1: Immunoturbidity

CRE2: Enzymatic Colorimetric Assay

NY State Available

Yes

Specimen

Specimen Type

Urine

Specimen Required

Patient Preparation: Heavy exercise should be avoided prior to collection.

Supplies: Sarstedt 5 mL Aliquot Tube (T914)

Container/Tube: Plastic tube

Specimen Volume: 5 mL

Collection Instructions:

1. Collect a random urine specimen.
2. No preservative.

Forms

If not ordering electronically, complete, print, and send 1 of the following forms with the specimen:

-[Kidney Transplant Test Request](#)

-[Renal Diagnostics Test Request](#) (T830)

Specimen Minimum Volume

1 mL

Reject Due To

All specimens will be evaluated at Mayo Clinic Laboratories for test suitability.

Specimen Stability Information

Specimen Type	Temperature	Time	Special Container
Urine	Refrigerated (preferred)	7 days	
	Ambient	7 days	
	Frozen	7 days	

Clinical & Interpretive**Clinical Information**

Diabetic nephropathy is a complication of diabetes and is characterized by proteinuria (normal urinary albumin excretion is <30 mg/day; overt proteinuria is >300 mg/day). Before overt proteinuria develops, albumin excretion increases in those diabetic patients who are destined to develop diabetic nephropathy. Therapeutic maneuvers (eg, aggressive blood pressure maintenance, particularly with angiotensin-converting enzyme inhibitors; aggressive blood sugar control; and possibly decreased protein intake) can significantly delay, or possibly prevent, development of nephropathy. Thus, there is a need to identify small, but abnormal, increases in the excretion of urinary albumin (in the range of 30-300 mg/day, ie, microalbuminuria).

The National Kidney Foundation guidelines for the management of patients with diabetes and microalbuminuria recommend that all type 1 diabetic patients older than 12 years and all type 2 diabetic patients younger than 70 years have their urine tested for microalbuminuria yearly when they are under stable glucose control.(1)

The preferred specimen is a 24-hour collection, but a random collection is acceptable. Studies have shown that correcting albumin for creatinine excretion rates has similar discriminatory value with respect to diabetic renal involvement. The albumin:creatinine ratio from a random urine specimen is also considered a valid screening tool.(2) Several studies have addressed whether the specimen needs to be a fasting urine, an exercised urine, or an overnight urine specimen. These studies have shown that the first-morning urine specimen is less sensitive, but more specific.

Studies also have shown that microalbuminuria is a marker of generalized vascular disease and is associated with stroke and heart disease.

Reference Values

Males: <17 mg/g creatinine

Females: <25 mg/g creatinine

Interpretation

In random urine specimens, normal urinary albumin excretion is below 17 mg/g creatinine for males and below 25 mg/g creatinine for females.(3)

Microalbuminuria is defined as an albumin:creatinine ratio of 17 to 299 for males and 25 to 299 for females.

A ratio of albumin:creatinine of 300 or higher is indicative of overt proteinuria.

Due to biologic variability, positive results should be confirmed by a second, first-morning random or 24-hour timed urine specimen. If there is discrepancy, a third specimen is recommended. When 2 out of 3 results are in the microalbuminuria range, this is evidence for incipient nephropathy and warrants increased efforts at glucose control, blood pressure control, and institution of therapy with an angiotensin-converting-enzyme (ACE) inhibitor (if the patient can tolerate it).

Cautions

Urine collected during menses may contain excess albumin due to blood contamination. Collection during this time should be avoided.

Heavy exercise may increase albumin excretion and should be avoided during collection. Reference values apply to a non-exercised state.

Bilirubin at 20 mg/dL reduces creatinine by 15% to 20%.

Clinical Reference

1. Bennett PH, Haffner S, Kasiske BL, et al: Screening and management of microalbuminuria in patients with diabetes mellitus: recommendations to the Scientific Advisory Board of the National Kidney Foundation from an ad hoc committee of the Council on Diabetes Mellitus of the National Kidney Foundation. *Am J Kidney Dis.* 1995 Jan;25:107-112. doi: 10.1016/0272-6386(95)90636-3
2. Krolewski AS, Laffel LM, Krolewski M, Quinn M, Warram JH: Glycosylated hemoglobin and the risk of microalbuminuria in patients with insulin-dependent diabetes mellitus. *N Engl J Med.* 1995 May 11;332:1251-1255. doi: 10.1056/NEJM199505113321902
3. Zelmanovitz T, Gross JL, Oliveira JR, Paggi A, Tatsch M, Azevedo MJ: The receiver operating characteristics curve in the evaluation of a random urine specimen as a screening test for diabetic nephropathy. *Diabetes Care.* 1997 April;20:516-519. doi: 10.2337/diacare.20.4.516
4. Miller GW, Bruns DE, Hertin GL, et al: Current issues in measurement and reporting of urinary albumin excretion. *Clin Chem.* 2009 Jan;55(1):24-38. doi: 10.1373/clinchem.2008.106567
5. Lamb EJ, Jones GRD: Kidney functions tests. In: Rifai N, Horvath AR, Wittwer CT, eds. *Tietz Textbook of Clinical Chemistry and Molecular Diagnostics.* 6th ed. Elsevier; 2018:480-488
6. Sacks DB: Diabetes mellitus. In: Rifai N, Horvath AR, Wittwer CT, eds. *Tietz Textbook of Clinical Chemistry and Molecular Diagnostics.* 6th ed. In: Elsevier; 2018:1197-1199

Performance

Method Description

Albumin is measured by immunoturbidimetry utilizing antibody to human albumin in an automated immunoprecipitation analysis system.(Package insert: ALBT2: Tina-Quant Albumin Gen 2. Roche Diagnostics V13.0 01/2020)

Creatinine is measured by the enzymatic method, which is based on the determination of sarcosine from creatinine with the aid of creatininase, creatinase, and sarcosine oxidase. The liberated hydrogen peroxide is measured via a modified

Trinder reaction using a colorimetric indicator. Optimization of the buffer system and the colorimetric indicator enables the creatinine concentration to be quantified both precisely and specifically.(Package insert: Creatinine plus v2. Roche Diagnostics; V15.0, 03/2019)

PDF Report

No

Day(s) Performed

Monday through Sunday

Report Available

1 to 3 days

Specimen Retention Time

7 days

Performing Laboratory Location

Mayo Clinic Laboratories - Rochester Main Campus

Fees & Codes**Fees**

- Authorized users can sign in to [Test Prices](#) for detailed fee information.
- Clients without access to Test Prices can contact [Customer Service](#) 24 hours a day, seven days a week.
- Prospective clients should contact their account representative. For assistance, contact [Customer Service](#).

Test Classification

This test has been modified from the manufacturer's instructions. Its performance characteristics were determined by Mayo Clinic in a manner consistent with CLIA requirements. This test has not been cleared or approved by the US Food and Drug Administration.

CPT Code Information

82043

82570

LOINC® Information

Test ID	Test Order Name	Order LOINC® Value
ALBR	Albumin, Random, U	9318-7
Result ID	Test Result Name	Result LOINC® Value
RALB1	Albumin, Random, U	89999-7
CRE2	Creatinine	2161-8
A_CR	Albumin/Creatinine Ratio	9318-7