



# Test Definition: BHDZ

Birt-Hogg-Dube Syndrome, FLCN, Full Gene Analysis, Varies

---

## Overview

### Useful For

Evaluating patients with a personal or family history suggestive of Birt-Hogg-Dube (BHD) syndrome

Establishing a diagnosis of BHD syndrome allowing for targeted cancer surveillance based on associated risks

Identifying variants within genes known to be associated with increased risk for BHD syndrome allowing for predictive testing of at-risk family members

### Genetics Test Information

This test utilizes next-generation sequencing to detect single nucleotide and copy number variants in the *FLCN* gene associated with Birt-Hogg-Dube (BHD) syndrome. See Method Description for additional details.

Identification of a disease-causing variant may assist with diagnosis, prognosis, clinical management, familial screening, and genetic counseling for BHD syndrome.

### Special Instructions

- [Molecular Genetics: Inherited Cancer Syndromes Patient Information](#)
- [Informed Consent for Genetic Testing](#)
- [Informed Consent for Genetic Testing \(Spanish\)](#)

### Method Name

Sequence Capture and Next-Generation Sequencing/Polymerase Chain Reaction (PCR)/Sanger Sequencing/Multiplex Ligation-Dependent Probe Amplification (MLPA)

### NY State Available

Yes

## Specimen

### Specimen Type

Varies

### Ordering Guidance

For a comprehensive hereditary renal cancer gene panel that includes testing for *FLCN*, consider RENC / Hereditary Renal Cancer Panel, Varies.

Testing for the *FLCN* gene as part of a customized panel is available. For more information see CGPH / Custom Gene Panel, Hereditary, Next-Generation Sequencing, Varies.

Targeted testing for familial variants (also called site-specific or known mutations testing) is available for this gene. For more information see FMTT / Familial Variant, Targeted Testing, Varies. To obtain more information about this testing option, call 800-533-1710.

**Specimen Required**

**Patient Preparation:** A previous bone marrow transplant from an allogenic donor will interfere with testing. For instructions for testing patients who have received a bone marrow transplant, call 800-533-1710.

**Submit only 1 of the following specimens:**

**Specimen Type:** Whole blood

**Container/Tube:**

**Preferred:** Lavender top (EDTA) or yellow top (ACD)

**Acceptable:** Green top (Sodium heparin)

**Specimen Volume:** 3 mL

**Collection Instructions:**

1. Invert several times to mix blood.
2. Send whole blood specimen in original tube. **Do not aliquot.**

**Specimen Stability Information:** Ambient 4 days/Refrigerated 4 days/Frozen 4 days

**Additional Information:**

1. Specimens are preferred to be received within 4 days of collection. Extraction will be attempted for samples received after 4 days and DNA yield will be evaluated to determine if testing may proceed.
2. To ensure minimum volume and concentration of DNA is met, the preferred volume of blood must be submitted. Testing may be canceled if DNA requirements are inadequate.

**Specimen Type:** Saliva

**Patient Preparation:** Patient should not eat, drink, smoke, or chew gum 30 minutes prior to collection.

**Supplies:**

DNA Saliva Kit High Yield (T1007)

Saliva Swab Collection Kit (T786)

**Container/Tube:**

**Preferred:** High-yield DNA saliva kit

**Acceptable:** Saliva swab

**Specimen Volume:** 1 Tube if using T1007 or 2 swabs if using T786

**Collection Instructions:** Collect and send specimen per kit instructions.

**Specimen Stability Information:** Ambient (preferred) 30 days/Refrigerated 30 days

**Additional Information:** Saliva specimens are acceptable but not recommended. Due to lower quantity/quality of DNA yielded from saliva, some aspects of the test may not perform as well as DNA extracted from a whole blood sample. When applicable, specific gene regions that were unable to be interrogated will be noted in the report. Alternatively, additional specimen may be required to complete testing.

**Forms**

1. **New York Clients-Informed consent is required.** Document on the request form or electronic order that a copy is on

file. The following documents are available:

[-Informed Consent for Genetic Testing \(T576\)](#)

[-Informed Consent for Genetic Testing \(Spanish\) \(T826\)](#)

2. [Molecular Genetics: Inherited Cancer Syndromes Patient Information Sheet \(T519\)](#)

3. If not ordering electronically, complete, print, and send a [Oncology Test Request \(T729\)](#) with the specimen.

**Specimen Minimum Volume**

Whole blood: 1 mL; Saliva: See Specimen Required

**Reject Due To**

All specimens will be evaluated at Mayo Clinic Laboratories for test suitability.

**Specimen Stability Information**

Specimen Type	Temperature	Time	Special Container
Varies	Varies		

**Clinical & Interpretive****Clinical Information**

Germline variants in the *FLCN* gene are associated with Birt-Hogg-Dube (BHD) syndrome. BHD syndrome is characterized by cutaneous manifestations (fibrofolliculomas, trichodiscomas/angiofibromas, perifollicular fibromas, and acrochordons), pulmonary cysts/history of pneumothorax, and various types of renal tumors. BHD syndrome is inherited in an autosomal dominant manner and the penetrance is considered to be very high.(1-6)

While there is no consensus on clinical surveillance of BHD syndrome, many recommendations have been put forth for the individual manifestations of the condition by different groups, such as the National Cancer Institute.(4-6)

**Reference Values**

An interpretive report will be provided.

**Interpretation**

All detected variants are evaluated according to American College of Medical Genetics and Genomics recommendations.(7) Variants are classified based on known, predicted, or possible pathogenicity and reported with interpretive comments detailing their potential or known significance.

**Cautions**

Clinical Correlations:

Test results should be interpreted in the context of clinical findings, family history, and other laboratory data.

Misinterpretation of results may occur if the information provided is inaccurate or incomplete.

If testing was performed because of a clinically significant family history, it is often useful to first test an affected family member. Detection of a reportable variant in an affected family member would allow for more informative testing of at-risk individuals.

To discuss the availability of additional testing options or for assistance in the interpretation of these results, contact the Mayo Clinic Laboratories genetic counselors at 800-533-1710.

**Technical Limitations:**

Next-generation sequencing may not detect all types of genomic variants. In rare cases, false-negative or false-positive results may occur. The depth of coverage may be variable for some target regions; assay performance below the minimum acceptable criteria or for failed regions will be noted. Given these limitations, negative results do not rule out the diagnosis of a genetic disorder. If a specific clinical disorder is suspected, evaluation by alternative methods can be considered.

There may be regions of genes that cannot be effectively evaluated by sequencing or deletion and duplication analysis as a result of technical limitations of the assay, including regions of homology, high guanine-cytosine (GC) content, and repetitive sequences. Confirmation of select reportable variants will be performed by alternate methodologies based on internal laboratory criteria.

This test is validated to detect 95% of deletions up to 75 base pairs (bp) and insertions up to 47 bp. Deletions-insertions (delins) of 40 or more bp, including mobile element insertions, may be less reliably detected than smaller delins.

This analysis targets single and multi-exon deletions/duplications; however, in some instances single exon resolution cannot be achieved due to isolated reduction in sequence coverage or inherent genomic complexity. Balanced structural rearrangements (such as translocations and inversions) may not be detected.

Deletion/duplication events that extend past the genes included on the panel may occur. In these instances, genes included in the ordered test are provided on the report and interpreted, and genomic breakpoints are reported if they are confirmed. However, copy number variants for genes not listed in the Method Description are typically not reported or interpreted for haploinsufficiency/triplosensitivity. CMACB / Chromosomal Microarray, Congenital, Blood; WESPR / Panel to Whole Exome Sequencing Reflex Test, Varies; or WGSDX / Whole Genome Sequencing for Hereditary Disorders, Varies is recommended for a full interpretation of deletions/duplications predicted to extend past the genes included on the panel.

Balanced structural rearrangements (such as translocations and inversions) may not be detected.

This test is not designed to detect low levels of mosaicism or differentiate between somatic and germline variants. If there is a possibility that any detected variant is somatic, additional testing may be necessary to clarify the significance of results.

For detailed information regarding gene specific performance and technical limitations, see Method Description or contact a laboratory genetic counselor.

If the patient has had an allogeneic hematopoietic stem cell transplant or a recent blood transfusion, results may be inaccurate due to the presence of donor DNA. Call Mayo Clinic Laboratories for instructions for testing patients who have received a bone marrow transplant.

**Reclassification of Variants:**

Currently, it is not standard practice for the laboratory to systematically review previously classified variants on a regular

---

basis. The laboratory encourages healthcare providers to contact the laboratory at any time to learn how the classification of a particular variant may have changed over time.

**Variant Evaluation:**

Evaluation and categorization of variants are performed using published American College of Medical Genetics and Genomics and the Association for Molecular Pathology recommendations as a guideline.<sup>(7)</sup> Other gene-specific guidelines may also be considered. Variants are classified based on known, predicted, or possible pathogenicity and reported with interpretive comments detailing their potential or known significance. Variants classified as benign or likely benign are not reported.

Multiple *in silico* evaluation tools may be used to assist in the interpretation of these results. The accuracy of predictions made by *in silico* evaluation tools is highly dependent upon the data available for a given gene, and periodic updates to these tools may cause predictions to change over time. Results from *in silico* evaluation tools should be interpreted with caution and professional clinical judgement.

**Clinical Reference**

1. Sattler EC, Steinlein OK: Birt-Hogg-Dube syndrome. In: Adam MP, Everman DB, Mirzaa GM, et al, eds. GeneReviews. [Internet]. University of Washington, Seattle; 2006. Updated January 30, 2020. Accessed April 26, 2024. Available at: [www.ncbi.nlm.nih.gov/books/NBK1522/](http://www.ncbi.nlm.nih.gov/books/NBK1522/)
2. Houweling AC, Gijzen LM, Jonker MA, et al. Renal cancer and pneumothorax risk in Birt-Hogg-Dube syndrome; an analysis of 115 FLCN mutation carriers from 35 BHD families. *Br J Cancer*. 2011;105(12):1912-1919
3. Schmidt LS, Nickerson ML, Warren MB, et al. Germline BHD-mutation spectrum and phenotype analysis of a large cohort of families with Birt-Hogg-Dube Syndrome. *Am J Hum Genet*. 2005;76(6):1023-1033
4. Stamatakis L, Metwalli AR, Middleton LA, Linehan WM. Diagnosis and management of BHD-associated kidney cancer. *Fam Cancer*. 2013;12(3):397-402
5. Farrant PBJ, Emerson R: Letter. hyfrecreation and curettage as a treatment for fibrofolliculomas in Birt-Hogg-Dube syndrome. *Dermatol Surg*. 2007;33(10):1287-1288
6. Kim D, Wysong A, Teng JM, Rahman Z. Laser-assisted delivery of topical rapamycin: mTOR inhibition for Birt-Hogg-Dube syndrome. *Dermatol Surg*. 2019;45(12):1713-1715
7. Richards S, Aziz N, Bale S, et al. Standards and guidelines for the interpretation of sequence variants: a joint consensus recommendation of the American College of Medical Genetics and Genomics and the Association for Molecular Pathology. *Genet Med*. 2015;17(5):405-424

**Performance****Method Description**

Next-generation sequencing (NGS) and/or Sanger sequencing are performed to test for the presence of variants in coding regions and intron/exon boundaries of the *FLCN* gene, as well as some other regions that have known pathogenic variants. The human genome reference GRCh37/hg19 build was used for sequence read alignment. At least 99% of the bases are covered at a read depth over 30X. Sensitivity is estimated at above 99% for single nucleotide variants, above 94% for deletions-insertions (delins) less than 40 base pairs (bp), above 95% for deletions up to 75 bp and insertions up to 47 bp. NGS and/or multiplex ligation-dependent probe amplification is performed to test for the presence of

deletions and duplications in the *FLCN* gene, including non-coding exons 1 through 3.

There may be regions of the *FLCN* gene that cannot be effectively evaluated by sequencing or deletion and duplication analysis as a result of technical limitations of the assay, including regions of homology, high guanine-cytosine (GC) content, and repetitive sequences. The reference transcript for *FLCN* gene is NM\_144997.7. Reference transcript numbers may be updated due to transcript re-versioning. Always refer to the final patient report for gene transcript information referenced at the time of testing. (Unpublished Mayo method)

Confirmation of select reportable variants may be performed by alternate methodologies based on internal laboratory criteria.

**PDF Report**

Supplemental

**Day(s) Performed**

Varies

**Report Available**

14 to 21 days

**Specimen Retention Time**

Whole blood: 2 weeks (if available); Extracted DNA: 3 months

**Performing Laboratory Location**

Mayo Clinic Laboratories - Rochester Main Campus

**Fees & Codes****Fees**

- Authorized users can sign in to [Test Prices](#) for detailed fee information.
- Clients without access to Test Prices can contact [Customer Service](#) 24 hours a day, seven days a week.
- Prospective clients should contact their account representative. For assistance, contact [Customer Service](#).

**Test Classification**

This test was developed and its performance characteristics determined by Mayo Clinic in a manner consistent with CLIA requirements. It has not been cleared or approved by the US Food and Drug Administration.

**CPT Code Information**

81479

**LOINC® Information**

Test ID	Test Order Name	Order LOINC® Value
---------	-----------------	--------------------

# Test Definition: BHDZ

Birt-Hogg-Dube Syndrome, FLCN, Full Gene Analysis, Varies

BHDZ	FLCN Full Gene Analysis	94232-6
------	-------------------------	---------

Result ID	Test Result Name	Result LOINC® Value
614635	Test Description	62364-5
614636	Specimen	31208-2
614637	Source	31208-2
614638	Result Summary	50397-9
614639	Result	82939-0
614640	Interpretation	69047-9
614641	Resources	99622-3
614642	Additional Information	48767-8
614643	Method	85069-3
614644	Genes Analyzed	48018-6
614645	Disclaimer	62364-5
614646	Released By	18771-6