

Overview

Useful For

Evaluating for chronic inflammatory demyelinating polyradiculoneuropathy (CIDP) and related demyelinating peripheral neuropathies

Profile Information

| Test Id | Reporting Name            | Available Separately | Always Performed |
|---------|---------------------------|----------------------|------------------|
| CIDPI   | CIDP/NP Interpretation, S | No                   | Yes              |
| CONCS   | Contactin-1 IgG CBA, S    | No                   | Yes              |
| NF4FS   | Neurofascin-155 IgG4, S   | No                   | Yes              |

Testing Algorithm

A thorough understanding of a patient's history along with clinical examination and laboratory testing are needed for a clinico-sero-pathological diagnosis of chronic demyelinating polyradiculoneuropathy (CIDP). To assess the probability of your patient having CIDP vs mimic disorders, see the [Chronic Inflammatory Demyelinating Polyradiculoneuropathy \(CIDP\) calculator](#).

Special Instructions

- [Acquired Neuropathy Diagnostic Algorithm](#)

Method Name

CONCS: Cell Binding Assay (CBA)  
NF4FS: Flow Cytometry (FCM)  
CIDPI: Medical Interpretation

NY State Available

Yes

Specimen

Specimen Type

Serum

Ordering Guidance

Multiple neurological phenotype-specific autoimmune/paraneoplastic evaluations are available. For more information as well as phenotype-specific testing options, refer to [Autoimmune Neurology Test Ordering Guide](#).

For a list of antibodies performed with each evaluation, see [Autoimmune Neurology Antibody Matrix](#).

Specimen Required

**Patient Preparation:** For optimal antibody detection, specimen collection is recommended before initiation of immunosuppressant medication or intravenous immunoglobulin treatment.

Collection Container/Tube:

**Preferred:** Red top

**Acceptable:** Serum gel

**Submission Container/Tube:** Plastic vial

**Specimen Volume:** 3 mL

**Collection Instructions:** Centrifuge and aliquot serum into a plastic vial.

Forms

If not ordering electronically, complete, print, and send a [Neurology Specialty Testing Client Test Request](#) (T732) with the specimen.

Specimen Minimum Volume

2 mL

Reject Due To

|                 |        |
|-----------------|--------|
| Gross hemolysis | Reject |
| Gross lipemia   | Reject |
| Gross icterus   | Reject |

Specimen Stability Information

| Specimen Type | Temperature              | Time     | Special Container |
|---------------|--------------------------|----------|-------------------|
| Serum         | Refrigerated (preferred) | 28 days  |                   |
|               | Ambient                  | 72 hours |                   |
|               | Frozen                   | 28 days  |                   |

Clinical & Interpretive

Clinical Information

Chronic inflammatory demyelinating polyradiculoneuropathy (CIDP) is an acquired, immune-mediated condition effecting peripheral nerves and nerve roots and is characterized by electrodiagnostic features of demyelination with a chronic onset that leads to significant disability. The prevalence of CIDP is estimated at approximately 2 to 4 cases per 100,000 persons. Although a rarer cause of polyneuropathy, it is important to recognize, as it is treatable with the appropriate use of immunomodulating therapies. Although the exact immunological trigger of CIDP remains unclear, a

subset of patients with suspected CIDP have been identified with autoantibodies targeting nodal-paranodal proteins. These patients share common immunopathological mechanisms of disease, clinical features, and treatment responses that are distinct from classic CIDP. A common target of these autoantibodies is the neurofascin-155 (NF155): contactin-1 (CNTN1) complex. NF155 is expressed at the paranodal loops of Schwann cells where it interacts with CNTN1 expressed on adjacent axons. This interaction stabilizes and allows the proper organization of the paranodal axoglial junction. Antibody-mediated disruption of this interaction in animal models recapitulates the pathophysiology observed in humans.

Neurofascin-155 IgG antibodies are present in approximately 5% to 10% of patients with CIDP-like presentations and, more rarely, in those with more acute forms of demyelinating polyradiculoneuropathy. NF155 IgG positive cases are more likely to present with distal weakness, gait disturbance, tremor, and dysarthria as compared to classic CIDP. Most patients who are seropositive for NF155 IgG have been reported to be refractory to intravenous immune globulin (IVIG) therapy and often require second-line treatment that includes B-cell depleting therapies such as rituximab. Studies in animal models, as well as the disease pathology, indicate NF155 IgG4 antibodies directly disrupt the paranodal axoglial junction ultimately leading to demyelination. IgG4 is the predominant antibody subclass found in these patients and is associated with poorer treatment responses to IVIG. The detection of NF155 IgG4 is a highly specific finding and has not been reported in other disease mimics such as hereditary neuropathies, distal acquired demyelinating symmetric neuropathy, and motor neuron disease.

Contactin-1 IgG antibodies are present in approximately 2% of patients with CIDP-like presentations. CNTN1 IgG positive cases are more likely to present with neuropathic pain, sensory ataxia, and subacute progressive demyelinating polyradiculoneuropathy or polyradiculopathy. The majority of seropositive patients have been reported to be refractory to treatment with IVIG. However, some of these patients respond well to B-cell depleting therapies such as rituximab. Up to half of CNTN1 IgG positive patients with CIDP or CIDP-like presentations have been reported to develop membranous nephropathy and, thus, screening for proteinuria may be warranted.

**Reference Values**

CONTACTIN-1 IgG:

Negative

NEUROFASCIN 155 IgG4:

Negative

**Interpretation**

Seropositivity for contactin-1 IgG is consistent with an immune-mediated demyelinating polyradiculoneuropathy/polyradiculopathy.

Seropositivity for neurofascin-155 IgG4 is consistent with an immune-mediated demyelinating polyradiculoneuropathy.

**Cautions**

A negative result does not exclude an immune mediated demyelinating neuropathy.

This test should only be utilized in the appropriate clinical context.

The use of immunotherapy prior to sample collection may negatively impact the sensitivity of this assay.

### Clinical Reference

1. Dubey D, Honorat JA, Shelly S, et al. Contactin-1 autoimmunity: Serologic, neurologic, and pathologic correlates. *Neurol Neuroimmunol Neuroinflamm*. 2020;7(4):e771
2. Cortese A, Lombardi R, Briani C, et al. Antibodies to neurofascin, contactin-1, and contactin-associated protein 1 in CIDP: Clinical relevance of IgG isotype. *Neurol Neuroimmunol Neuroinflamm*. 2019;7(1):e639
3. Manso C, Querol L, Mekaouche M, Illa I, Devaux JJ. Contactin-1 IgG4 antibodies cause paranode dismantling and conduction defects. *Brain*. 2016;139(Pt 6):1700-1712
4. Le Quintrec M, Teisseire M, Bec N, et al: Contactin-1 is a novel target antigen in membranous nephropathy associated with chronic inflammatory demyelinating polyneuropathy. *Kidney Int*. 2021 Dec;100(6):1240-1249
5. Ogata H, Yamasaki R, Hiwatashi A, et al: Characterization of IgG4 anti-neurofascin 155 antibody-positive polyneuropathy. *Ann Clin Transl Neurol*. 2015(10):960-971
6. Cortese A, Lombardi R, Briani C, et al. Antibodies to neurofascin, contactin-1, and contactin-associated protein 1 in CIDP: Clinical relevance of IgG isotype. *Neurol Neuroimmunol Neuroinflamm*. 2020;7(1):e639
7. Querol L, Nogales-Gadea G, Rojas-Garcia R, et al. Neurofascin IgG4 antibodies in CIDP associate with disabling tremor and poor response to IVIg. *Neurology*. 2014;82(10):879-886
8. Shelly S, Klein CJ, Dyck PJB, et al. Neurofascin-155 immunoglobulin subtypes: Clinicopathologic associations and neurologic outcomes. *Neurology*. 2021;97(24):e2392-e2403

### Performance

#### Method Description

Contactin-1 IgG:

Patient sample is applied to a composite slide containing transfected and nontransfected EU90 cells. After incubation and washing, fluorescein-conjugated goat-antihuman IgG is applied to detect the presence of patient IgG binding.(Package insert: IIFT: Neurology Mosaics, Instructions for the indirect immunofluorescence test. EUROIMMUN; FA\_112d-1\_A\_UK\_C13, 02/25/2019)

Neurofascin-155 IgG4:

This cell-binding assay utilizes flow cytometry to detect neurofascin 155 (NF155) IgG4 antibodies in patient sample. Briefly, a stable HEK293 cell line expressing human NF155 on the cell surface is premixed with parental HEK293 cells that do not express human NF155. The 2 cell populations are distinguished using a green fluorescent protein marker, which is only expressed in NF155 expressing cells. The mixture of cells is incubated with diluted patient sera to allow antibodies present in the sample to bind target antigens. Next the cells are incubated with a human IgG4 specific secondary antibody conjugated to AlexaFluor 647 to detect cell bound human IgG4 antibodies. The AlexaFluor 647 signal intensity of the different cell populations is measured using a flow cytometer. The IBI (IgG binding index) is then calculated as the median fluorescent intensity (MFI) of AlexaFluor 647 of the NF155 expressing cells divided by the MFI of the parental non-NF155 expressing cells. When the IBI is greater than or equal to 2.0 the result is considered positive for NF155 IgG4 antibodies.(Unpublished Mayo method)

PDF Report

No

Day(s) Performed

CONCS: Monday through Thursday, Sunday  
NF4FS: Monday, Friday

Report Available

5 to 10 days

Specimen Retention Time

28 days

Performing Laboratory Location

Mayo Clinic Laboratories - Rochester Main Campus

Fees & Codes

Fees

- Authorized users can sign in to [Test Prices](#) for detailed fee information.
- Clients without access to Test Prices can contact [Customer Service](#) 24 hours a day, seven days a week.
- Prospective clients should contact their account representative. For assistance, contact [Customer Service](#).

Test Classification

This test was developed and its performance characteristics determined by Mayo Clinic in a manner consistent with CLIA requirements. It has not been cleared or approved by the US Food and Drug Administration.

CPT Code Information

86255 x 2

LOINC® Information

| Test ID | Test Order Name       | Order LOINC® Value |
|---------|-----------------------|--------------------|
| CIDP    | CIDP/NP Evaluation, S | 101447-1           |

| Result ID | Test Result Name          | Result LOINC® Value |
|-----------|---------------------------|---------------------|
| 614591    | Neurofascin-155 IgG4, S   | 100845-7            |
| 616444    | CIDP/NP Interpretation, S | 69048-7             |
| 616442    | Contactin-1 IgG CBA, S    | 101448-9            |