

Overview

Useful For

Establishing a molecular diagnosis for patients with hereditary spastic paraplegia

Identifying variants within genes known to be associated with hereditary spastic paraplegia, allowing for predictive testing of at-risk family members

Genetics Test Information

This test utilizes next-generation sequencing to detect single nucleotide and copy number variants in 128 genes associated with hereditary spastic paraplegia: *ABCD1, ACO2, AFG3L2, ALDH18A1, AMPD2, AP4B1, AP4E1, AP4M1, AP4S1, AP5Z1, APOPT1 (COA8), ARG1, ARL6IP1, ASNS, ATL1, AUH, B4GALNT1, BICD2, BOLA3, BSCL2, C12orf65 (MTRFR), COQ7, CPT1C, CTC1, CTSD, CYP27A1, CYP2U1, CYP7B1, DARS1, DARS2, DDHD1, DDHD2, DLD, EARS2, ENTPD1, ERLIN1, ERLIN2, EXOSC3, FA2H, FAR1, FARS2, FUCA1, FXN, GALC, GBA2, GBE1, GJC2, GLB1, GM2A, GPT2, HACE1, HEXA, HIBCH, HSPD1, HTRA1, IBA57, IFIH1, IRF2BPL, ISCA2, KDM5C, KIDINS220, KIF1A, KIF5A, L1CAM, L2HGDH, LYRM7, MAG, MARS2, MED17, MTFMT, NIPA1, NT5C2, NUBPL, OPA3, PANK2, PDHX, PEX16, PGAP1, PLA2G6, PLP1, PNP, PNPLA6, PNPLA8, PRF1, PRUNE1, PSAP, PYCR2, RAB18, RAB3GAP1, RAB3GAP2, RARS1, REEP1, REEP2, RNASEH2A, RNASEH2B, RNASEH2C, RTN2, SACS, SAMHD1, SDHAF1, SERAC1, SLC12A6, SLC16A2, SLC19A3, SLC33A1, SLC6A8, SOX2, SPART, SPAST, SPG11, SPG21, SPG7, SPTAN1, SUMF1, TACO1, TBC1D20, TECPR2, TFG, TREX1, TTC19, TYROBP, UBAP1, UBQLN2, VAMP1, VPS13D, WASHC5, ZFYVE26, and ZFYVE27. For more information see Method Description and [Targeted Genes and Methodology Details for Inherited Spastic Paraplegia Gene Panel](#).*

Identification of a disease-causing variant may assist with diagnosis, prognosis, clinical management, recurrence risk assessment, familial screening, and genetic counseling for hereditary spastic paraplegia.

Special Instructions

- [Informed Consent for Genetic Testing](#)
- [Hereditary Peripheral Neuropathy Diagnostic Algorithm](#)
- [Molecular Genetics: Neurology Patient Information](#)
- [Informed Consent for Genetic Testing \(Spanish\)](#)
- [Targeted Genes and Methodology Details for Inherited Spastic Paraplegia Gene Panel](#)

Method Name

Sequence Capture and Targeted Next-Generation Sequencing (NGS) followed by Polymerase Chain Reaction (PCR) and Sanger Sequencing

NY State Available

Yes

Specimen

Specimen Type

Varies

Ordering Guidance

Targeted testing for familial variants (also called site-specific or known mutations testing) is available for the genes on this panel. See FMTT / Familial Variant, Targeted Testing, Varies. To obtain more information about this testing option, call 800-533-1710.

Customization of this panel and single gene analysis for any gene present on this panel are available. For more information see CGPH / Custom Gene Panel, Hereditary, Next-Generation Sequencing, Varies.

Specimen Required

Patient Preparation: A previous bone marrow transplant from an allogenic donor will interfere with testing. For instructions for testing patients who have received a bone marrow transplant, call 800-533-1710.

Submit only 1 of the following specimens:

Specimen Type: Whole blood

Container/Tube:

Preferred: Lavender top (EDTA) or yellow top (ACD)

Acceptable: None

Specimen Volume: 3 mL

Collection Instructions:

1. Invert several times to mix blood.
2. Send whole blood specimen in original tube. **Do not aliquot.**

Specimen Stability Information: Ambient (preferred) 4 days/Refrigerated 4 days/Frozen 4 days

Additional Information:

1. Specimens are preferred to be received within 4 days of collection. Extraction will be attempted for specimens received after 4 days, and DNA yield will be evaluated to determine if testing may proceed.
2. To ensure minimum volume and concentration of DNA is met, the requested volume must be submitted. Testing may be canceled if DNA requirements are inadequate.

Specimen Type: Saliva

Patient Preparation: Patient should not eat, drink, smoke, or chew gum 30 minutes prior to collection.

Supplies: Saliva Swab Collection Kit (T786)

Specimen Volume: 1 Swab

Collection Instructions: Collect and send specimen per kit instructions.

Specimen Stability Information: Ambient (preferred) 30 days/Refrigerated 30 days

Additional Information: Saliva specimens are acceptable but not recommended. Due to lower quantity/quality of DNA yielded from saliva, some aspects of the test may not perform as well as DNA extracted from a whole blood sample. When applicable, specific gene regions that were unable to be interrogated will be noted in the report. Alternatively, additional specimen may be required to complete testing.

Specimen Type: Extracted DNA

Container/Tube:

Preferred: Screw Cap Micro Tube, 2 mL with skirted conical base

Acceptable: Matrix tube, 1 mL

Collection Instructions:

- 1. The preferred volume is at least 100 mCL at a concentration of 75 ng/mCL.
- 2. Include concentration and volume on tube.

Specimen Stability Information: Frozen (preferred) 1 year/Ambient/Refrigerated

Additional Information: DNA must be extracted in a CLIA-certified laboratory or equivalent and must be extracted from a specimen type listed as acceptable for this test (including applicable anticoagulants). Our laboratory has experience with Chemagic, Puregene, Autopure, MagnaPure, and EZ1 extraction platforms and cannot guarantee that all extraction methods are compatible with this test. If testing fails, one repeat will be attempted, and if unsuccessful, the test will be reported as failed and a charge will be applied. If applicable, specific gene regions that were unable to be interrogated due to DNA quality will be noted in the report.

Forms

- 1. **New York Clients-Informed consent is required.** Document on the request form or electronic order that a copy is on file. The following documents are available:
 - [Informed Consent for Genetic Testing](#) (T576)
 - [Informed Consent for Genetic Testing \(Spanish\)](#) (T826)
- 2. [Molecular Genetics: Neurology Patient Information](#)
- 3. If not ordering electronically, complete, print, and send a [Neurology Specialty Testing Client Test Request](#) (T732) with the specimen.

Specimen Minimum Volume

See Specimen Required

Reject Due To

All specimens will be evaluated at Mayo Clinic Laboratories for test suitability.

Specimen Stability Information

Specimen Type	Temperature	Time	Special Container
Varies	Varies		

Clinical & Interpretive

Clinical Information

Hereditary spastic paraplegias (HSP) are a group of neurodegenerative disorders characterized by progressive lower extremity weakness and spasticity, both of which can be variable. Other common neurological symptoms include ataxia, cognitive impairment, neuropathy, seizures, and dysarthria. If onset of symptoms occurs in very early childhood, symptoms may be nonprogressive and resemble spastic digenic cerebral palsy. If the onset of symptoms occurs in later childhood or after, symptoms usually progress slowly and steadily.

Clinically HSP are classified in an uncomplicated or pure form and a complicated or complex form. The uncomplicated form presents with progressive lower-extremity spastic weakness, corticospinal tract signs, variable hypertonic urinary bladder disturbance, and disturbance in vibration sense and proprioception. The complicated form is characterized by

the impairments present in uncomplicated HSP plus other system involvement or other neurologic findings. Additionally, the complicated form usually follows an autosomal recessive inheritance pattern, while the uncomplicated form predominantly follows an autosomal dominant inheritance pattern.

Given HSP are a heterogeneous group of disorders, multigene panels can be an efficient and cost-effective way to establish a molecular diagnosis for symptomatic individuals.

Reference Values

An interpretive report will be provided.

Interpretation

All detected variants are evaluated according to American College of Medical Genetics and Genomics recommendations.(1) Variants are classified based on known, predicted, or possible pathogenicity and reported with interpretive comments detailing their potential or known significance.

Cautions

Clinical Correlations:

Test results should be interpreted in the context of clinical findings, family history, and other laboratory data. Misinterpretation of results may occur if the information provided is inaccurate or incomplete.

If testing was performed because of a clinically significant family history, it is often useful to first test an affected family member. Detection of a reportable variant in an affected family member would allow for more informative testing of at-risk individuals.

To discuss the availability of additional testing options or for assistance in the interpretation of these results, contact the Mayo Clinic Laboratories genetic counselors at 800-533-1710.

Technical Limitations:

Next-generation sequencing may not detect all types of genomic variants. In rare cases, false-negative or false-positive results may occur. The depth of coverage may be variable for some target regions; assay performance below the minimum acceptable criteria or for failed regions will be noted. Given these limitations, negative results do not rule out the diagnosis of a genetic disorder. If a specific clinical disorder is suspected, evaluation by alternative methods can be considered.

There may be regions of genes that cannot be effectively evaluated by sequencing or deletion and duplication analysis as a result of technical limitations of the assay, including regions of homology, high guanine-cytosine (GC) content, and repetitive sequences. Confirmation of select reportable variants will be performed by alternate methodologies based on internal laboratory criteria.

This test is validated to detect 95% of deletions up to 75 base pairs (bp) and insertions up to 47 bp. Deletions-insertions (delins) of 40 or more bp, including mobile element insertions, may be less reliably detected than smaller delins.

Deletion/Duplication Analysis:

This analysis targets single and multi-exon deletions/duplications; however, in some instances single exon resolution cannot be achieved due to isolated reduction in sequence coverage or inherent genomic complexity. Balanced structural rearrangements (such as translocations and inversions) may not be detected.

This test is not designed to detect low levels of mosaicism or to differentiate between somatic and germline variants. If there is a possibility that any detected variant is somatic, additional testing may be necessary to clarify the significance of results.

Genes may be added or removed based on updated clinical relevance. For detailed information regarding gene specific performance and technical limitations, see Method Description or contact a laboratory genetic counselor.

If the patient has had an allogeneic hematopoietic stem cell transplant or a recent blood transfusion, results may be inaccurate due to the presence of donor DNA. Call Mayo Clinic Laboratories for instructions for testing patients who have received a bone marrow transplant.

Reclassification of Variants:

Currently, it is not standard practice for the laboratory to systematically review previously classified variants on a regular basis. The laboratory encourages healthcare professional to contact the laboratory at any time to learn how the classification of a particular variant may have changed over time. Due to broadening genetic knowledge, it is possible that the laboratory may discover new information of relevance to the patient. Should that occur, the laboratory may issue an amended report.

Variant Evaluation:

Evaluation and categorization of variants are performed using published American College of Medical Genetics and Genomics and the Association for Molecular Pathology recommendations as a guideline.(1) Other gene-specific guidelines may also be considered. Variants are classified based on known, predicted, or possible pathogenicity and reported with interpretive comments detailing their potential or known significance. Variants classified as benign or likely benign are not reported.

Multiple in silico evaluation tools may be used to assist in the interpretation of these results. The accuracy of predictions made by in silico evaluation tools is highly dependent upon the data available for a given gene, and periodic updates to these tools may cause predictions to change over time. Results from in silico evaluation tools should be interpreted with caution and professional clinical judgment.

Rarely, incidental or secondary findings may implicate another predisposition or presence of active disease. These findings will be carefully reviewed to determine whether they will be reported.

Clinical Reference

1. Richards S, Aziz N, Bale S, et al. Standards and guidelines for the interpretation of sequence variants: a joint consensus recommendation of the American College of Medical Genetics and Genomics and the Association for Molecular Pathology. Genet Med. 2015;17(5):405-424
2. Murala S, Nagarajan E, Bollu PC. Hereditary spastic paraplegia. Neurol Sci. 2021;42(3):883-894

Performance

Method Description

Next-generation sequencing (NGS) and/or Sanger sequencing are performed to test for the presence of variants in coding regions and intron/exon boundaries of the genes analyzed, as well as some other regions that have known disease-causing variants. The human genome reference GRCh37/hg19 build was used for sequence read alignment. At least 99% of the bases are covered at a read depth over 30X. Sensitivity is estimated at above 99% for single nucleotide variants, above 94% for deletion-insertions (delins) less than 40 base pairs (bp), above 95% for deletions up to 75 bp and insertions up to 47 bp. NGS and/or a polymerase chain reaction-based quantitative method is performed to test for the presence of deletions and duplications in the genes analyzed.

There may be regions of genes that cannot be effectively evaluated by sequencing or deletion and duplication analysis as a result of technical limitations of the assay, including regions of homology, high guanine-cytosine (GC) content, and repetitive sequences. See [Targeted Genes and Methodology Details for Inherited Spastic Paraplegia Gene Panel](#) for details regarding the targeted genes analyzed for each test and specific gene regions not routinely covered. (Unpublished Mayo method)

Confirmation of select reportable variants may be performed by alternate methodologies based on internal laboratory criteria.

Genes analyzed: *ABCD1, ACO2, AFG3L2, ALDH18A1, AMPD2, AP4B1, AP4E1, AP4M1, AP4S1, AP5Z1, APOPT1 (COA8), ARG1, ARL6IP1, ASNS, ATL1, AUH, B4GALNT1, BICD2, BOLA3, BSCL2, C12orf65 (MTRFR), COQ7, CPT1C, CTC1, CTSD, CYP27A1, CYP2U1, CYP7B1, DARS1, DARS2, DDHD1, DDHD2, DLD, EARS2, ENTPD1, ERLIN1, ERLIN2, EXOSC3, FA2H, FAR1, FARS2, FUCA1, FXN, GALC, GBA2, GBE1, GJC2, GLB1, GM2A, GPT2, HACE1, HEXA, HIBCH, HSPD1, HTRA1, IBA57, IFIH1, IRF2BPL, ISCA2, KDM5C, KIDINS220, KIF1A, KIF5A, L1CAM, L2HGDH, LYRM7, MAG, MARS2, MED17, MTFMT, NIPA1, NT5C2, NUBPL, OPA3, PANK2, PDHX, PEX16, PGAP1, PLA2G6, PLP1, PNP, PNPLA6, PNPLA8, PRF1, PRUNE1, PSAP, PYCR2, RAB18, RAB3GAP1, RAB3GAP2, RARS1, REEP1, REEP2, RNASEH2A, RNASEH2B, RNASEH2C, RTN2, SACS, SAMHD1, SDHAF1, SERAC1, SLC12A6, SLC16A2, SLC19A3, SLC33A1, SLC6A8, SOX2, SPART, SPAST, SPG11, SPG21, SPG7, SPTAN1, SUMF1, TACO1, TBC1D20, TECPR2, TFG, TREX1, TTC19, TYROBP, UBAP1, UBQLN2, VAMP1, VPS13D, WASHC5, ZFYVE26, and ZFYVE27*

PDF Report

Supplemental

Day(s) Performed

Varies

Report Available

21 to 35 days

Specimen Retention Time

Whole blood: 30 days (if available); Extracted DNA: 3 months

Performing Laboratory Location

Mayo Clinic Laboratories - Rochester Main Campus

Fees & Codes

Fees

- Authorized users can sign in to [Test Prices](#) for detailed fee information.
- Clients without access to Test Prices can contact [Customer Service](#) 24 hours a day, seven days a week.
- Prospective clients should contact their account representative. For assistance, contact [Customer Service](#).

Test Classification

This test was developed and its performance characteristics determined by Mayo Clinic in a manner consistent with CLIA requirements. It has not been cleared or approved by the US Food and Drug Administration.

CPT Code Information

81448

LOINC® Information

Test ID	Test Order Name	Order LOINC® Value
ISPP	Spastic Paraplegia Gene Panel	103730-8

Result ID	Test Result Name	Result LOINC® Value
617611	Test Description	62364-5
617612	Specimen	31208-2
617613	Source	31208-2
617614	Result Summary	50397-9
617615	Result	82939-0
617616	Interpretation	69047-9
618183	Additional Results	82939-0
617617	Resources	99622-3
617618	Additional Information	48767-8
617619	Method	85069-3
617620	Genes Analyzed	48018-6
617621	Disclaimer	62364-5
617622	Released By	18771-6