

CD200 (OX2) Immunostain, Technical Component Only

Overview

Useful For

Identifying the presence of CD200 (OX2) protein within chronic lymphocytic leukemia, small lymphocytic lymphoma, hairy cell leukemia, and primary mediastinal large B-cell lymphoma

Reflex Tests

Test Id	Reporting Name	Available Separately	Always Performed
IHTOI	IHC Initial, Tech Only	No	No
IHTOA	IHC Additional, Tech Only	No	No

Testing Algorithm

For the initial technical component only immunohistochemical (IHC) stain performed, the appropriate bill-only test ID will be reflexed and charged (IHTOI). For each additional technical component only IHC stain performed, an additional bill-only test ID will be reflexed and charged (IHTOA).

Method Name

Immunohistochemistry (IHC)

NY State Available

Yes

Specimen

Specimen Type TECHONLY

Ordering Guidance

This test includes only technical performance of the stain (no pathologist interpretation is performed). If diagnostic consultation by a pathologist is required, order PATHC / Pathology Consultation.

Shipping Instructions

Attach the green pathology address label and the pink Immunostain Technical Only label included in the kit to the outside of the transport container.

Specimen Required

Specimen Type: Tissue Supplies: Immunostain Technical Only Envelope (T693) Container/Tube: Immunostain Technical Only Envelope



CD200 (OX2) Immunostain, Technical Component Only

Preferred:

Formalin-fixed, paraffin-embedded tissue block

OR

2 Unstained, positively charged glass slides (25- x 75- x 1-mm) per test ordered; sections 4-microns thick **Acceptable**: None

Digital Image Access

1. Information on accessing digital images of immunohistochemical (IHC) stains and the manual requisition form can be accessed through this website: <u>https://news.mayocliniclabs.com/ihc-stains/</u>

2. Clients ordering stains using a manual requisition form will not have access to digital images.

3. Clients wishing to access digital images must place the order for IHC stains electronically. Information regarding digital imaging can be accessed through this website: <u>https://news.mayocliniclabs.com/ihc-stains/#FAQ</u>

Forms

If not ordering electronically, complete, print, and send a <u>Immunohistochemical (IHC)/In Situ Hybridization (ISH) Stains</u> <u>Request</u> (T763) with the specimen.

Reject Due To

Wet/frozen	Reject
tissue	
Cytology	
smears	
Nonformalin	
fixed tissue	
Nonparaffin	
embedded	
tissue	
Noncharged	
slides	
ProbeOn slides	
Snowcoat	
slides	

Specimen Stability Information

Specimen Type	Temperature	Time	Special Container
TECHONLY	Ambient (preferred)		
	Refrigerated		

Clinical & Interpretive

Clinical Information



CD200 (OX2) Immunostain, Technical Component Only

CD200 (OX2) is a type I immunoglobulin superfamily membrane glycoprotein that exhibits membranous and cytoplasmic staining within multiple cell types (B cells, subset of T cells, dendritic cells, endothelial cells and in the peripheral/central nervous system). CD200 (OX2) is a diagnostic marker to distinguish small lymphocytic lymphoma versus mantle cell lymphoma; hairy cell leukemia versus splenic marginal zone lymphoma and primary mediastinal large B-cell lymphoma versus large B-cell lymphoma not otherwise specified.

Interpretation

This test does not include pathologist interpretation, only technical performance of the stain. If interpretation is required, order PATHC / Pathology Consultation for a full diagnostic evaluation or second opinion of the case.

The positive and negative controls are verified as showing appropriate immunoreactivity.

Interpretation of this test should be performed in the context of the patient's clinical history and other diagnostic tests by a qualified pathologist.

Cautions

Age of a cut paraffin section can affect immunoreactivity. Stability thresholds vary widely among published literature and are antigen dependent. Best practice is for paraffin sections to be cut within 6 weeks.

The charge of glass slides can be affected by environmental factors and subsequently may alter slide staining. Sending unsuitable glass slides can result in inconsistent staining due to poor slide surface chemistry.

Best practices for storage of positively charged slides: -Minimize time slides are stored after being unpackaged -Limit exposure to high humidity and heat -Minimize exposure to plastics

Clinical Reference

1. Pillai V, Pozdnyakova O, Charest K, et al. CD200 flow cytometric assessment and semiquantitative immunohistochemical staining distinguishes hairy cell leukemia from hairy cell leukemia-variant and other B-cell lymphoproliferative disorders. Am J Clin Pathol. 2013;140:536-543

 Dorfman DM, Shahsafaei A, and Alonso MA. Utility of CD200 immunostaining in the diagnosis of primary mediastinal large B cell lymphoma: comparison with MAL, CD23, and other markers. Mod Pathol. 2012;25:1637-1643
Dorfman, DM and Shahsafaei A. CD200 (OX-2 membrane glycoprotein) expression in B cell-derived neoplasms. Am J Clin Pathol. 2010;134:726-733

4. Palumbo GA, Parrinello N, Fargione G, et. Al. CD200 expression may help in differential diagnosis between mantle cell lymphoma and B-cell chronic lymphocytic leukemia. Leuk Res. 2009;33:1212-1216

Performance

Method Description

Immunohistochemistry on sections of paraffin-embedded tissue.(Unpublished Mayo method)



CD200 (OX2) Immunostain, Technical Component Only

PDF Report

No

Day(s) Performed

Monday through Friday

Report Available 1 to 3 days

Specimen Retention Time

Until staining is complete.

Performing Laboratory Location Mayo Clinic Laboratories - Rochester Main Campus

Fees & Codes

Fees

- Authorized users can sign in to <u>Test Prices</u> for detailed fee information.
- Clients without access to Test Prices can contact <u>Customer Service</u> 24 hours a day, seven days a week.
- Prospective clients should contact their account representative. For assistance, contact <u>Customer Service</u>.

Test Classification

This test has been cleared, approved, or is exempt by the US Food and Drug Administration and is used per manufacturer's instructions. Performance characteristics were verified by Mayo Clinic in a manner consistent with CLIA requirements.

CPT Code Information

88342-TC, primary 88341-TC, if additional IHC

LOINC[®] Information

CD200 (OX2) IHC, Tech Only Order only;no result	Test ID	Test Order Name	Order LOINC [®] Value
	CD200	CD200 (OX2) IHC, Tech Only	Order only;no result

Result ID	Test Result Name	Result LOINC [®] Value
619518	CD200 (OX2) IHC, Tech Only	Bill only; no result