

MayoComplete Endometrial Carcinoma Panel, Next-Generation Sequencing, Tumor

### Overview

#### **Useful For**

Primarily for determining if patients will respond to targeted therapy

Assessment of microsatellite instability for immunotherapy decisions

Molecular-based classification of endometrial carcinoma

## **Genetics Test Information**

This test uses targeted next-generation sequencing to determine microsatellite instability (MSI) status and evaluate for somatic mutations within the ATM, ATR, BRCA1, BRCA2, CDK12, CTNNB1, EPCAM, ERBB2, FBXW7, L1CAM, MLH1, MSH2, MSH6, NBN, PALB2, PMS2, POLE, PPP2R1A, and TP53 genes. See <u>Targeted Genes and Methodology Details for MayoComplete Endometrial Panel</u> for details regarding the targeted gene regions evaluated by this test.

This test is performed to evaluate for somatic mutations within solid tumor samples. This test **does not assess** for germline alterations within the genes listed.

#### **Additional Tests**

Test Id	Reporting Name	Available Separately	Always Performed
SLIRV	Slide Review in MG	No, (Bill Only)	Yes

## **Testing Algorithm**

When this test is ordered, slide review will always be performed at an additional charge.

## **Special Instructions**

- Tissue Requirements for Solid Tumor Next-Generation Sequencing
- Targeted Genes and Methodology Details for MayoComplete Endometrial Panel

### **Highlights**

This panel, performed on formalin-fixed, paraffin-embedded tumor or cytology slides, includes a curated list of 19 genes important for the clinical management of patients with endometrial cancer. This includes molecular classification of endometrial carcinoma based on current guidelines and World Health Organization recommendations.(1)

This test evaluates mismatch repair genes and microsatellite instability (MSI) status (MSS, MSI-H), as this is often clinically actionable for determining the efficacy of immunotherapy in solid tumors.

## **Method Name**

Sequence Capture and Targeted Next-Generation Sequencing (NGS)

# **NY State Available**



MayoComplete Endometrial Carcinoma Panel, Next-Generation Sequencing, Tumor

Yes

# **Specimen**

# **Specimen Type**

Varies

## **Ordering Guidance**

Multiple oncology (cancer) gene panels are available. For more information see <u>Hematology, Oncology, and Hereditary</u> <u>Test Selection Guide</u>.

## **Necessary Information**

A pathology report (final or preliminary), at minimum containing the following information, must accompany specimen for testing to be performed:

- 1. Patient name
- 2. Block number-must be on all blocks, slides, and paperwork (can be handwritten on the paperwork)
- 3. Tissue collection date
- 4. Source of the tissue

### Specimen Required

This assay requires at least 20% tumor nuclei.

- -Preferred amount of tumor area with sufficient percent tumor nuclei: tissue 216 mm(2)
- -Minimum amount of tumor area: tissue 36 mm(2)
- -These amounts are cumulative over up to 10 unstained slides and must have adequate percent tumor nuclei.
- -Tissue fixation: 10% neutral buffered formalin, not decalcified
- -For specimen preparation guidance, see <u>Tissue Requirement for Solid Tumor Next-Generation Sequencing</u>. In this document, the sizes are given as 4 mm x 4 mm x 10 slides as preferred: approximate/equivalent to 144 mm(2) and the minimum as 3 mm x 1 mm x 10 slides: approximate/equivalent to 36 mm(2).

**Preferred:** Submit 3, if available, or 2 of the following specimens. **Acceptable:** Submit **at least one** of the following specimens.

**Specimen Type:** Tissue block

Collection Instructions: Submit a formalin-fixed, paraffin-embedded tissue block with acceptable amount of tumor

tissue.

Specimen Type: Tissue slide

Slides: 1 Hematoxylin and eosin-stained and 10 unstained

**Collection Instructions**: Submit the followings slides:

1 Slide stained with hematoxylin and eosin

AND



MayoComplete Endometrial Carcinoma Panel, Next-Generation Sequencing, Tumor

10 Unstained, nonbaked slides with 5-micron thick sections of the tumor tissue.

Note: The total amount of required tumor nuclei can be obtained by scraping up to 10 slides from the same block.

Additional Information: Unused unstained slides will not be returned.

**Specimen Type:** Cytology slide (direct smears or ThinPrep)

Slides: 1 to 3 Slides

Collection Instructions: Submit 1 to 3 slides stained and coverslipped with a preferred total of 5000 nucleated cells, or a

minimum of at least 3000 nucleated cells.

**Note:** Glass coverslips are preferred; plastic coverslips are acceptable but will result in longer turnaround times. **Additional Information**: Cytology slides will not be returned. An image of the slides will be stored per regulatory

requirements.

#### Forms

If not ordering electronically, complete, print, and send an Oncology Test Request (T729) with the specimen.

## Specimen Minimum Volume

See Specimen Required

## Reject Due To

All specimens will be evaluated at Mayo Clinic Laboratories for test suitability

## **Specimen Stability Information**

Specimen Type	Temperature	Time	Special Container
Varies	Ambient (preferred)		
	Refrigerated		

# Clinical & Interpretive

#### **Clinical Information**

This test uses formalin-fixed paraffin-embedded tissue or cytology slides to assess for somatic mutations involving the following genes, known to be associated with endometrial cancer: *ATM, ATR, BRCA1, BRCA2, CDK12, CTNNB1, EPCAM, ERBB2, FBXW7, L1CAM, MLH1, MSH2, MSH6, NBN, PALB2, PMS2, POLE, PPP2R1A,* and *TP53*. The testing also includes microsatellite instability status, a biomarker for determining efficacy of immunotherapy in solid tumors. The results of this test can be useful for molecular tumor classification, assessing prognosis and guiding treatment of individuals with endometrial tumors. The data can also be useful in determining clinical trial eligibility.

# **Reference Values**

An interpretive report will be provided.

## Interpretation

The interpretation of molecular biomarker analysis includes an overview of the results and the associated diagnostic, prognostic, and therapeutic implications.



MayoComplete Endometrial Carcinoma Panel, Next-Generation Sequencing, Tumor

#### **Cautions**

This test cannot differentiate between somatic and germline alterations. Additional testing may be necessary to clarify the significance of results if there is a potential hereditary risk.

DNA variants of uncertain significance may be identified.

A negative result does not rule out the presence of a variant that may be present below the limits of detection of this assay. In a specimen with 20% or more tumor content, the analytical sensitivity of this assay for sequence reportable alterations is 5% mutant allele frequency with a minimum coverage of 500X.

Point mutations and small deletion-insertion mutations (delins) will be detected in the ATM, ATR, BRCA1, BRCA2, CDK12, CTNNB1, EPCAM, ERBB2, FBXW7, L1CAM, MLH1, MSH2, MSH6, NBN, PALB2, PMS2, POLE, PPP2R1A, and TP53 genes only. This test may detect single exon deletions but does not detect multi-exon deletions, duplications, larger-scale genomic copy number variants, copy neutral loss of heterozygosity, or epigenetic modifications such as promoter methylation. Delins of 1,000 bp or less are detectable with at least 50 or more supporting reads.

This test cannot reliably determine if a variant identified in *PMS2* exons 11-15 originated from *PMS2* or the highly homologous pseudogene *PMS2CL*. In the instance that a reportable variant is detected in *PMS2* exons 11-15, additional testing will be recommended in the patient report.

Variant allele frequency (VAF) is the percentage of sequencing reads supporting a specific variant divided by the total sequencing reads at that position. In somatic testing, VAF should be interpreted in the context of several factors, including, but not limited to, tumor purity/heterogeneity/copy number status (ploidy, gains/losses, loss of heterozygosity) and sequencing artifact/misalignment.(2,3)

Rare alterations (ie, polymorphisms) may be present that could lead to false-negative or false-positive results.

The presence or absence of a variant may not be predictive of response to therapy in all patients.

Test results should be interpreted in the context of clinical, tumor sampling, histopathological, and other laboratory data. If results obtained do not match other clinical or laboratory findings, contact the laboratory for discussion. Misinterpretation of results may occur if the information provided is inaccurate or incomplete.

Reliable results are dependent on adequate specimen collection and processing. This test has been validated on cytology slides and formalin-fixed, paraffin-embedded tissues; other types of fixatives are discouraged. Improper treatment of tissues, such as decalcification, may cause polymerase chain reaction failure.

## **Supportive Data**

Performance Characteristics:

The limit of detection for calling a somatic variant (single nucleotide variants [SNV] and deletions-insertions [delins]) is 5% variant allele frequency and having at least 500x deduplicated coverage.

Verification studies demonstrated concordance between this test and the reference method for detection of SNV and delins is 98.5% (673/683) and 98.4% (122/124) of variants, respectively. Concordance for the detection of delins was



MayoComplete Endometrial Carcinoma Panel, Next-Generation Sequencing, Tumor

99.0% (100/101) in variants 1 to 10 base pairs (bp) in size, 93.3% (14/15) in variants 11 to 50 bp in size, and 100% (8/8) in variants over50 bp in size.

Microsatellite instability (MSI) evaluation is accurate at a tumor purity of at least 10% for colorectal tumors and 20% for other tumor types. During verification studies, 98% (200/204) concordance for MSI status was observed between this test and the reference method.

To ensure accuracy, this test will be performed on cases that are estimated by a pathologist to have at least 20% tumor cells.

#### Clinical Reference

- 1. WHO Classification of Tumours Editorial Board, eds. Female genital tumours. 5th ed. World Health Organization; 2020. WHO Classification of Tumours. Vol 4
- 2. Strom SP. Current practices and guidelines for clinical next-generation sequencing oncology testing. Cancer Biol Med. 2016;13(1):3-11. doi:10.28092/j.issn.2095-3941.2016.0004
- 3. Spurr L, Li M, Alomran N, et al. Systematic pan-cancer analysis of somatic allele frequency. Sci Rep. 2018;8(1):7735. Published 2018 May 16. doi:10.1038/s41598-018-25462-0
- 4. Marcus L, Lemery SJ, Keegan P, Pazdur R. FDA Approval Summary: Pembrolizumab for the treatment of microsatellite instability-high solid tumors. Clin Cancer Res. 2019;25(13):3753-3758
- 5. Leon-Castillo A, Britton H, McConechy MK, et al. Interpretation of somatic POLE mutations in endometrial carcinoma. J Pathol. 2020;250(3):323-335
- 6. Cancer Genome Atlas Research Network; Kandoth C, Schultz N, Cherniack AD, et al. Integrated genomic characterization of endometrial carcinoma. Nature. 2013;497(7447):67-73

## **Performance**

#### **Method Description**

Next-generation sequencing is performed to determine microsatellite instability status and evaluate the presence of a mutation in most coding regions of the ATM, ATR, BRCA1, BRCA2, CDK12, CTNNB1, EPCAM, ERBB2, FBXW7, L1CAM, MLH1, MSH2, MSH6, NBN, PALB2, PMS2, POLE, PPP2R1A, and TP53 genes. See Targeted Genes and Methodology Details for MayoComplete Endometrial Panel for details regarding the targeted gene regions identified by this test. (Unpublished Mayo method)

A pathology review and macro dissection to enrich for tumor cells is performed prior to slide scraping.

### PDF Report

No

## Day(s) Performed

Monday through Friday

## Report Available

12 to 20 days



MayoComplete Endometrial Carcinoma Panel, Next-Generation Sequencing, Tumor

## **Specimen Retention Time**

Tissue blocks: Unused portions of blocks will be returned; Tissue slides: Unused slides are stored for at least 5 years;

Extracted DNA: 3 months

## **Performing Laboratory Location**

Mayo Clinic Laboratories - Rochester Main Campus

## **Fees & Codes**

# **Fees**

- Authorized users can sign in to <u>Test Prices</u> for detailed fee information.
- Clients without access to Test Prices can contact <u>Customer Service</u> 24 hours a day, seven days a week.
- Prospective clients should contact their account representative. For assistance, contact <u>Customer Service</u>.

### **Test Classification**

This test was developed and its performance characteristics determined by Mayo Clinic in a manner consistent with CLIA requirements. It has not been cleared or approved by the US Food and Drug Administration.

# **CPT Code Information**

88381-Microdissection, manual 81457

#### **LOINC®** Information

Test ID	Test Order Name	Order LOINC® Value
MCECP	MayoComplete Endometrial Panel	105590-4

Result ID	Test Result Name	Result LOINC® Value
619632	Result	82939-0
619633	Interpretation	69047-9
619634	Additional Information	48767-8
619635	Specimen	31208-2
619636	Tissue ID	80398-1
619637	Method	85069-3
619638	Disclaimer	62364-5
619639	Released By	18771-6