

# **Test Definition: MYGLS**

Myoglobin, Serum

# **Overview**

### **Useful For**

Assessing muscle damage from any cause

#### **Method Name**

Electrochemiluminescent Immunoassay (ECLIA)

### **NY State Available**

Yes

# **Specimen**

# **Specimen Type**

Serum

## **Specimen Required**

**Collection Container/Tube:** 

**Preferred:** Serum gel **Acceptable:** Red top

Submission Container/Tube: Plastic vial

Specimen Volume: 1 mL

**Collection Instructions:** Centrifuge and aliquot serum into plastic vial.

## **Specimen Minimum Volume**

0.6 mL

## **Reject Due To**

Gross	Reject
hemolysis	
Gross lipemia	Reject
Gross icterus	Reject

# **Specimen Stability Information**

Specimen Type	Temperature	Time	Special Container
Serum	Refrigerated (preferred)	14 days	
	Ambient	7 days	
	Frozen	365 days	



# **Test Definition: MYGLS**

Myoglobin, Serum

## Clinical & Interpretive

#### **Clinical Information**

Myoglobin is a heme protein found in smooth and skeletal muscles. Serum myoglobin reflects a balance between intravascular release of myoglobin from muscle and renal clearance.

Previously serum myoglobin had been advocated as a sensitive marker for early acute myocardial injury (eg, acute myocardial infarction: AMI). However, more recent studies indicate that newer markers (eg, troponin) provide superior diagnostic utility in detecting early myocardial injury.

Elevation of serum myoglobin may occur as a result of muscle trauma, resuscitation, myopathies, AMI, shock, strenuous body activity, or decreased elimination during renal insufficiency. Extreme elevations occur in rhabdomyolysis. Creatine kinase is released from muscle and used more commonly for this purpose.

#### **Reference Values**

Males: 0 to 72 mcg/L Females: 0 to 58 mcg/L

## Interpretation

Elevated myoglobin levels are seen in conditions of acute muscle injury.

#### Cautions

Elevation is nonspecific for acute myocardial infarction. The test is of no value in this regard in the presence of kidney failure, rhabdomyolysis, extensive trauma, acute peripheral vascular occlusion, or after seizures.

Serum levels rise in renal insufficiency.

In very rare cases, gammopathy, in particular type IgM (Waldenstrom macroglobulinemia), may cause unreliable results.

#### **Clinical Reference**

- 1. Lamb EJ, Jones GRD. Kidney functions tests. In: Rifai N, Horvath AR, Wittwer CT, eds. Tietz Textbook of Clinical Chemistry and Molecular Diagnostics. 6th ed. Elsevier; 2018:489
- 2. Cappenllini MD, Lo SF, Swinkels DW. Hemoglobin, iron, bilirubin. In: Rifai N, Horvath AR, Wittwer CT, eds. Tietz Textbook of Clinical Chemistry and Molecular Diagnostics. 6th ed. Elsevier; 2018:747

#### **Performance**

#### **Method Description**

This myoglobin test is a sandwich-principal assay. The first incubation is 9 mcL of sample, a biotinylated monoclonal myoglobin-specific antibody, and a monoclonal myoglobin-specific antibody labeled with a ruthenium complex, which react to form a sandwich complex. In the second incubation, the complex becomes bound to the solid phase via interaction of biotin and streptavidin after addition of streptavidin-coated microparticles. The reaction mixture is aspirated into the measuring cell where the microparticles are magnetically captured onto the surface of the electrode.



# **Test Definition: MYGLS**

Myoglobin, Serum

Results are determined via a calibration curve, which is instrument specifically generated by 2-point calibration, and a master curve provided via the cobas link.(Package insert: Elecsys Myoglobin. Roche Diagnostics; 10/2022)

## **PDF Report**

No

## Day(s) Performed

Monday through Sunday

## **Report Available**

1 to 2 days

## **Specimen Retention Time**

7 days

### **Performing Laboratory Location**

Mayo Clinic Laboratories - Rochester Main Campus

#### **Fees & Codes**

#### **Fees**

- Authorized users can sign in to <u>Test Prices</u> for detailed fee information.
- Clients without access to Test Prices can contact <u>Customer Service</u> 24 hours a day, seven days a week.
- Prospective clients should contact their account representative. For assistance, contact <u>Customer Service</u>.

## **Test Classification**

This test has been cleared, approved, or is exempt by the US Food and Drug Administration and is used per manufacturer's instructions. Performance characteristics were verified by Mayo Clinic in a manner consistent with CLIA requirements.

# **CPT Code Information**

83874

#### **LOINC®** Information

Test ID	Test Order Name	Order LOINC® Value
MYGLS	Myoglobin, S	2639-3
_		

Result ID	Test Result Name	Result LOINC® Value
MYGLS	Myoglobin, S	2639-3