



# Test Definition: ADMI

ADAMTS13 Interpretation

## Overview

### Useful For

Technical interpretation of testing performed to assist in the diagnosis of congenital, immune, or acquired thrombotic thrombocytopenic purpura

### Method Name

Only orderable as part of profile. For more information see ADAMP / ADAMTS13 Activity with Reflex Inhibitor Profile, Plasma

Technical Interpretation

### NY State Available

Yes

## Specimen

### Specimen Type

Plasma Na Cit

### Specimen Required

Only orderable as part of a profile. For more information see ADAMP / ADAMTS13 Activity with Reflex Inhibitor Profile, Plasma.

### Reject Due To

All specimens will be evaluated at Mayo Clinic Laboratories for test suitability.

### Specimen Stability Information

Specimen Type	Temperature	Time	Special Container
Plasma Na Cit	Frozen		

## Clinical & Interpretive

### Clinical Information

Thrombotic thrombocytopenic purpura (TTP), a rare (estimated incidence of 3.7 cases per million) and potentially fatal thrombotic microangiopathy syndrome, is characterized by a pentad of symptoms: thrombocytopenia, microangiopathic hemolytic anemia (intravascular hemolysis and presence of peripheral blood schistocytes), neurological symptoms, fever, and kidney dysfunction. A large majority of patients initially present with thrombocytopenia and peripheral blood

---

evidence of microangiopathy and, in the absence of any other potential explanation for such findings, satisfy criteria for early initiation of plasma exchange, which is critical for patient survival. TTP may rarely be congenital (Upshaw-Shulman syndrome) but, far more commonly, is acquired. Acquired TTP may be considered primary or idiopathic (the most frequent type) or associated with distinctive clinical conditions (secondary TTP) such as medications, hematopoietic stem cell or solid organ transplantation, sepsis, and malignancy.

The isolation and characterization of an IgG autoantibody frequently found in patients with idiopathic TTP clarified the basis of this entity and led to the isolation and characterization of a metalloprotease called ADAMTS13 (a disintegrin and metalloprotease with thrombospondin type 1 motif 13 repeats), which is the target for the IgG autoantibody, leading to a functional deficiency of ADAMTS13. ADAMTS13 cleaves the ultra-high-molecular-weight multimers of von Willebrand factor (VWF) at the peptide bond Tyr1605-Met1606 to disrupt VWF-induced platelet aggregation. The IgG antibody prevents this cleavage and leads to TTP. Although the diagnosis of TTP may be confirmed with ADAMTS13 activity and inhibition studies, the decision to initiate plasma exchange should not be delayed pending results of this assay.

ADAMTS13 activity results can have an impact on overall survival, ultimate clinical outcome, responsiveness to plasma exchange, and relapse are still controversial in recent literature. Therefore, clinical correlation is essential.

**Reference Values**

Only orderable as part of profile. For more information see ADAMP / ADAMTS13 Activity with Reflex Inhibitor Profile, Plasma.

ADAMTS13 ACTIVITY ASSAY

> or =70%

ADAMTS13 INHIBITOR TITER

<0.5 BU

**Interpretation**

Interpretive comments are provided.

Less than 10% ADAMTS13 activity is highly indicative of thrombotic thrombocytopenic purpura (TTP) in an appropriate clinical setting.

**Cautions**

No significant cautionary statements

**Clinical Reference**

1. Sadler JE. Von Willebrand factor, ADAMTS13, and thrombotic thrombocytopenic purpura. *Blood*. 2008;112(1):11-18. doi:10.1182/blood-2008-02-078170
2. George JN. How I treat patients with thrombotic thrombocytopenic purpura: 2010. *Blood*. 2010;116(20):4060-4069. doi:10.1182/blood-2010-07-271445
3. Upshaw JD: Congenital deficiency of a factor in normal plasma that reverses microangiopathic hemolysis and thrombocytopenia. *N Engl J Med*. 1978;298(24):1350-1352. doi:10.1056/NEJM197806152982407
4. Chiasakul T, Cuker A. Clinical and laboratory diagnosis of TTP: an integrated approach. *Hematology Am Soc Hematol Educ Program*. 2018;2018(1):530-538. doi:10.1182/asheducation-2018.1.530
5. Mackie I, Mancini I, Muia J, et al. International Council for Standardization in Haematology (ICSH) recommendations for laboratory measurement of ADAMTS13. *Int J Lab Hematol*. 2020;42(6):685-696. doi:10.1111/ijlh.13295

## Performance

### Method Description

Computer-generated interpretive comments will be provided based on the ADAMTS13 activity result.

### PDF Report

No

### Day(s) Performed

Monday through Sunday

### Report Available

3 to 5 days

### Specimen Retention Time

7 days

### Performing Laboratory Location

Mayo Clinic Laboratories - Rochester Main Campus

## Fees & Codes

### Fees

- Authorized users can sign in to [Test Prices](#) for detailed fee information.
- Clients without access to Test Prices can contact [Customer Service](#) 24 hours a day, seven days a week.
- Prospective clients should contact their account representative. For assistance, contact [Customer Service](#).

### Test Classification

Not Applicable

### LOINC® Information

Test ID	Test Order Name	Order LOINC® Value
ADMI	ADAMTS13 Interpretation	No LOINC Needed

Result ID	Test Result Name	Result LOINC® Value
621149	ADAMTS13 Interpretation	69049-5
621150	ADAMTS13 Interpretation	No LOINC Needed