

Nectin-4, Semi-Quantitative Immunohistochemistry, Manual

Overview

Useful For

Identifying nectin-4 overexpression in bladder cancer and other cancers

Method Name

Immunohistochemistry (IHC)

NY State Available

Yes

Specimen

Specimen Type

Special

Shipping Instructions

Attach the green "Attention Pathology" address label (T498) to the outside of the transport container before putting into the courier mailer.

Necessary Information

A pathology/diagnostic report and a brief history are required.

Specimen Required

Specimen Type: Tissue

Supplies: Pathology Packaging Kit (T554)

Collection Instructions: Formalin-fixed, paraffin-embedded tissue block; or 3 unstained glass, positively-charged slides

with 4-microns formalin-fixed, paraffin-embedded tissue

Additional Information: One slide will be stained with hematoxylin and eosin and returned.

Forms

If not ordering electronically, complete, print, and send a <u>Immunohistochemical (IHC)/In Situ Hybridization (ISH) Stains</u>
Request (T763) with the specimen.

Reject Due To

Wet/frozen	Reject
tissue	
Decalcified	
paraffin	
embedded	



Nectin-4, Semi-Quantitative Immunohistochemistry, Manual

tissue
Cytology
smears
Non-formalin
fixed tissue
Nonparaffin
embedded
tissue
Noncharged
slides
ProbeOn slides
Snowcoat
slides

Specimen Stability Information

Specimen Type	Temperature	Time	Special Container
Special	Ambient (preferred)		
	Refrigerated		

Clinical & Interpretive

Clinical Information

Nectin-4, also known as poliovirus receptor-like 4 (PVRL4), is overexpressed in many cancers, in particular bladder cancer. An antibody drug conjugate, enfortumab vedotin, targets the cell surface protein nectin-4. Some studies suggest that tumors with high expression of nectin-4 are more likely responsive to drug therapy.

Reference Values

An interpretive report will be provided.

Interpretation

This test does not include pathologist interpretation, only technical performance of the stain. If interpretation is required, order PATHC / Pathology Consultation for a full diagnostic evaluation or second opinion of the case.

The positive and negative controls are verified as showing appropriate immunoreactivity.

Interpretation of this test should be performed in the context of the patient's clinical history and other diagnostic tests by a qualified pathologist.

Cautions

This test has been validated for nondecalcified paraffin-embedded tissue specimens fixed in 10% neutral buffered formalin. Recommended fixation time is between 6 and 48 hours. This test has not been validated on tissues subjected to the decalcification process or use of alternative fixatives.



Nectin-4, Semi-Quantitative Immunohistochemistry, Manual

Age of a cut paraffin section can affect immunoreactivity. Stability thresholds vary widely among published literature and are antigen dependent. Best practice is for paraffin sections to be cut within 6 weeks.

Clinical Reference

- 1. Heath EI, Rosenberg JE. The biology and rationale of targeting nectin-4 in urothelial carcinoma. Nat Rev Urol. 2021;18(2):93-103
- 2. Bouleftour W, Guillot A, Magne N. The anti-nectin 4: A promising tumor cells target. A systematic review. Mol Cancer Ther. 2022;21(4):493-501
- 3. Takano A, Ishikawa N, Nishino R, et al. Identification of nectin-4 oncoprotein as a diagnostic and therapeutic target for lung cancer. Cancer Res. 2009;69(16):6694-6703

Performance

Method Description

Immunohistochemistry on sections of paraffin-embedded tissue.(Unpublished Mayo method)

PDF Report

No

Day(s) Performed

Monday through Friday

Report Available

5 to 7 days

Specimen Retention Time

Until reported

Performing Laboratory Location

Mayo Clinic Laboratories - Rochester Main Campus

Fees & Codes

Fees

- Authorized users can sign in to <u>Test Prices</u> for detailed fee information.
- Clients without access to Test Prices can contact <u>Customer Service</u> 24 hours a day, seven days a week.
- Prospective clients should contact their account representative. For assistance, contact <u>Customer Service</u>.

Test Classification

This test was developed and its performance characteristics determined by Mayo Clinic in a manner consistent with CLIA



Order LOINC® Value

80398-1

Nectin-4, Semi-Quantitative Immunohistochemistry, Manual

requirements. It has not been cleared or approved by the US Food and Drug Administration.

Test Order Name

Case Number

CPT Code Information

88360

Test ID

622266

LOINC® Information

NECT4	Nectin-4, SemiQuant IHC, Manual	In Process			
Result ID	Test Result Name	Result LOINC® Value			
622261	Interpretation	59465-5			
622262	Participated in the Interpretation	No LOINC Needed			
622263	Report electronically signed by	19139-5			
622264	Material Received	81178-6			
622265	Disclaimer	62364-5			