

## Overview

### Useful For

Evaluation of patients with loss of response to guselkumab, with recurrence of symptoms, or low or undetectable serum guselkumab measured at trough

### Method Name

Only orderable as part of profile. For more information see GUSAP / Guselkumab Quantitation with Antibodies, Serum.

Electrochemiluminescent-Bridging Immunoassay (ECLIA)

### NY State Available

Yes

## Specimen

### Specimen Type

Serum

### Specimen Required

Only orderable as part of profile. For more information see GUSAP / Guselkumab Quantitation with Antibodies, Serum.

**Patient Preparation:** For 12 hours before specimen collection, patient **should not** take multivitamins or dietary supplements (eg, hair, skin, and nail supplements) containing biotin (vitamin B7).

**Supplies:** Sarstedt Aliquot Tube 5 mL (T914)

#### Container/Tube:

**Preferred:** Serum gel

**Acceptable:** Red top

**Submission Container/Tube:** Plastic vial

**Specimen Volume:** 0.6 mL serum

#### Collection Instructions:

1. Draw blood immediately before next scheduled dose (trough specimen).
2. Within 2 hours of collection, centrifuge and aliquot serum into a plastic vial.

### Specimen Minimum Volume

Serum: 0.5 mL

### Reject Due To

Gross hemolysis	OK
-----------------	----

Gross lipemia	Reject
Gross icterus	OK
Heat-treated specimens	Reject

## Specimen Stability Information

Specimen Type	Temperature	Time	Special Container
Serum	Refrigerated (preferred)	28 days	
	Ambient	14 days	
	Frozen	28 days	

## Clinical & Interpretive

### Clinical Information

Guselkumab (Tremfya; Johnson and Johnson) is a fully human IgG1 lambda therapeutic monoclonal antibody used for the treatment of moderate to severe ulcerative colitis (UC) and Crohn disease (CD), as well as plaque psoriasis and psoriatic arthritis. Guselkumab targets interleukin (IL) 23A (IL-23p19) binding with high affinity to the p19 subunit and inhibiting further action.

Therapeutic drug monitoring (TDM) has become standard of care in the gastroenterology practice for biologic therapies used in CD and UC. TDM is routinely used to assess loss of response to therapy and proactively manage patients taking tumor necrosis factor inhibitors (eg, infliximab and adalimumab), alpha-4-beta7 integrins (vedolizumab), and IL-12/23 blockers (ustekinumab). With the approval of guselkumab for inflammatory bowel disease, TDM is expected to play an important role in managing loss of response to therapy and guide decision making for use of monotherapy or combination therapy.

Guselkumab, like other therapeutic monoclonal antibodies, is immunogenic. Clinical trials have shown that antibodies-to-guselkumab occur at rates of about 6% to 9% for plaque psoriasis, 2% for psoriatic arthritis, 11% for UC, and 5% for CD. The presence of anti-drug antibodies against therapeutic monoclonal antibodies has been shown to impact clinical efficacy, either by accelerated clearance or by inhibition of target binding. Assessment for the presence of antibodies to guselkumab (ATG) may be important for the management of patients, especially for those individuals with sub-therapeutic trough concentrations of guselkumab. For those individuals demonstrating loss of response in the context of sub-therapeutic drug concentrations, the presence of ATG may indicate the need to transition to another treatment approach. In contrast, those individuals with sub-therapeutic drug concentrations in the absence of detectable ATG may benefit from dose escalation.

### Reference Values

Only orderable as part of profile. For more information see GUSAP / Guselkumab Quantitation with Antibodies, Serum.

Antibodies to guselkumab  
 <9.8 ng/mL

### Interpretation

The presence of detectable anti-guselkumab antibodies may be associated with increased guselkumab clearance and lower circulating concentrations of guselkumab in serum. Low trough concentrations of guselkumab may be correlated with loss of response to the drug.

**Cautions**

Clinical management decisions for patients receiving guselkumab treatment should not be based solely on quantitation of guselkumab or assessment of antibodies to guselkumab (ATG). Test results must be interpreted within the clinical context of the patient.

Therapeutic ranges have not been established for guselkumab quantitation. Therapeutic concentrations of guselkumab may vary according to the disease (eg, Crohn disease vs psoriatic arthritis vs psoriasis).

Interference with the ATG assay, in the form of depressed signal, was observed in samples containing more than 200 ng/mL biotin.

**Clinical Reference**

1. Janssen Biotech, Inc. Highlights of prescribing information: Tremfya (guselkumab) 2017. Updated September 2025. Accessed October 2, 2025. Available at [www.janssenlabels.com/package-insert/product-monograph/prescribing-information/TREMFYA-pi.pdf](http://www.janssenlabels.com/package-insert/product-monograph/prescribing-information/TREMFYA-pi.pdf)
2. The efficacy and safety of guselkumab induction therapy in patients with moderately to severely active ulcerative colitis: Results from the Phase 3 QUASAR Induction Study. *Gastroenterol Hepatol (N Y)*. 2023;19(7 Suppl 3):9-10
3. Peyrin-Biroulet L, Allegretti JR, Rubin DT, et al. Guselkumab in patients with moderately to severely active ulcerative colitis: QUASAR Phase 2b Induction Study. *Gastroenterology*. 2023;165(6):1443-1457. doi:10.1053/j.gastro.2023.08.038
4. Danese S, Panaccione R, Feagan BG, et al. Efficacy and safety of 48 weeks of guselkumab for patients with Crohn's disease: maintenance results from the phase 2, randomized, double-blind GALAXI-1 trial. *Lancet Gastroenterol Hepatol*. 2024;9(2):133-146
5. Shao J, Vetter M, Vermeulen A, et al. Combination therapy with guselkumab and golimumab in patients with moderately to severely active ulcerative colitis: Pharmacokinetics, immunogenicity and drug-drug interactions. *Clin Pharmacol Ther*. 2024;115(6):1418-1427
6. Ladwig PM, Barnidge DR, Willrich MAV. Mass spectrometry approaches for identification and quantitation of therapeutic monoclonal antibodies in the clinical laboratory. *Clin Vaccine Immunol*. 2017;24(5):e00545-16
7. Sharma K, da Silva BC, Hanauer SB. The role of immunogenicity in optimizing biological therapies for inflammatory bowel disease. *Expert Rev Gastroenterol Hepatol*. 2025;19(3):243-258

**Performance****Method Description**

Testing for antibodies to guselkumab is accomplished using a laboratory-developed immunoassay.(Unpublished Mayo method)

**PDF Report**

No

**Day(s) Performed**

---

Thursday

**Report Available**

2 to 8 days

**Specimen Retention Time**

14 days

**Performing Laboratory Location**

Mayo Clinic Laboratories - Rochester Superior Drive

**Fees & Codes****Fees**

- Authorized users can sign in to [Test Prices](#) for detailed fee information.
- Clients without access to Test Prices can contact [Customer Service](#) 24 hours a day, seven days a week.
- Prospective clients should contact their account representative. For assistance, contact [Customer Service](#).

**Test Classification**

This test was developed and its performance characteristics determined by Mayo Clinic in a manner consistent with CLIA requirements. It has not been cleared or approved by the US Food and Drug Administration.

**CPT Code Information**

83520

**LOINC® Information**

Test ID	Test Order Name	Order LOINC® Value
GUSAB	Guselkumab Ab, S	No LOINC Needed

Result ID	Test Result Name	Result LOINC® Value
623122	Guselkumab Ab, S	In Process
623291	GUSAB Interpretation	77202-0