

Hydromorphone Confirmation, Random, Urine

## Overview

#### **Useful For**

Detection and quantification of hydromorphone in urine

#### **Method Name**

Liquid Chromatography Tandem Mass Spectrometry (LC-MS/MS)

#### **NY State Available**

Yes

# **Specimen**

# Specimen Type

Urine

# **Ordering Guidance**

For situations where chain of custody is required, a Chain of Custody Kit (T282) is available. For chain-of-custody testing, order OPATX / Opiates Confirmation, Chain of Custody, Random, Urine.

Additional drug panels and specific requests are available; call 800-533-1710.

# **Additional Testing Requirements**

**If urine creatinine is required** or adulteration of the specimen is suspected, order ADULT / Adulterants Survey, Random, Urine.

# **Specimen Required**

**Supplies:** Sarstedt Aliquot Tube, 5 mL (T914) **Collection Container/Tube:** Plastic urine container **Submission Container/Tube:** Plastic, 5-mL tube

**Specimen Volume:** 3 mL **Collection Instructions:** 

- 1. Collect a random urine specimen.
- 2. No preservative.

## **Additional Information:**

- 1. No specimen substitutions.
- 2. STAT requests are **not** accepted for this test.
- 3. Submitting less than 1 mL will compromise our ability to perform all necessary testing.

## **Forms**

If not ordering electronically, complete, print, and send a Therapeutics Test Request (T831) with the specimen.



Hydromorphone Confirmation, Random, Urine

# Specimen Minimum Volume

2.5 mL

# **Reject Due To**

Gross	OK
hemolysis	
Gross icterus	OK

# **Specimen Stability Information**

Specimen Type	Temperature	Time	Special Container
Urine	Refrigerated (preferred)	28 days	
	Ambient	14 days	
	Frozen	28 days	

# Clinical & Interpretive

#### **Clinical Information**

Opiates are the natural or synthetic drugs that have a morphine-like pharmacological action. Medically, opiates are used primarily for relief of pain. Opiates include morphine and drugs structurally similar to morphine (eg, codeine, hydrocodone, hydromorphone, oxycodone).

Hydrocodone exhibits a complex pattern of metabolism including O-demethylation, N-demethylation, and 6-keto reduction to the 6-beta hydroxymetabolites. Hydromorphone is a metabolite of hydrocodone. The presence of hydrocodone indicates exposure within 2 to 3 days prior to specimen collection.

Hydromorphone is metabolized primarily in the liver and is excreted primarily as the glucuronidated conjugate, with small amounts of parent drug and minor amounts of 6-hydroxy reduction metabolites. The presence of hydromorphone indicates exposure within 2 to 3 days prior to specimen collection. Hydromorphone is also a metabolite of hydrocodone; therefore, the presence of hydromorphone could also indicate exposure to hydrocodone.

# **Reference Values**

Negative

Positive results are reported with a quantitative result.

Cutoff concentrations by liquid chromatography tandem mass spectrometry:

Hydromorphone: 25 ng/mL

#### Interpretation

This procedure reports the total urine concentration; this is the sum of the unconjugated and conjugated forms of the parent drug.

#### **Cautions**

Other drugs in the opioid class, such as fentanyl, meperidine, methadone, and opiate antagonists such as naloxone, are



Hydromorphone Confirmation, Random, Urine

not detected.

#### **Clinical Reference**

- 1. Gutstein HB, Akil H. Opioid analgesics. In: Brunton LL, Lazo JS, Parker KL, eds. Goodman and Gilman's: The Pharmacological Basis of Therapeutics. 11th ed. McGraw-Hill; 2006:chap 21
- 2. Baselt RC, ed: Disposition of Toxic Drugs and Chemical in Man. 9th ed. Biomedical Publications; 2011
- 3. Hackett LP, Dusci LJ, Ilett KF, Chiswell GM. Optimizing the hydrolysis of codeine and morphine glucuronides in urine. Ther Drug Monit. 2002;24(5):652-657. doi:10.1097/00007691-200210000-00012
- 4. Langman LJ, Bechtel LK, Holstege CP. Clinical toxicology. In: Rifai N, Chiu RWK, Young I, Burnham CAD, Wittwer CT, eds Tietz Textbook of Laboratory Medicine. 7th ed. Elsevier; 2023:chap 43

#### **Performance**

#### **Method Description**

Confirmation with quantification by liquid chromatography/mass spectrometry (LC-MS/MS).(Unpublished Mayo method)

## **PDF Report**

No

#### Day(s) Performed

Monday through Friday

# **Report Available**

2 to 5 days

# **Specimen Retention Time**

14 days

## **Performing Laboratory Location**

Mayo Clinic Laboratories - Rochester Superior Drive

#### **Fees & Codes**

# Fees

- Authorized users can sign in to <u>Test Prices</u> for detailed fee information.
- Clients without access to Test Prices can contact <u>Customer Service</u> 24 hours a day, seven days a week.
- Prospective clients should contact their account representative. For assistance, contact <u>Customer Service</u>.

# **Test Classification**

This test was developed and its performance characteristics determined by Mayo Clinic in a manner consistent with CLIA requirements. It has not been cleared or approved by the US Food and Drug Administration.



Hydromorphone Confirmation, Random, Urine

# **CPT Code Information**

80361

G0480 (if appropriate)

# **LOINC®** Information

Test ID	Test Order Name	Order LOINC® Value
HYDMU	Hydromorphone Confirmation, U	16998-7

Result ID	Test Result Name	Result LOINC® Value
62615	Hydromorphone-by LC-MS/MS	16998-7
36025	Hydromorphone Interpretation	69050-3