Test Definition: CD20I

CD20 Immunostain, Technical Component Only

Overview

Useful For
Classification of lymphomas

Reflex Tests

Test Id	Reporting Name	Available Separately	Always Performed
IHTOI	IHC Initial, Tech Only	No	No
IHTOA	IHC Additional, Tech Only	No	No

Testing Algorithm
For the initial technical component only immunohistochemical (IHC) stain performed, the appropriate bill-only test ID will be reflexed and charged (IHTOI). For each additional technical component only IHC stain performed, an additional bill-only test ID will be reflexed and charged (IHTOA).

Method Name
Immunohistochemistry (IHC)

NY State Available
Yes

Specimen

Specimen Type
TECHONLY

Ordering Guidance
This test includes only technical performance of the stain; **no pathologist interpretation is provided**.

Technical component only stains **should not** be ordered with PATHC / Pathology Consultation. If ordered with PATHC, the technical component stains **will be canceled**. Any immunohistochemistry (IHC)/in situ hybridization (ISH) stain performed as a part of the PATHC will be performed at the reviewing pathologist's discretion at an additional charge.

Shipping Instructions
Attach the green pathology address label and the pink Immunostain Technical Only label included in the kit to the outside of the transport container.

Specimen Required
Specimen Type: Tissue

Test Definition: CD20I

CD20 Immunostain, Technical Component Only

Supplies: Immunostain Technical Only Envelope (T693)
Container/Tube: Immunostain Technical Only Envelope
Submit:
-Formalin-fixed, paraffin-embedded tissue block
OR
-2 Unstained, positively charged glass slides (25- x 75- x 1-mm) per test ordered; sections 4-microns thick
Additional Information: Broken or cracked unstained slides will be rejected. Stained slides that were broken and repaired (ie, double mounted) at the client site will not be rejected, as, in most cases, they can still be reviewed under the microscope. However, due to the double mounting, digital scanning cannot be performed.

Digital Image Access

1. Information on accessing digital images of immunohistochemical (IHC) stains and the manual requisition form can be accessed through this website: https://news.mayocliniclabs.com/ihc-stains/
2. Clients ordering stains using a manual requisition form will not have access to digital images.
3. Clients wishing to access digital images must place the order for IHC stains electronically. Information regarding digital imaging can be accessed through this website: https://news.mayocliniclabs.com/ihc-stains/#FAQ

Forms

If not ordering electronically, complete, print, and send a Immunohistochemical (IHC)/In Situ Hybridization (ISH) Stains Request (T763) with the specimen.

Reject Due To

Wet/frozen tissue	Reject
Cytology smears	Reject
Nonformalin fixed tissue	Reject
Nonparaffin embedded tissue	Reject
Noncharged slides	Reject
ProbeOn slides	Reject
Snowcoat slides	Reject
Broken or cracked unstained slides	Reject
Broken or cracked stained slides	Reject

that cannot be repaired	

Specimen Stability Information

Specimen Type	Temperature	Time	Special Container
TECHONLY	Ambient (preferred)		
	Refrigerated		

Clinical & Interpretive

Clinical Information

CD20 is a phosphorylated protein preferentially expressed by mature B lymphocytes. CD20 is not expressed by most normal plasma cells. It is one of the most specific B-cell lineage-associated antigens used in the diagnosis of B-cell lymphomas.

Interpretation

This test does not include pathologist interpretation, only technical performance of the stain. If interpretation is required, order PATHC / Pathology Consultation for a full diagnostic evaluation or second opinion of the case.

The positive and negative controls are verified as showing appropriate immunoreactivity. If a control tissue is not included on the slide, a scanned image of the relevant quality control tissue is available upon request; call 855-516-8404.

Interpretation of this test should be performed in the context of the patient's clinical history and other diagnostic tests by a qualified pathologist.

Cautions

Age of a cut paraffin section can affect immunoreactivity. Stability thresholds vary widely among published literature and are antigen dependent. Best practice is for paraffin sections to be cut within 6 weeks.

The charge of glass slides can be affected by environmental factors and subsequently may alter slide staining. Sending unsuitable glass slides can result in inconsistent staining due to poor slide surface chemistry.

Best practices for storage of positively charged slides:
-Minimize time slides are stored after being unpackaged
-Limit exposure to high humidity and heat
-Minimize exposure to plastics

Broken or cracked unstained slides are not compatible with the instrumentation used in our performing laboratories and will be rejected. Additional material may be requested.

Attempts will be made to repair broken or cracked stained slides; however, if they are unable to be repaired (ie, shattered in transit), they will be rejected and discarded due to safety concerns.

Clinical Reference

1. Choi CH, Park YH, Lim JH, et al. Prognostic implication of semi-quantitative immunohistochemical assessment of CD20 expression in diffuse large B-cell lymphoma. J Pathol Transl Med. 2016;50(2):96-103. doi:10.4132/jptm.2016.01.12
2. Duman BB, Sahin B, Ergin M, Guvenc B. Loss of CD20 antigen expression after rituximab therapy of CD20 positive B cell lymphoma (diffuse large B cell extranodal marginal zone lymphoma combination): A case report and review of the literature. Med Oncol. 2012;29(2):1223-1226. doi:10.1007/s12032-011-9955-3
3. Cang S, Mukhi N, Wang K, Liu D. Novel CD20 monoclonal antibodies for lymphoma therapy. J Hematol Oncol. 2012;5:64. doi:10.1186/1756-8722-5-64
4. Magaki S, Hojat SA, Wei B, So A, Yong WH. An introduction to the performance of immunohistochemistry. Methods Mol Biol. 2019;1897:289-298. doi:10.1007/978-1-4939-8935-5_25

Performance

Method Description

Immunohistochemistry on sections of paraffin-embedded tissue.(Unpublished Mayo method)

PDF Report

No

Day(s) Performed

Monday through Friday

Report Available

1 to 3 days

Specimen Retention Time

Until staining is complete.

Performing Laboratory Location

Mayo Clinic Laboratories - Rochester Main Campus

Fees & Codes

Fees

- Authorized users can sign in to Test Prices for detailed fee information.
- Clients without access to Test Prices can contact Customer Service 24 hours a day, seven days a week.
- Prospective clients should contact their account representative. For assistance, contact Customer Service.

Test Classification

This test was developed and its performance characteristics determined by Mayo Clinic in a manner consistent with CLIA

requirements. It has not been cleared or approved by the US Food and Drug Administration.

CPT Code Information

88342-TC, primary
88341-TC, if additional IHC

LOINC® Information

Test ID	Test Order Name	Order LOINC® Value
CD20I	CD20 IHC, Tech Only	No LOINC Needed

Result ID	Test Result Name	Result LOINC® Value
70692	CD20 IHC, Tech Only	Bill only; no result