

IgM Immunostain, Technical Component Only

### **Overview**

### **Useful For**

Classification of lymphomas and multiple myeloma

#### **Reflex Tests**

| Test Id | Reporting Name            | Available Separately | Always Performed |
|---------|---------------------------|----------------------|------------------|
| IHTOI   | IHC Initial, Tech Only    | No                   | No               |
| IHTOA   | IHC Additional, Tech Only | No                   | No               |

### **Testing Algorithm**

For the initial technical component only immunohistochemical (IHC) stain performed, the appropriate bill-only test ID will be reflexed and charged (IHTOI). For each additional technical component only IHC stain performed, an additional bill-only test ID will be reflexed and charged (IHTOA).

### **Method Name**

Immunohistochemistry (IHC)

### **NY State Available**

Yes

## Specimen

### **Specimen Type**

**TECHONLY** 

## **Ordering Guidance**

This test includes only technical performance of the stain (no pathologist interpretation is performed). If diagnostic consultation by a pathologist is required order PATHC / Pathology Consultation.

### **Shipping Instructions**

Attach the green "Attention Pathology" address label (T498) and the pink Immunostain Technical Only label included in the kit to the outside of the transport container.

## **Specimen Required**

**Specimen Type:** Tissue

**Supplies:** Immunostain Technical Only Envelope (T693) **Container/Tube:** Immunostain Technical Only Envelope

**Preferred:** 

-Formalin-fixed, paraffin-embedded tissue block



IgM Immunostain, Technical Component Only

OR

-2 Unstained, positively charged glass slides (25- x 75- x 1-mm) per test ordered; sections 4-microns thick **Acceptable**: None

## **Digital Image Access**

- 1. Information on accessing digital images of immunohistochemical (IHC) stains and the manual requisition form can be accessed through this website: https://news.mayocliniclabs.com/pathology/digital-imaging/
- 2. Clients ordering stains using a manual requisition form will not have access to digital images.
- 3. Clients wishing to access digital images must place the order for IHC stains electronically. Information regarding digital imaging can be accessed through this website: <a href="https://news.mayocliniclabs.com/pathology/digital-imaging/#section3">https://news.mayocliniclabs.com/pathology/digital-imaging/#section3</a>

#### **Forms**

If not ordering electronically, complete, print, and send a <u>Immunohistochemical (IHC)/In Situ Hybridization (ISH) Stains</u>
Request (T763) with the specimen.

## **Reject Due To**

| Wet/frozen     | Reject |
|----------------|--------|
| tissue         |        |
| Cytology       |        |
| smears         |        |
| Nonformalin    |        |
| fixed tissue   |        |
| Nonparaffin    |        |
| embedded       |        |
| tissue         |        |
| Noncharged     |        |
| slides         |        |
| ProbeOn slides |        |
| Snowcoat       |        |
| slides         |        |

## **Specimen Stability Information**

| Specimen Type | Temperature         | Time | Special Container |
|---------------|---------------------|------|-------------------|
| TECHONLY      | Ambient (preferred) |      |                   |
|               | Refrigerated        |      |                   |

## **Clinical & Interpretive**

## **Clinical Information**

The immunoglobulin molecules (antibodies) function as surface receptors for antigens for B lymphocytes and as secretory products of plasma cells forming the humoral arm of the immune system. IgM represents one of the immunoglobulin heavy chain types. IgM positive-B lymphocytes are normally present in the follicular mantle zones and



IgM Immunostain, Technical Component Only

germinal centers. Immunoreactivity is a specific marker for B lymphocytes and plasma cells. Diagnostically, staining for immunoglobulin heavy chains can aid in the diagnosis and classification of small B cell malignant lymphomas and multiple myeloma.

### Interpretation

This test does not include pathologist interpretation, only technical performance of the stain. If interpretation is required, order PATHC / Pathology Consultation for a full diagnostic evaluation or second opinion of the case.

The positive and negative controls are verified as showing appropriate immunoreactivity.

Interpretation of this test should be performed in the context of the patient's clinical history and other diagnostic tests by a qualified pathologist.

#### **Cautions**

Age of a cut paraffin section can affect immunoreactivity. Stability thresholds vary widely among published literature and are antigen dependent. Best practice is for paraffin sections to be cut within 6 weeks.

The charge of glass slides can be affected by environmental factors and subsequently may alter slide staining. Sending unsuitable glass slides can result in inconsistent staining due to poor slide surface chemistry.

Best practices for storage of positively charged slides:

- -Minimize time slides are stored after being unpackaged
- -Limit exposure to high humidity and heat
- -Minimize exposure to plastics

### **Clinical Reference**

- 1. Moreira RK, Revetta F, Koehler E, Washington MK. Diagnostic utility of IgG and IgM immunohistochemistry in autoimmune liver disease. World J Gastroenterol. 2010;16(4):453-457. doi:10.3748/wjg.v16.i4.453
- 2. Shubham S, Bhardwaj M, Mahapatra HS. Comparative evaluation of immunoperoxidase versus immunofluorescent techniques in interpretation of kidney biopsies. Indian J Pathol Microbiol. 2016;59(3):305-309. doi:10.4103/0377-4929.188105
- 3. Zhu Z, Zhang M, Shao W, et al. Immunoglobulin M, a novel molecule of myocardial cells of mice. Int J Biochem Cell Biol. 2017;88:172-180. doi:10.1016/j.biocel.2017.04.003
- 4. Magaki S, Hojat SA, Wei B, So A, Yong WH. An introduction to the performance of immunohistochemistry. Methods Mol Biol. 2019;1897:289-298. doi:10.1007/978-1-4939-8935-5\_25

### **Performance**

### **Method Description**

Immunohistochemistry on sections of paraffin-embedded tissue. (Unpublished Mayo method)

### **PDF Report**

No



IgM Immunostain, Technical Component Only

### Day(s) Performed

Monday through Friday

### **Report Available**

1 to 3 days

### **Specimen Retention Time**

Until staining is complete.

### **Performing Laboratory Location**

Mayo Clinic Laboratories - Rochester Main Campus

### **Fees & Codes**

#### **Fees**

- Authorized users can sign in to <u>Test Prices</u> for detailed fee information.
- Clients without access to Test Prices can contact <u>Customer Service</u> 24 hours a day, seven days a week.
- Prospective clients should contact their account representative. For assistance, contact <u>Customer Service</u>.

### **Test Classification**

This test has been cleared, approved, or is exempt by the US Food and Drug Administration and is used per manufacturer's instructions. Performance characteristics were verified by Mayo Clinic in a manner consistent with CLIA requirements.

### **CPT Code Information**

88342-TC, primary 88341-TC, if additional IHC

### **LOINC®** Information

| Test ID | Test Order Name    | Order LOINC® Value   |
|---------|--------------------|----------------------|
| IGMI    | IgM IHC, Tech Only | Order only;no result |

| Result ID | Test Result Name   | Result LOINC® Value  |
|-----------|--------------------|----------------------|
| 70785     | IgM IHC, Tech Only | Bill only; no result |