

## Overview

### Useful For

Determining overexpression of HER2 protein of gastric and esophageal adenocarcinoma in formalin-fixed, paraffin-embedded tissue sections (no reflex to FISH testing)

### Method Name

Ventana Pathway Immunoperoxidase Stain with Manual Semi-Quantitative Immunohistochemistry

### NY State Available

Yes

## Specimen

### Specimen Type

Special

### Ordering Guidance

**If ordering for diagnostic purposes:** order PATHC / Pathology Consultation and then request the stains.

For specimens such as intestine, liver, colon, which do not contain metastatic adenocarcinoma from the stomach or esophagus, order H2BR / *HER2* Amplification, Miscellaneous Tumor, FISH, Tissue.

For breast cancer specimens, order HERBA / *HER2*, Breast, Quantitative Immunohistochemistry, Automated with *HER2* FISH Reflex or HERBN / *HER2*, Breast, Quantitative Immunohistochemistry, Automated, No Reflex.

### Shipping Instructions

Attach the green pathology address label included in the kit to the outside of the transport container.

Ship ambient.

### Necessary Information

**Include accompanying pathology report stating the final diagnosis.** If not available, a preliminary diagnosis is acceptable **only** if it refers to invasive or metastatic gastric or esophageal adenocarcinoma.

### Specimen Required

**Only formalin-fixed, paraffin-embedded (FFPE) gastric or esophageal adenocarcinoma specimens will be accepted.**

Fixation in 10% neutral-buffered formalin is preferred. The performance and quality of immunohistochemical stains in 10% neutral-buffered FFPE tissue depends on proper fixation. It is recommended (not required) for surgical specimens to be fixed between 18 and 24 hours and biopsy specimens between 6 and 8 hours.

# Test Definition: HERGN

HER2, Gastric/Esophageal, Semi-Quantitative  
Immunohistochemistry, Manual, No Reflex

**If being ordered for prognostic purposes:**

- Specimen Type:** Gastric or esophageal adenocarcinoma
- Supplies:** Pathology Packaging Kit (T554)
- Preferred:** Paraffin-embedded tissue block containing invasive gastric or esophageal adenocarcinoma tissue
- Additional Information:** Paraffin blocks will be returned with final report.

- Acceptable:** Slides
- Specimen Volume:** 5
- Collection Instructions:** 5 Unstained sections, containing gastric or esophageal adenocarcinoma, on charged slides cut at 4 microns less than 1 month ago.

**Forms**

If not ordering electronically, complete, print, and send an [Immunohistochemical \(IHC\)/In Situ Hybridization \(ISH\) Stains Request](#) (T763)

- Reject Due To**
- No specimen should be rejected.

**Specimen Stability Information**

Specimen Type	Temperature	Time	Special Container
Special	Ambient (preferred)		
	Refrigerated		

**Clinical & Interpretive**

**Clinical Information**

The *HER2* (official gene name *ERBB2*) proto-oncogene encodes a membrane receptor with tyrosine kinase activity and homology to the epidermal growth factor receptor.

Amplification and overexpression of the *HER2* gene have been associated with a shorter disease-free survival and shorter overall survival in gastric and gastroesophageal junction cancers, as well as breast, endometrial, and ovarian cancer.(1,2)

**Reference Values**

Reported as negative (0, 1+), equivocal (2+), and positive (3+)

**Interpretation**

Results are reported as positive (3+ HER2 protein expression), equivocal (2+), or negative (0 or 1+)

**Cautions**

No significant cautionary statements

---

**Clinical Reference**

1. Pergam M, Slamon D: Biological rationale for *HER2/neu (c-erbB2)* as a target for monoclonal therapy. *Semin Oncol* 2000;27(5):13-19
2. Gravalos C, Jimeno A: HER2 in gastric cancer: a new prognostic factor and a novel therapeutic target. *Ann Oncol* 2008 Sep;19(9):1523-1529
3. Meza-Junco J, Au HJ, Sawyer MB: Trastuzumab for gastric cancer. *Expert Opin Biol Ther* 2009;9(12):1543-1551

**Performance****Method Description**

Testing is performed on formalin-fixed paraffin-embedded tissue sections using Ventana Pathway Immunoperoxidase HER2 (4B5) rabbit monoclonal primary antibody and a proprietary detection system. No expression (HER2 score of 0), low expression (HER2 score of 1+) and high expression (HER2 score of 3+) controls are used. (Package insert: PATHWAY anti-HER-2/neu [4B5] Rabbit Monoclonal Primary Antibody; Ventana Medical Systems Inc 3/16/2012)

**Scoring:**

Scoring is performed for surgical and biopsy specimens according to the following article: Ruschoff J, Dietel M, Baretton G, et al: HER2 diagnostics in gastric cancer-guideline validation and development of standardized immunohistochemical testing. *Virchows Arch* 2010 Sep;457(3):299-307

**Surgical Specimen:**

Score of 0 is no reactivity or membranous reactivity (staining) in <10% of invasive tumor cells. Score of 1+ is faint/barely perceptible membranous reactivity (staining) in > or =10% of invasive tumor cells; cells are reactive (stained) only in part of their membrane. Score of 2+ is weak to moderate complete, basolateral, or lateral membranous reactivity (staining) in > or =10% of invasive tumor cells. Score of 3+ is strong complete, basolateral, or lateral membranous reactivity (staining) in > or =10% of invasive tumor cells.

**Biopsy Specimen:**

Score of 0 is no reactivity or no membranous reactivity (staining) in any invasive tumor cells. Score of 1+ is tumor cell cluster with a faint/barely perceptible membranous reactivity (staining) irrespective of percentage of invasive tumor cells stained. Score of 2+ is tumor cell cluster with a weak to moderate complete, basolateral, or lateral membranous reactivity (staining) irrespective of percentage of invasive tumor cells stained. Score of 3+ is tumor cell cluster with a strong complete, basolateral, or lateral membranous reactivity (staining) irrespective of percentage of invasive tumor cells stained.

\*Tumor cells cluster is defined as a cluster of 5 or more tumor cells by Ruschoff and colleagues (2010). There is no percentage cutoff in biopsy specimens for upper GI tract HER2 scoring.

**PDF Report**

No

**Day(s) Performed**

Monday through Friday

Report Available

4 to 6 days

Specimen Retention Time

Until 1 week after results are reported. Materials made at Mayo Clinic may be retained at Mayo Clinic indefinitely.

Performing Laboratory Location

Mayo Clinic Laboratories - Rochester Main Campus

Fees & Codes

Fees

- Authorized users can sign in to [Test Prices](#) for detailed fee information.
- Clients without access to Test Prices can contact [Customer Service](#) 24 hours a day, seven days a week.
- Prospective clients should contact their account representative. For assistance, contact [Customer Service](#).

Test Classification

This test has been modified from the manufacturer's instructions. Its performance characteristics were determined by Mayo Clinic in a manner consistent with CLIA requirements. This test has not been cleared or approved by the US Food and Drug Administration.

CPT Code Information

88360

LOINC® Information

Test ID	Test Order Name	Order LOINC® Value
HERGN	HER Gastric/Esoph IHC NO Reflex	Obsolete

Result ID	Test Result Name	Result LOINC® Value
MA021	Tumor classification	21918-8
70990	Interpretation	50595-8
70991	Participated in the Interpretation	No LOINC Needed
70992	Report electronically signed by	19139-5
70994	Material Received	81178-6
71626	Disclaimer	62364-5
71840	Case Number	80398-1