

Test Definition: FMARC

Bupivacaine (Marcaine)

Overview

Method Name

Liquid Chromatography/ Tandem Mass Spectrometry (LC/MS/MS)

NY State Available

Yes

Specimen

Specimen Type

Serum Red

Specimen Required

Draw blood in a plain red-top tube(s), **serum gel tube is not acceptable**. Spin down and send 1 mL of serum frozen in a plastic vial.

Specimen Minimum Volume

0.5 mL

Reject Due To

Hemolysis	NA NA
Lipemia	NA
Icterus	NA
Other	NA

Specimen Stability Information

Specimen Type	Temperature	Time	Special Container
Serum Red	Frozen (preferred)	180 days	
	Ambient	72 hours	
	Refrigerated	72 hours	

Clinical & Interpretive

Reference Values

Reference Range: 0.1 - 4.0 ug/mL



Test Definition: FMARC

Bupivacaine (Marcaine)

Performance

PDF Report

No

Day(s) Performed

Monday through Sunday

Report Available

5 to 11 days

Performing Laboratory Location

Medtox Laboratories, Inc.

Fees & Codes

Fees

- Authorized users can sign in to <u>Test Prices</u> for detailed fee information.
- Clients without access to Test Prices can contact <u>Customer Service</u> 24 hours a day, seven days a week.
- Prospective clients should contact their account representative. For assistance, contact <u>Customer Service</u>.

Test Classification

This test was developed and its performance characteristics determined by LabCorp. It has not been cleared or approved by the Food and Drug Administration.

CPT Code Information

80299

LOINC® Information

Test ID	Test Order Name	Order LOINC® Value
FMARC	Bupivacaine (Marcaine)	3412-4

Result ID	Test Result Name	Result LOINC® Value
Z5052	Bupivacaine	3412-4