



# Test Definition: FMARC

Bupivacaine (Marcaine)

## Overview

### Method Name

Liquid Chromatography/ Tandem Mass Spectrometry (LC/MS/MS)

### NY State Available

Yes

## Specimen

### Specimen Type

Serum Red

### Specimen Required

Draw blood in a plain red-top tube(s), **serum gel tube is not acceptable**. Spin down and send 1 mL of serum frozen in a plastic vial.

### Specimen Minimum Volume

0.5 mL

### Reject Due To

Hemolysis	NA
Lipemia	NA
Icterus	NA
Other	NA

### Specimen Stability Information

Specimen Type	Temperature	Time	Special Container
Serum Red	Frozen (preferred)	180 days	
	Ambient	72 hours	
	Refrigerated	72 hours	

## Clinical & Interpretive

### Reference Values

Reference Range: 0.1 - 4.0 ug/mL

**Performance****PDF Report**

No

**Day(s) Performed**

Monday through Sunday

**Report Available**

5 to 11 days

**Performing Laboratory Location**

Medtox Laboratories, Inc.

**Fees & Codes****Fees**

- Authorized users can sign in to [Test Prices](#) for detailed fee information.
- Clients without access to Test Prices can contact [Customer Service](#) 24 hours a day, seven days a week.
- Prospective clients should contact their account representative. For assistance, contact [Customer Service](#).

**Test Classification**

This test was developed and its performance characteristics determined by LabCorp. It has not been cleared or approved by the Food and Drug Administration.

**CPT Code Information**

80299

**LOINC® Information**

Test ID	Test Order Name	Order LOINC® Value
FMARC	Bupivacaine (Marcaine)	3412-4

Result ID	Test Result Name	Result LOINC® Value
Z5052	Bupivacaine	3412-4